



Louisiana

Outpatient Pulmonary Rehabilitation

Policy # 00621

Original Effective Date: 10/01/2018

Current Effective Date: 08/09/2021

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Note: Lung and Lobar Lung Transplant is addressed separately in medical policy 00414.

When Services Are Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Based on review of available data, the Company may consider a single course of pulmonary rehabilitation (PR) in the outpatient ambulatory care setting for treatment of chronic pulmonary disease (e.g., moderate-to-severe chronic obstructive pulmonary disease (COPD), bronchiectasis, cystic fibrosis, interstitial lung disease) for patients with moderate-to-severe disease who are experiencing disabling symptoms and significantly diminished quality of life despite optimal medical management to be **eligible for coverage.****

Based on review of available data, the Company may consider a single course of PR in an outpatient ambulatory care setting as a preoperative conditioning component for those considered appropriate candidates for lung volume reduction surgery (LVRS) or for lung transplantation (see medical policy 00414) to be **eligible for coverage.****

Based on review of available data, the Company may consider PR programs following lung transplantation to be **eligible for coverage.****

Note: When a single course of outpatient PR is considered medically necessary, a maximum of 3 sessions per week for 12 weeks will be approved.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Outpatient Pulmonary Rehabilitation

Policy # 00621

Original Effective Date: 10/01/2018

Current Effective Date: 08/09/2021

When Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Based on review of available data, the Company considers PR programs following other types of lung surgery, included but not limited to LVRS and surgical resection of lung cancer to be **investigational.***

Based on review of available data, the Company considers multiple courses of PR, either as maintenance therapy in patients who initially respond, or in patients who fail to respond, or whose response to an initial rehabilitation program has diminished over time to be **investigational.***

Based on review of available data, the Company considers home-based PR programs to be **investigational.***

Based on review of available data, the Company considers PR programs in all other situations to be **investigational.***

Policy Guidelines

A pulmonary rehabilitation outpatient program is a comprehensive program that generally includes team assessment, patient training, psychosocial intervention, exercise training, and follow-up. The overall length of the program and the total number of visits for each component may vary from program to program.

Team assessment includes input from a physician, respiratory care practitioner, nurse, and psychologist, among others.

Patient training includes breathing retraining, bronchial hygiene, medications, and proper nutrition.

Psychosocial intervention addresses support system and dependency issues.

Exercise training includes strengthening and conditioning, and may include stair climbing, inspiratory muscle training, treadmill walking, cycle training (with or without ergometer), and

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Outpatient Pulmonary Rehabilitation

Policy # 00621

Original Effective Date: 10/01/2018

Current Effective Date: 08/09/2021

supported and unsupported arm exercise training. Exercise conditioning is an essential component of pulmonary rehabilitation. Education in disease management techniques without exercise conditioning does not improve health outcomes of patients who have chronic obstructive pulmonary disease.

Follow-up to a comprehensive outpatient pulmonary rehabilitation program may include supervised home exercise conditioning.

Candidates for pulmonary rehabilitation should be medically stable and not limited by another serious or unstable medical condition. Contraindications to pulmonary rehabilitation include severe psychiatric disturbance (eg, dementia, organic brain syndrome) and significant or unstable medical conditions (eg, heart failure, acute cor pulmonale, substance abuse, significant liver dysfunction, metastatic cancer, disabling stroke).

Background/Overview

In 2013, the American Thoracic Society and the European Respiratory Society defined pulmonary rehabilitation as a “comprehensive intervention based on a thorough patient assessment followed by patient-tailored therapies that include, but are not limited to exercise training, education, and behavior change.” Pulmonary rehabilitation programs are intended to improve patient functioning and quality of life. Most research has focused on patients with chronic obstructive pulmonary disease, although there has been some interest in patients with asthma, cystic fibrosis, or bronchiectasis.

Pulmonary rehabilitation is also routinely offered to patients awaiting lung transplantation and lung volume reduction surgery. Pulmonary rehabilitation before lung surgery may stabilize or improve patients’ exercise tolerance, teach patients techniques that will help them recover after the procedure, and allow health care providers to identify individuals who might be suboptimal surgical candidates due to noncompliance, poor health, or other reasons.’

Rationale/Source

Pulmonary rehabilitation is a multidisciplinary approach to reducing symptoms and improving quality of life in patients with compromised lung function. Pulmonary rehabilitation programs

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Outpatient Pulmonary Rehabilitation

Policy # 00621

Original Effective Date: 10/01/2018

Current Effective Date: 08/09/2021

generally include a patient assessment followed by therapeutic interventions including exercise training, education, and behavior change.

Summary of Evidence

Chronic Pulmonary Disease Rehabilitation

For individuals with moderate-to-severe chronic obstructive pulmonary disease (COPD) who receive a single course of outpatient pulmonary rehabilitation, the evidence includes numerous systematic reviews of randomized controlled trials (RCTs). Relevant outcomes are symptoms, functional outcomes, and quality of life. The published studies found improved outcomes (ie, functional ability, quality of life) in patients with moderate-to-severe COPD who underwent a comprehensive pulmonary rehabilitation program in the outpatient setting. Among the many randomized trials, the structure of the pulmonary rehabilitation programs varied, so it is not possible to provide guidance on the optimal components or duration of a pulmonary rehabilitation program. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with idiopathic pulmonary fibrosis who receive a single course of outpatient pulmonary rehabilitation, the evidence includes 2 systematic reviews of RCTs. Relevant outcomes are symptoms, functional outcomes, and quality of life. The number of controlled studies is limited. One small RCT evaluated a comprehensive pulmonary rehabilitation program in patients with idiopathic pulmonary fibrosis; at 3 months postintervention, outcomes did not differ between groups that did and did not receive pulmonary rehabilitation. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with bronchiectasis who receive a single course of outpatient pulmonary rehabilitation, the evidence includes a systematic review of RCTs. Relevant outcomes are symptoms, functional outcomes, and quality of life. The systematic review of 4 RCTs on pulmonary rehabilitation for patients with bronchiectasis found that some, but not all, outcomes, improved more with pulmonary rehabilitation than with nonexercise control conditions immediately after the intervention. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Outpatient Pulmonary Rehabilitation

Policy # 00621

Original Effective Date: 10/01/2018

Current Effective Date: 08/09/2021

Although most published evidence on outpatient pulmonary rehabilitation for chronic pulmonary diseases assesses COPD, observational studies have reported on outcomes from pulmonary rehabilitation for other chronic pulmonary diseases. Clinical guidelines from pulmonary organizations have supported the use of outpatient pulmonary rehabilitation for individuals who are experiencing disabling symptoms and have significantly diminished quality of life despite optimal medical management. Therefore, outpatient pulmonary rehabilitation may be considered medically necessary for this population.

Preparation for Lung Surgery

For individuals with scheduled lung surgery for volume reduction, transplantation, or resection who receive a single course of outpatient pulmonary rehabilitation, the evidence includes RCTs and observational studies. Relevant outcomes are symptoms, functional outcomes, and quality of life. There is a lack of large RCTs comparing pulmonary rehabilitation with no pulmonary rehabilitation for preoperative candidates undergoing lung volume reduction surgery, lung transplantation, or lung cancer resection. Moreover, the available studies have evaluated exercise programs, but not necessarily comprehensive pulmonary rehabilitation programs. Also, the few small RCTs, and observational studies have only reported short-term outcomes and inconsistent evidence of benefit even on these outcomes. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Findings from the National Emphysema Treatment Trial have suggested that pulmonary rehabilitation is an appropriate component of care for patients with COPD before undergoing lung volume reduction surgery. Also, pulmonary rehabilitation is considered standard of care in patients undergoing lung transplantation to maximize preoperative pulmonary status. Thus, pulmonary rehabilitation may be considered medically necessary for patients considered appropriate candidates for lung volume reduction surgery or lung transplantation.

Pulmonary Rehabilitation After Lung Surgery

For individuals who have had lung volume reduction surgery who receive a single course of outpatient pulmonary rehabilitation, the evidence includes a case series. Relevant outcomes are symptoms, functional outcomes, and quality of life. No published RCTs were identified. The case series evaluated a comprehensive pulmonary rehabilitation program after lung volume reduction surgery in 49 patients who had not received preoperative pulmonary rehabilitation. Health-related quality of life was higher at 3 to 6 months and 12 to 18 months postsurgery. The series did not

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Outpatient Pulmonary Rehabilitation

Policy # 00621

Original Effective Date: 10/01/2018

Current Effective Date: 08/09/2021

provide data on patients who underwent lung volume reduction surgery and did not have postoperative pulmonary rehabilitation, or patients who had preoperative pulmonary rehabilitation. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have had lung transplantation who receive a single course of outpatient pulmonary rehabilitation, the evidence includes RCTs, a systematic review, and a case series. Relevant outcomes are symptoms, functional outcomes, and quality of life. Neither of the 2 RCTs identified in a 2010 systematic review reported on functional outcomes, but uncontrolled studies have reported improvements in functional outcomes. An RCT, published after the systematic review, found that patients who had a postsurgical exercise intervention walked more 1 year post discharge than before and had a significantly greater 6-minute walk distance. Findings on other outcomes were mixed. The most recent RCT (2017) did not identify a difference in outcomes with longer duration of pulmonary rehabilitation. Case series data also support improvements in 6-minute walk distance after postoperative pulmonary rehabilitation. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have had lung cancer resection who receive a single course of outpatient pulmonary rehabilitation, the evidence includes 2 RCTs. Relevant outcomes are symptoms, functional outcomes, and quality of life. One small RCT evaluated a comprehensive pulmonary rehabilitation program in patients who underwent thoracotomy for lung cancer. The trial was terminated early, had a high dropout rate, and reported mixed findings. An exercise-only intervention in patients who had lung cancer surgery had mixed findings and did not evaluate functional outcomes. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Repeat or Maintenance Pulmonary Rehabilitation

For individuals who have had an initial course of pulmonary rehabilitation who receive repeat or maintenance outpatient pulmonary rehabilitation, the evidence includes an RCT. Relevant outcomes are symptoms, functional outcomes, and quality of life. This small RCT had methodologic limitations and did not report inpatient and outpatient outcomes separately; it also lasted only 3 weeks. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Outpatient Pulmonary Rehabilitation

Policy # 00621

Original Effective Date: 10/01/2018

Current Effective Date: 08/09/2021

Home-Based Pulmonary Rehabilitation

For individuals who have an indication for outpatient pulmonary rehabilitation who receive a single course of home-based pulmonary rehabilitation, the evidence includes RCTs and systematic reviews. Relevant outcomes are symptoms, functional outcomes, and quality of life. Most studies of home-based pulmonary rehabilitation have compared outcomes with standard care. Very few have compared home-based pulmonary rehabilitation with the hospital- or clinic-based pulmonary rehabilitation, and the available studies are mostly of low quality. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Supplemental Information

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in ‘Supplemental Information’ if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American Thoracic Society and European Respiratory Society

A 2015 joint statement on pulmonary rehabilitation was issued by the American Thoracic Society and the European Respiratory Society. The statement included the following relevant conclusions:

- “Pulmonary rehabilitation (PR) has demonstrated physiological, symptom-reducing, psychosocial, and health economic benefits in multiple outcome areas for patients with chronic respiratory diseases.”
- “The evidence indicates that patients who benefit from PR include not only persons with moderate to severe airflow limitation but also those with mild to moderate airflow limitation with symptom-limited exercise tolerance, those after hospitalization for COPD [chronic obstructive pulmonary disease] exacerbation, and those with symptomatic non-COPD respiratory conditions.”
- “Patients graduating from a PR program stand to benefit from a home, community-based, or program-based maintenance exercise program to support the continuation of positive exercise behavior.”

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Outpatient Pulmonary Rehabilitation

Policy # 00621

Original Effective Date: 10/01/2018

Current Effective Date: 08/09/2021

In 2017, the Society issued a joint statement on the management of COPD exacerbation. For patients hospitalized with a COPD exacerbation, they suggest “the initiation of pulmonary rehabilitation within 3 weeks after hospital discharge” (strength: conditional; quality of evidence: very low). In addition, “[they] suggest not initiating pulmonary rehabilitation during hospitalisation” (strength: conditional; quality of evidence: very low).

American College of Physicians

In 2011, Joint guidelines on the management of stable COPD were issued by the American College of Physicians, the American College of Chest Physicians, American Thoracic Society, and European Respiratory Society. The guidelines recommended that “clinicians should prescribe pulmonary rehabilitation for symptomatic patients with an FEV [forced expiratory volume] <50% predicted (Grade: strong recommendation, moderate-quality evidence). Clinicians may consider pulmonary rehabilitation for symptomatic or exercise-limited patients with an FEV >50% predicted (Grade: weak recommendation, moderate-quality evidence).”

United States Preventive Services Task Force Recommendations

Not applicable

Medicare National Coverage

In 2007, the Centers for Medicare & Medicaid Services affirmed its position that a national coverage determination for pulmonary rehabilitation is not appropriate.

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 1.

Table 1. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
NCT03326089	Short and Long-term Effects of Oxygen Supplemented Pulmonary Rehabilitation in Idiopathic Pulmonary Fibrosis	20	Sep 2022
<i>Unpublished</i>			

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Outpatient Pulmonary Rehabilitation

Policy # 00621

Original Effective Date: 10/01/2018

Current Effective Date: 08/09/2021

NCT03299504	Factors Predicting Success in Lung Transplant Recipients Who Have Participated in the COLTT Program (Daily Intensive Post-hospitalization Rehabilitation): A Retrospective Review	105	Apr 2018 (last updated 08/24/18)
NCT03244137	Effects of Pulmonary Rehabilitation on Cognitive Function in Patients With Severe to Very Severe Chronic Obstructive Pulmonary Disease	100	Dec 2019 (last updated 01/07/20)
NCT02426437	Examining Pulmonary Rehabilitation on Discharged COPD Patients	150	Dec 2020
NCT02842463	Use of the 6-minute Stepper Test to Individualise Pulmonary Rehabilitation in Patients With Mild to Moderate Chronic Obstructive Pulmonary Disease	80	June 2020
NCT03244137	Effects of Pulmonary Rehabilitation on Cognitive Function in Patients With Severe to Very Severe Chronic Obstructive Pulmonary Disease	100	Dec 2019 (completed)
NCT02887521	Pulmonary Rehabilitation Before Lung Cancer Resection	194	Oct 2019 (completed)

NCT: national clinical trial.

References

1. Blue Cross and Blue Shield Association, Medical Policy Reference Manual, “Outpatient Pulmonary Rehabilitation”, 8.03.05, April 2021.
2. Spruit MA, Singh SJ, Garvey C, et al. An official American Thoracic Society/European Respiratory Society statement: key concepts and advances in pulmonary rehabilitation. *Am J Respir Crit Care Med.* Oct 15 2013; 188(8): e13-64. PMID 24127811
3. Puhan MA, Gimeno-Santos E, Cates CJ, et al. Pulmonary rehabilitation following exacerbations of chronic obstructive pulmonary disease. *Cochrane Database Syst Rev.* Dec 08 2016; 12: CD005305. PMID 27930803
4. McCarthy B, Casey D, Devane D, et al. Pulmonary rehabilitation for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev.* Feb 23 2015; (2): CD003793. PMID 25705944

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Outpatient Pulmonary Rehabilitation

Policy # 00621

Original Effective Date: 10/01/2018

Current Effective Date: 08/09/2021

5. Rugbjerg M, Iepsen UW, Jorgensen KJ, et al. Effectiveness of pulmonary rehabilitation in COPD with mild symptoms: a systematic review with meta-analyses. *Int J Chron Obstruct Pulmon Dis.* 2015; 10: 791-801. PMID 25945044
6. Roman M, Larraz C, Gomez A, et al. Efficacy of pulmonary rehabilitation in patients with moderate chronic obstructive pulmonary disease: a randomized controlled trial. *BMC Fam Pract.* Feb 11 2013; 14: 21. PMID 23399113
7. Gottlieb V, Lyngso AM, Nybo B, et al. Pulmonary rehabilitation for moderate COPD (GOLD 2)--does it have an effect?. *COPD.* Oct 2011; 8(5): 380-6. PMID 21936683
8. Liu X-D, Jin H-Z, Ng B-P, et al. Therapeutic effects of qigong in patients with COPD: a randomized controlled trial. *Hong Kong J Occup Ther.* Aug 9 2012;22(1):38-46. <https://journals.sagepub.com/doi/pdf/10.1016/j.hkjot.2012.06.002>.
9. van Wetering CR, Hoogendoorn M, Mol SJ, et al. Short- and long-term efficacy of a community-based COPD management programme in less advanced COPD: a randomised controlled trial. *Thorax.* Jan 2010; 65(1): 7-13. PMID 19703824
10. Yu X, Li X, Wang L, et al. Pulmonary Rehabilitation for Exercise Tolerance and Quality of Life in IPF Patients: A Systematic Review and Meta-Analysis. *Biomed Res Int.* 2019; 2019: 8498603. PMID 31016200
11. Cheng L, Tan B, Yin Y, et al. Short- and long-term effects of pulmonary rehabilitation for idiopathic pulmonary fibrosis: a systematic review and meta-analysis. *Clin Rehabil.* Oct 2018; 32(10): 1299-1307. PMID 29843523
12. Lee AL, Hill CJ, McDonald CF, et al. Pulmonary Rehabilitation in Individuals With Non-Cystic Fibrosis Bronchiectasis: A Systematic Review. *Arch Phys Med Rehabil.* Apr 2017; 98(4): 774-782.e1. PMID 27320420
13. Fishman A, Martinez F, Naunheim K, et al. A randomized trial comparing lung-volume-reduction surgery with medical therapy for severe emphysema. *N Engl J Med.* May 22 2003; 348(21): 2059-73. PMID 12759479
14. Hoffman M, Chaves G, Ribeiro-Samora GA, et al. Effects of pulmonary rehabilitation in lung transplant candidates: a systematic review. *BMJ Open.* Feb 03 2017; 7(2): e013445. PMID 28159852
15. Morano MT, Araujo AS, Nascimento FB, et al. Preoperative pulmonary rehabilitation versus chest physical therapy in patients undergoing lung cancer resection: a pilot randomized controlled trial. *Arch Phys Med Rehabil.* Jan 2013; 94(1): 53-8. PMID 22926460

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Outpatient Pulmonary Rehabilitation

Policy # 00621

Original Effective Date: 10/01/2018

Current Effective Date: 08/09/2021

16. Benzo R, Wigle D, Novotny P, et al. Preoperative pulmonary rehabilitation before lung cancer resection: results from two randomized studies. *Lung Cancer*. Dec 2011; 74(3): 441-5. PMID 21663994
17. Bradley A, Marshall A, Stonehewer L, et al. Pulmonary rehabilitation programme for patients undergoing curative lung cancer surgery. *Eur J Cardiothorac Surg*. Oct 2013; 44(4): e266-71. PMID 23959742
18. Beling J. Improved health-related quality of life after lung volume reduction surgery and pulmonary rehabilitation. *Cardiopulm Phys Ther J*. Sep 2009; 20(3): 16-22. PMID 20467519
19. Wickerson L, Mathur S, Brooks D. Exercise training after lung transplantation: a systematic review. *J Heart Lung Transplant*. May 2010; 29(5): 497-503. PMID 20133160
20. Langer D, Burtin C, Schepers L, et al. Exercise training after lung transplantation improves participation in daily activity: a randomized controlled trial. *Am J Transplant*. Jun 2012; 12(6): 1584-92. PMID 22390625
21. Fuller LM, Button B, Tarrant B, et al. Longer Versus Shorter Duration of Supervised Rehabilitation After Lung Transplantation: A Randomized Trial. *Arch Phys Med Rehabil*. Feb 2017; 98(2): 220-226.e3. PMID 27697429
22. Munro PE, Holland AE, Bailey M, et al. Pulmonary rehabilitation following lung transplantation. *Transplant Proc*. Jan-Feb 2009; 41(1): 292-5. PMID 19249538
23. Stigt JA, Uil SM, van Riesen SJ, et al. A randomized controlled trial of postthoracotomy pulmonary rehabilitation in patients with resectable lung cancer. *J Thorac Oncol*. Feb 2013; 8(2): 214-21. PMID 23238118
24. Edvardsen E, Skjonsberg OH, Holme I, et al. High-intensity training following lung cancer surgery: a randomised controlled trial. *Thorax*. Mar 2015; 70(3): 244-50. PMID 25323620
25. Carr SJ, Hill K, Brooks D, et al. Pulmonary rehabilitation after acute exacerbation of chronic obstructive pulmonary disease in patients who previously completed a pulmonary rehabilitation program. *J Cardiopulm Rehabil Prev*. Sep-Oct 2009; 29(5): 318-24. PMID 19561523
26. COPD Working Group. Pulmonary rehabilitation for patients with chronic pulmonary disease (COPD): an evidence-based analysis. *Ont Health Technol Assess Ser*. 2012; 12(6): 1-75. PMID 23074434
27. Guell MR, Cejudo P, Ortega F, et al. Benefits of Long-Term Pulmonary Rehabilitation Maintenance Program in Patients with Severe Chronic Obstructive Pulmonary Disease. Three-Year Follow-up. *Am J Respir Crit Care Med*. Mar 01 2017; 195(5): 622-629. PMID 27611807

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Outpatient Pulmonary Rehabilitation

Policy # 00621

Original Effective Date: 10/01/2018

Current Effective Date: 08/09/2021

28. Wilson AM, Browne P, Olive S, et al. The effects of maintenance schedules following pulmonary rehabilitation in patients with chronic obstructive pulmonary disease: a randomised controlled trial. *BMJ Open*. Mar 11 2015; 5(3): e005921. PMID 25762226
29. Liu XL, Tan JY, Wang T, et al. Effectiveness of home-based pulmonary rehabilitation for patients with chronic obstructive pulmonary disease: a meta-analysis of randomized controlled trials. *Rehabil Nurs*. Jan-Feb 2014; 39(1): 36-59. PMID 23780865
30. Vieira DS, Maltais F, Bourbeau J. Home-based pulmonary rehabilitation in chronic obstructive pulmonary disease patients. *Curr Opin Pulm Med*. Mar 2010; 16(2): 134-43. PMID 20104176
31. Neves LF, Reis MH, Goncalves TR. Home or community-based pulmonary rehabilitation for individuals with chronic obstructive pulmonary disease: a systematic review and meta-analysis. *Cad Saude Publica*. Jun 20 2016; 32(6). PMID 27333130
32. Maltais F, Bourbeau J, Shapiro S, et al. Effects of home-based pulmonary rehabilitation in patients with chronic obstructive pulmonary disease: a randomized trial. *Ann Intern Med*. Dec 16 2008; 149(12): 869-78. PMID 19075206
33. Rochester CL, Vogiatzis I, Holland AE, et al. An Official American Thoracic Society/European Respiratory Society Policy Statement: Enhancing Implementation, Use, and Delivery of Pulmonary Rehabilitation. *Am J Respir Crit Care Med*. Dec 01 2015; 192(11): 1373-86. PMID 26623686
34. Wedzicha JA, Miravittles M, Hurst JR, et al. Management of COPD exacerbations: a European Respiratory Society/American Thoracic Society guideline. *Eur Respir J*. Mar 2017; 49(3). PMID 28298398
35. Qaseem A, Wilt TJ, Weinberger SE, et al. Diagnosis and management of stable chronic obstructive pulmonary disease: a clinical practice guideline update from the American College of Physicians, American College of Chest Physicians, American Thoracic Society, and European Respiratory Society. *Ann Intern Med*. Aug 02 2011; 155(3): 179-91. PMID 21810710
36. Centers for Medicare & Medicaid Services. National Coverage Determination (NCD) for Pulmonary Rehabilitation Services (240.8). 2008; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCAId=130&NcaName=Smoking+%26+Tobacco+Use+Cessation+Counseling&ExpandComments=n&CommentPeriod=0&NCDId=320&NCSelection=NCA%7CCAL%7CNC%7CMEDCAC%7CTA%7CMCD&KeyWord=Pulmonary+Rehabilitation&KeyWordLookUp=Doc&KeyWordSearchType=And&kq=true>.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Outpatient Pulmonary Rehabilitation

Policy # 00621

Original Effective Date: 10/01/2018

Current Effective Date: 08/09/2021

Policy History

Original Effective Date: 10/01/2018

Current Effective Date: 08/09/2021

07/05/2018 Medical Policy Committee review

07/11/2018 Medical Policy Implementation Committee approval. New policy.

07/03/2019 Medical Policy Committee review

07/18/2019 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

07/02/2020 Medical Policy Committee review

07/08/2020 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

09/16/2020 Coding update

07/01/2021 Medical Policy Committee review

07/14/2021 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

Next Scheduled Review Date: 07/2022

Coding

The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)†, copyright 2020 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Outpatient Pulmonary Rehabilitation

Policy # 00621

Original Effective Date: 10/01/2018

Current Effective Date: 08/09/2021

Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	No codes
HCPCS	G0237, G0238, G0239, G0302, G0303, G0304, G0305, G0424, S9473
ICD-10 Diagnosis	C96.6, D38.0-D38.6, D84.1, D86.0-D86.9, E84.0-E84.9, I11.0-I11.9, I27.0, J41.0-J41.8, J43.0-J43.9, J44.0-J44.9, J47.0-J47.9, J84.0-J84.9, J99, M34.0-M34.9, P27.0-P27.9, Q21.0, Z86.711 Added codes eff 10/1/2020: J84.170, J84.178

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
 1. Consultation with the Blue Cross and Blue Shield Association technology assessment program (TEC) or other nonaffiliated technology evaluation center(s);
 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Outpatient Pulmonary Rehabilitation

Policy # 00621

Original Effective Date: 10/01/2018

Current Effective Date: 08/09/2021

3. Reference to federal regulations.

****Medically Necessary (or “Medical Necessity”)** - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. In accordance with nationally accepted standards of medical practice;
- B. Clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and
- C. Not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.

For these purposes, “nationally accepted standards of medical practice” means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

‡ Indicated trademarks are the registered trademarks of their respective owners.

NOTICE: If the Patient’s health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.