



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Note: Hematopoietic Cell Transplantation for Primary Amyloidosis or Waldenstrom's Macroglobulinemia is addressed separately in medical policy 00138.

When Services Are Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Multiple Myeloma

When Services Are Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Based on review of available data, the Company may consider a single or second (salvage) autologous hematopoietic cell transplantation (HCT) to treat multiple myeloma to be **eligible for coverage**.**

Based on review of available data, the Company may consider tandem** autologous-autologous hematopoietic cell transplantation (HCT) to treat multiple myeloma be **eligible for coverage**.**

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

****Tandem transplantation refers to a planned infusion (transplant) of previously harvested hematopoietic stem cells with a repeat hematopoietic cell infusion (transplant) that is performed within 6 months of the initial transplant. This is distinguished from a repeat transplantation requested or performed more than 6 months after the first transplant, and is used as salvage therapy after failure of initial transplantation or relapsed disease.**

Based on review of available data, the Company may consider tandem transplantation with an initial round of autologous hematopoietic cell transplantation (HCT) followed by a non-marrow-ablative conditioning regimen and allogeneic hematopoietic cell transplantation (HCT) (i.e., reduced-intensity conditioning transplant) to treat newly diagnosed multiple myeloma patients to be **eligible for coverage.****

When Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Based on review of available data, the Company considers allogeneic hematopoietic cell transplantation (HCT), myeloablative or nonmyeloablative, as initial therapy of newly diagnosed multiple myeloma or as salvage therapy to be **investigational.***

POEMS Syndrome

When Services Are Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Based on review of available data, the Company may consider autologous hematopoietic cell transplantation (HCT) to treat disseminated POEMS syndrome to be **eligible for coverage.****

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

When Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Based on review of available data, the Company considers allogeneic and tandem hematopoietic cell transplantation (HCT) to treat POEMS syndrome to be **investigational**.*

Policy Guidelines

The International Working Group on Myeloma has updated the European Group for Blood and Marrow Transplant criteria to describe a complete response to multiple myeloma therapy. The criteria include negative immunofixation on the serum and urine; disappearance of soft tissue plasmacytomas; and 5% or fewer plasma cells in bone marrow aspiration.

Patients with disseminated POEMS syndrome may have diffuse sclerotic lesions or disseminated bone marrow involvement.

Background/Overview

Multiple Myeloma

Multiple myeloma (MM) is a systemic malignancy of plasma cells that represents approximately 10% of all hematologic cancers. It is treatable but rarely curable. At diagnosis, most patients have generalized disease, and the selection of treatment is influenced by patient age, general health, prior therapy, and the presence of disease complications.

The disease is staged by estimating tumor mass, based on various clinical parameters such as hemoglobin, serum calcium, number of lytic bone lesions, and the presence or absence of renal failure. Multiple myeloma usually evolves from an asymptomatic premalignant stage (termed *monoclonal gammopathy of undetermined significance*). Treatment is usually reserved for patients with symptomatic disease (usually progressive myeloma), whereas asymptomatic patients are observed because there is little evidence that early treatment of asymptomatic MM prolongs survival compared with therapy delivered at the time of symptoms or end-organ damage. In some patients, an intermediate asymptomatic but more advanced premalignant stage is recognized

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

and referred to as smoldering MM. The overall risk of disease progression from smoldering to symptomatic MM is 10% per year for the first 5 years, approximately 3% per year for the next 5 years, and 1% for the next 10 years.

POEMS Syndrome

POEMS syndrome (also known as osteosclerotic myeloma, Crow-Fukase syndrome, or Takatsuki syndrome) is a rare, paraneoplastic disorder secondary to a plasma cell dyscrasia. This complex, multiorgan disease was first described in 1938, but the acronym POEMS was coined in 1980, reflecting hallmark characteristics of the syndrome: polyneuropathy, organomegaly, endocrinopathy, M protein, and skin changes. No single test establishes the presence of POEMS syndrome. Its pathogenesis is undefined, although some evidence has suggested it is mediated by an imbalance of proinflammatory cytokines including interleukin (IL)-1 β , IL-6, and tumor necrosis factor α ; vascular endothelial growth factor may also be involved. However, specific criteria have been established, and the syndrome may entail other findings in the constellation of signs and symptoms, as shown in Table 1. Both mandatory major criteria, at least 1 of the other major criteria, and at least 1 of the minor criteria are necessary for diagnosis.

Table 1. Criteria and Associations for POEMS Syndrome

Mandatory Major Criteria	Other Major Criteria	Minor Criteria	Other Symptoms and Signs
Polyneuropathy	Castleman disease	Organomegaly (splenomegaly, hepatomegaly, lymphadenopathy)	Pulmonary hypertension/restrictive lung disease
Monoclonal plasma-proliferative disorder	Sclerotic bone lesions	Edema (edema, pleural effusion, ascites)	Clubbing
	Vascular endothelial	Endocrinopathy (adrenal, thyroid, pituitary,	Thrombotic diatheses

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

Mandatory Major Criteria	Other Major Criteria	Minor Criteria	Other Symptoms and Signs
	growth factor elevation	gonadal, parathyroid, pancreatic)	
		Skin changes (hyperpigmentation, hypertrichosis, plethora, hemangiomas, white nails)	Weight loss
		Papilledema	Low vitamin B ₁₂ levels
		Thrombosis/polycythemia	Diarrhea
			Hyperhidrosis

The prevalence of POEMS syndrome is unclear. A national survey in Japan showed a prevalence of about 0.3 per 100,000. Other large series have been described in the United States, France, China, and India. In general, patients with POEMS have superior overall survival compared with that of MM (nearly 14 years in a large series). However, given the rarity of POEMS, there is a paucity of RCT evidence for POEMS therapies. Numerous approaches have been tried, including ionizing radiation, plasmapheresis, intravenous immunoglobulin, interferon- α , corticosteroids, alkylating agents, tamoxifen, trans-retinoic acid, and high-dose chemotherapy with autologous hematopoietic cell transplantation (HCT) support. Optimal treatment involves eliminating the plasma cell clone (eg, by surgical excision or local radiotherapy for an isolated plasmacytoma) or systemic chemotherapy in patients with disseminated disease (eg, medullary disease or multiple plasmacytomas). Given the underlying plasma cell dyscrasia of POEMS syndrome, newer approaches to MM, including bortezomib, lenalidomide, and thalidomide, have also been investigated.

Hematopoietic Cell Transplantation

Hematopoietic cell transplantation is a procedure in which hematopoietic stem cells are intravenously infused to restore bone marrow and immune function in cancer patients who receive

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

bone marrow-toxic doses of cytotoxic drugs with or without whole-body radiotherapy. Hematopoietic stem cells may be obtained from the transplant recipient (autologous HCT) or a donor (allogeneic HCT [allo-HCT]). They can be harvested from bone marrow, peripheral blood, or umbilical cord blood shortly after delivery of neonates.

Immunologic compatibility between infused hematopoietic stem cells and the recipient is not an issue in autologous HCT. In allo-HCT, immunologic compatibility between donor and patient is a critical factor for achieving a successful outcome. Compatibility is established by typing of human leukocyte antigens (HLA) using cellular, serologic, or molecular techniques. HLA refers to the gene complex expressed at the HLA-A, -B, and -DR (antigen-D related) loci on each arm of chromosome 6. An acceptable donor will match the patient at all or most of the HLA loci.

Conditioning for Hematopoietic Cell Transplantation

Conventional Conditioning

The conventional (“classical”) practice of allo-HCT involves administration of cytotoxic agents (e.g., cyclophosphamide, busulfan) with or without total body irradiation at doses sufficient to cause bone marrow ablation in the recipient. The beneficial treatment effect of this procedure is due to a combination of the initial eradication of malignant cells and subsequent graft-versus-malignancy effect mediated by non-self-immunologic effector cells. While the slower graft-versus-malignancy effect is considered the potentially curative component, it may be overwhelmed by existing disease in the absence of pretransplant conditioning. Intense conditioning regimens are limited to patients who are sufficiently medically fit to tolerate substantial adverse effects. These include opportunistic infections secondary to loss of endogenous bone marrow function and organ damage or failure caused by cytotoxic drugs. Subsequent to graft infusion in allo-HCT, immunosuppressant drugs are required to minimize graft rejection and graft-versus-host disease, which increases susceptibility to opportunistic infections.

The success of autologous HCT is predicated on the potential of cytotoxic chemotherapy, with or without radiotherapy, to eradicate cancerous cells from the blood and bone marrow. This permits subsequent engraftment and repopulation of the bone marrow with presumably normal hematopoietic stem cells obtained from the patient before undergoing bone marrow ablation. Therefore, autologous HCT is typically performed as consolidation therapy when the patient’s disease is in complete remission. Patients who undergo autologous HCT are also susceptible to

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

chemotherapy-related toxicities and opportunistic infections before engraftment, but not graft-versus-host disease.

Reduced-Intensity Conditioning Allogeneic Hematopoietic Cell Transplantation

Reduced-intensity conditioning (RIC) refers to the pretransplant use of lower doses of cytotoxic drugs or less intense regimens of radiotherapy than are used in traditional full-dose myeloablative conditioning treatments. Although the definition of RIC is variable, with numerous versions employed, all regimens seek to balance the competing effects of relapse due to residual disease and non-relapse mortality. The goal of RIC is to reduce disease burden and to minimize associated treatment-related morbidity and non-relapse mortality in the period during which the beneficial graft-versus-malignancy effect of allogeneic transplantation develops. Reduced-intensity conditioning regimens range from nearly total myeloablative to minimally myeloablative with lymphoablation, with intensity tailored to specific diseases and patient condition. Patients who undergo RIC with allo-HCT initially demonstrate donor cell engraftment and bone marrow mixed chimerism. Most will subsequently convert to full-donor chimerism. In this review, the term reduced-intensity conditioning will refer to all conditioning regimens intended to be nonmyeloablative.

Multiple Myeloma Treatment Overview

In the prechemotherapy era, the median survival for a patient diagnosed with MM was approximately 7 months. After the introduction of chemotherapy (eg, the alkylating agent melphalan in the 1960s), prognosis improved, with a median survival of 24 to 30 months and 10-year survival of 3%. In a large group of patients with newly diagnosed MM, there was no difference in overall survival reported during a 24-year period from 1971 to 1994, with a trend toward improvement during 1995 to 2000, and a statistically significant benefit in overall survival during 2001 to 2006. These data suggested that autologous HCT was responsible for the trends during 1994 to 2000, while novel agents have contributed to the improvement since 2001.

The introduction of novel agents and better prognostic indicators has been the major advances in the treatment of this disease. Novel agents such as the proteasome inhibitors (eg, bortezomib), the monoclonal antibody daratumumab, and the immunomodulatory derivatives thalidomide and lenalidomide first showed efficacy in relapsed and refractory myeloma and now have been integrated into first-line regimens. With the introduction of these novel treatments, it is now expected that most

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

patients with MM will respond to initial therapy, and only a small minority will have refractory disease.

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

The U.S. FDA regulates human cells and tissues intended for implantation, transplantation, or infusion through the Center for Biologics Evaluation and Research, under Code of Federal Regulation, Title 21, parts 1270 and 1271. Hematopoietic stem cells are included in these regulations.

Rationale/Source

Description

Multiple myeloma (MM) is a systemic malignancy of plasma cells that represents approximately 10% of all hematologic cancers. POEMS syndrome, characterized by polyneuropathy, organomegaly, endocrinopathy, M protein, and skin changes, is a rare, paraneoplastic disorder secondary to a plasma cell dyscrasia. Plasma cell dyscrasias are treatable but rarely curable. In some cases, autologous or allogeneic HCT is considered as therapy.

Summary of Evidence

Newly Diagnosed Multiple Myeloma

For individuals who have newly diagnosed MM who receive autologous HCT as initial treatment, the evidence includes several prospective randomized controlled trials (RCTs) that compared high-dose chemotherapy plus autologous HCT to standard chemotherapy regimens or regimens containing newer MM agents. Relevant outcomes are overall survival (OS) and treatment-related morbidity. In general, the evidence has suggested OS rates are improved with autologous HCT compared with conventional chemotherapy in this setting. Limitations of the published evidence include patient heterogeneity, variability in treatment protocols, short follow-up periods, inconsistency in reporting important health outcomes, and inconsistency in reporting or collecting outcomes. Recent RCTs comparing high-dose chemotherapy plus autologous HCT to regimens that include novel MM agents have also shown that high-dose chemotherapy plus autologous HCT

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

improves progression-free survival. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have newly diagnosed MM who receive tandem autologous HCT, the evidence includes several RCTs. Relevant outcomes are OS and treatment-related morbidity. Compared with single autologous HCT, RCTs have generally found that tandem autologous HCT improve OS and recurrence-free survival in newly diagnosed MM. Two recent RCTs found conflicting results on the benefit of tandem autologous HCT versus single autologous HCT; however, the study that found no additional benefit with tandem autologous HCT had a higher rate of nonadherence to the second planned HCT. Differences in initial therapy regimens between trials may also have led to conflicting results. Several RCTs compared reduced-intensity conditioning allogeneic HCT (allo-HCT) following a first autologous HCT with single or tandem autologous transplants. The RCTs were based on genetic randomization (ie, patients with a human leukocyte antigen-identical sibling were offered reduced-intensity conditioning allo-HCT following autologous HCT, whereas other patients underwent either 1 or 2 autologous transplants). Although the body of evidence has shown inconsistencies regarding OS and disease-free survival rates, some studies have shown a survival benefit with tandem autologous HCT followed by reduced-intensity conditioning allo-HCT, although at the cost of higher transplant-related mortality compared with conventional treatments. Factors across studies that may account for differing trial results include different study designs, nonuniform preparative regimens, different patient characteristics (including risk stratification), and criteria for advancing to a second transplant. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have newly diagnosed MM who receive allo-HCT as initial or salvage treatment, the evidence includes nonrandomized studies. Relevant outcomes are OS and treatment-related morbidity. Studies have reported on patients with both myeloablative conditioning and reduced-intensity conditioning. Limitations of the published evidence include patient sample heterogeneity, variability in treatment protocols, short follow-up periods, inconsistency in reporting important health outcomes, and inconsistency in reporting or collecting outcomes. Nonmyeloablative allo-HCT as first-line therapy is associated with lower transplant-related mortality but a greater risk of relapse; convincing evidence is lacking that allo-HCT improves survival better than autologous HCT. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

Relapsed or Refractory Multiple Myeloma

For individuals who have relapsed MM after failing an autologous HCT who receive autologous HCT, the evidence includes RCTs, retrospective studies, and reviews summarizing recent studies on a second autologous HCT in relapsed myeloma. Relevant outcomes are OS and treatment-related morbidity. Despite some limitations of the published evidence, including patient sample heterogeneity, variability in treatment protocols, and short follow-up periods, the available trial evidence has suggested OS rates are improved with autologous HCT compared with conventional chemotherapy or continuous lenalidomide plus dexamethasone in this setting. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have refractory multiple myeloma after failing a first HCT who receive tandem autologous HCT, the evidence includes systematic reviews and a retrospective study. Relevant outcomes are OS and treatment-related morbidity. The evidence has shown tandem autologous HCT improves OS rates in this setting. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

POEMS Syndrome

For individuals who have POEMS syndrome who receive HCT, the evidence includes retrospective cohort studies, case reports, and case series. Relevant outcomes are OS and treatment-related morbidity. No RCTs of HCT of any type have been performed in patients with POEMS syndrome of any severity, nor is it likely such studies will be performed because of the rarity of this condition. Available case reports and series are subject to selection bias and are heterogeneous concerning treatment approaches and peritransplant support. However, for patients with disseminated POEMS syndrome, a chain of evidence and contextual factors related to the disease and MM would suggest improvement in health outcomes with autologous HCT. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

Additional Information

2017 Input

In response to requests, input was received from 1 specialty medical society, 1 academic medical center, and 2 Blue Distinction Centers for Transplant while this policy was under review in 2017.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

There was a consensus that allogeneic HCT is investigational for newly diagnosed multiple myeloma and as salvage therapy after primary graft failure and for the primary progressive disease.

2013 Input

In response to requests, input was received from 3 academic medical centers and 6 Blue Distinction Centers for Transplant while this policy was under review in 2013. There was near-consensus that autologous HCT is medically necessary for POEMS syndrome and near-consensus that allogeneic and tandem HCT are investigational for POEMS syndrome.

Supplemental Information

Clinical Input From Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

2017 Input

In response to requests, input was received from 1 specialty medical society, 1 academic medical center, and 2 Blue Distinction Centers for Transplant while this policy was under review in 2017. There was a consensus that allogeneic HCT is investigational for newly diagnosed multiple myeloma and as salvage therapy after primary graft failure and for the primary progressive disease.

2013 Input

In response to requests, input was received from 3 academic medical centers and 6 Blue Distinction Centers for Transplant while this policy was under review in 2013. There was near-consensus that autologous HCT is medically necessary for POEMS syndrome and near-consensus that allogeneic and tandem HCT are investigational for POEMS syndrome.

Practice Guidelines and Position Statements

American Society of Clinical Oncology

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

In 2019, the American Society of Clinical Oncology (ASCO) published practice guidelines for the treatment of MM. The guidelines recommend offering up-front transplant to all eligible patients, although delayed HCT may be considered in select patients. Salvage or delayed HCT may be used as consolidation at first relapse in patients who choose not to proceed with HCT initially. Tandem autologous HCT and allogeneic HCT (allo-HCT) should not be routinely recommended. However, up-front tandem autologous HCT can be considered for select high-risk patients or those with a suboptimal response to the initial transplant; allo-HCT may be considered in select high-risk patients in the context of a clinical trial. For relapsed MM, autologous HCT, if not received after primary induction therapy, should be offered to transplant-eligible patients. Repeat HCT may be considered in relapsed MM if progression-free survival after the first transplant was 18 months or greater.

American Society for Blood and Marrow Transplantation

In 2015, the American Society for Blood and Marrow Transplantation (ASBMT; now referred to as the American Society for Transplantation and Cellular Therapy) published evidence-based guidelines on the use of HCT in patients with MM. The ASBMT recognized that much of the evidence from RCTs summarized in the 2015 guidelines came from trials that predated the novel triple-therapy induction regimens. Furthermore, advances in supportive care and earlier disease detection have increasingly influenced decision making and allow individual tailoring of therapy. The ASBMT guidelines did not address POEMS or other plasma cell dyscrasias besides MM.

The ASBMT and 3 other groups (2015) published joint guidelines based on an expert consensus conference. These guidelines contained the following recommendations for HCT as salvage therapy:

"...autologous HCT: (1) In transplantation-eligible patients relapsing after primary therapy that did NOT include an autologous HCT, high-dose therapy with HCT as part of salvage therapy should be considered standard; (2) High-dose therapy and autologous HCT should be considered appropriate therapy for any patients relapsing after primary therapy that includes an autologous HCT with initial remission duration of more than 18 months; (3) High-dose therapy and autologous HCT can be used as bridging strategy to allogeneic HCT; (4) The role of post salvage HCT maintenance needs to be explored in the context of well-designed prospective trials that should include new agents, such as monoclonal antibodies, -modulating agents, and oral proteasome inhibitors; (5) Autologous HCT consolidation should be explored as a strategy to develop novel conditioning regimens or post-HCT strategies in patients with short remission (less than 18 months remissions) after primary therapy

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

(and (6) Prospective randomized trials need to be performed to define the role of salvage autologous HCT in patients with MM [multiple myeloma] relapsing after primary therapy comparing to 'best non-HCT' therapy.

Regarding allogeneic HCT... (1) Allogeneic HCT should be considered appropriate therapy for any eligible patient with early relapse (less than 24 months) after primary therapy that included an autologous HCT and/or with high-risk features (ie, cytogenetics, extramedullary disease, plasma cell leukemia, or high lactate dehydrogenase); (2) Allogeneic HCT should be performed in the context of a clinical trial if possible; (3) The role of post allogeneic HCT maintenance therapy needs to be explored in the context of well-designed prospective trials; and (4) Prospective randomized trials need to be performed to define the role of salvage allogeneic HCT in patients with MM relapsing after primary therapy."

International Myeloma Working Group

The 2010 conclusions and recommendations of the International Myeloma Working Group consensus statement on the current status of allo-HCT for MM are as follows: myeloablative allo-HCT may cure a minority of patients but is associated with high transplant-related mortality, but could be evaluated in well-designed prospective clinical trials. Nonmyeloablative allo-HCT as first-line therapy is associated with lower transplant-related mortality but a greater risk of relapse and convincing evidence is lacking that allo-HCT improves survival compared with autologous HCT.

National Comprehensive Cancer Network

Autologous Hematopoietic Cell Transplantation

The NCCN guidelines (v.3.2021) state that autologous HCT is the preferred option after induction therapy in transplant-eligible patients, but a delayed HCT after early stem cell collection and storage is appropriate as well (category 1 recommendation). A repeat HCT can be considered for refractory/progressive disease after primary treatment in patients with prolonged response to initial HCT.

Tandem Hematopoietic Cell Transplantation

The NCCN recommends collecting enough stem cells for 2 transplants in younger patients if tandem transplant or salvage transplant would be considered. A tandem transplant with or without

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

maintenance therapy can be considered for all patients who are candidates for HCT and is an option for patients who do not achieve at least a very good partial response after the first autologous HCT and those with high-risk features.

Allogeneic Hematopoietic Cell Transplantation

The NCCN states the following for allo-HCT: "Allogeneic HCT includes either myeloablative or nonmyeloablative (ie, "mini" transplant) transplants. Allogeneic HCT has been investigated as an alternative to autologous HCT to avoid the contamination of reinfused autologous tumor cells, but also to take advantage of the beneficial graft-versus-tumor effect associated with allogeneic transplants. However, lack of a suitable donor and increased morbidity has limited this approach, particularly for the typical older MM population". The guidelines also note that allo-HCT should be done in the context of a clinical trial when possible.

POEMS Syndrome

The NCCN guidelines recommend autologous HCT in patients with POEMS syndrome who are eligible as sole therapy or as consolidation therapy after induction therapy.

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

Medicare has the following national coverage determination for the use of HCT for MM.

“Effective ... January ... 2016, allogeneic HSCT [hematopoietic stem cell transplantation] for multiple myeloma is covered by Medicare only for beneficiaries with Durie-Salmon Stage II or III multiple myeloma, or International Staging System (ISS) Stage II or Stage III multiple myeloma, and participating in an approved prospective clinical study that meets the criteria below. There must be appropriate statistical techniques to control for selection bias and confounding by age, duration of diagnosis, disease classification, International Myeloma Working Group (IMWG) classification, ISS stage, comorbid conditions, type of preparative/conditioning regimen, graft vs. host disease (GVHD) prophylaxis, donor type and cell source.

A prospective clinical study seeking Medicare coverage for allogeneic HSCT for multiple myeloma pursuant to CED must address the following question:

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

Compared to patients who do not receive allogeneic HSCT, do Medicare beneficiaries with multiple myeloma who receive allogeneic HSCT have improved outcomes as indicated by:

- Graft vs. host disease (acute and chronic);
- Other transplant-related adverse events;
- Overall survival; and
- (optional) Quality of life?"

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 2.

Table 2. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT01208662 ^a	A Randomized Phase III Study Comparing Conventional Dose Treatment Using a Combination of Lenalidomide, Bortezomib, and Dexamethasone (RVD) to High-dose Treatment With Peripheral Stem Cell Transplant in the Initial Management of Myeloma in Patients up to 65 Years of Age	660	Sep 2023
<i>Unpublished</i>			
NCT02322320	Continued, Long-Term Follow-Up and Lenalidomide Maintenance Therapy for Patients on BMT CTN 0702 (BMT CTN #Q07LT)	450	Jun 2019 (Completed)
NCT00998270	Autologous Bone Marrow Transplantation (BMT) Compared With Allogeneic BMT in Multiple Myeloma	185	Oct 2017 (unknown)

NCT: national clinical trial.

^a Denotes industry-sponsored or cosponsored trial.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

References

1. Blue Cross and Blue Shield Association, Medical Policy Reference Manual, “Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome”, 8.01.17, 2:2021.
2. Kyle RA, Rajkumar SV. Multiple myeloma. *Blood*. Mar 15 2008; 111(6): 2962-72. PMID 18332230
3. Palumbo A, Rajkumar SV. Treatment of newly diagnosed myeloma. *Leukemia*. Mar 2009; 23(3): 449-56. PMID 19005483
4. Durie BG, Harousseau JL, Miguel JS, et al. International uniform response criteria for multiple myeloma. *Leukemia*. Sep 2006; 20(9): 1467-73. PMID 16855634
5. Dispenzieri A. Long-term outcomes after autologous stem cell transplantation in patients with POEMS syndrome. *Clin Adv Hematol Oncol*. Nov 2012; 10(11): 744-6. PMID 23271262
6. Dispenzieri A. POEMS syndrome: update on diagnosis, risk-stratification, and management. *Am J Hematol*. Aug 2012; 87(8): 804-14. PMID 22806697
7. Bardwick PA, Zvaifler NJ, Gill GN, et al. Plasma cell dyscrasia with polyneuropathy, organomegaly, endocrinopathy, M protein, and skin changes: the POEMS syndrome. Report on two cases and a review of the literature. *Medicine (Baltimore)*. Jul 1980; 59(4): 311-22. PMID 6248720
8. Dispenzieri A, Kyle RA, Lacy MQ, et al. POEMS syndrome: definitions and long-term outcome. *Blood*. Apr 01 2003; 101(7): 2496-506. PMID 12456500
9. Dispenzieri A. POEMS Syndrome: 2019 Update on diagnosis, risk-stratification, and management. *Am J Hematol*. Jul 2019; 94(7): 812-827. PMID 31012139
10. Nasu S, Misawa S, Sekiguchi Y, et al. Different neurological and physiological profiles in POEMS syndrome and chronic inflammatory demyelinating polyneuropathy. *J Neurol Neurosurg Psychiatry*. May 2012; 83(5): 476-9. PMID 22338030
11. Reece DE. Recent trends in the management of newly diagnosed multiple myeloma. *Curr Opin Hematol*. Jul 2009; 16(4): 306-12. PMID 19491669
12. Reece D HJ, Gertz MA. Myeloma Management 2009: Nontransplant therapy of myeloma, high-dose therapy for myeloma, and a personalised care plan for treatment of myeloma. 2009 American Society of Clinical Oncology Annual Meeting Educational Handbook. 2009:502-509.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

13. Qiao SK, Guo XN, Ren JH, et al. Efficacy and Safety of Lenalidomide in the Treatment of Multiple Myeloma: A Systematic Review and Meta-analysis of Randomized Controlled Trials. *Chin Med J (Engl)*. May 05 2015; 128(9): 1215-22. PMID 25947406
14. Rajkumar SV, Kumar S. Multiple myeloma current treatment algorithms. *Blood Cancer J*. Sep 28 2020; 10(9): 94. PMID 32989217
15. Fonseca R. Strategies for risk-adapted therapy in myeloma. *Hematology Am Soc Hematol Educ Program*. 2007: 304-10. PMID 18024644
16. Rajkumar SV. Multiple myeloma: 2020 update on diagnosis, risk-stratification and management. *Am J Hematol*. May 2020; 95(5): 548-567. PMID 32212178
17. Cavo M, Gay F, Beksac M, et al. Autologous haematopoietic stem-cell transplantation versus bortezomib-melphalan-prednisone, with or without bortezomib-lenalidomide-dexamethasone consolidation therapy, and lenalidomide maintenance for newly diagnosed multiple myeloma (EMN02/HO95): a multicentre, randomised, open-label, phase 3 study. *Lancet Haematol*. Jun 2020; 7(6): e456-e468. PMID 32359506
18. Attal M, Lauwers-Cances V, Hulin C, et al. Lenalidomide, Bortezomib, and Dexamethasone with Transplantation for Myeloma. *N Engl J Med*. Apr 06 2017; 376(14): 1311-1320. PMID 28379796
19. Gay F, Oliva S, Petrucci MT, et al. Chemotherapy plus lenalidomide versus autologous transplantation, followed by lenalidomide plus prednisone versus lenalidomide maintenance, in patients with multiple myeloma: a randomised, multicentre, phase 3 trial. *Lancet Oncol*. Dec 2015; 16(16): 1617-29. PMID 26596670
20. Attal M, Harousseau JL. The role of high-dose therapy with autologous stem cell support in the era of novel agents. *Semin Hematol*. Apr 2009; 46(2): 127-32. PMID 19389496
21. Attal M, Harousseau JL, Stoppa AM, et al. A prospective, randomized trial of autologous bone marrow transplantation and chemotherapy in multiple myeloma. *Intergroupe Francais du Myelome*. *N Engl J Med*. Jul 11 1996; 335(2): 91-7. PMID 8649495
22. Barlogie B, Kyle RA, Anderson KC, et al. Standard chemotherapy compared with high-dose chemoradiotherapy for multiple myeloma: final results of phase III US Intergroup Trial S9321. *J Clin Oncol*. Feb 20 2006; 24(6): 929-36. PMID 16432076
23. Blade J, Rosinol L, Sureda A, et al. High-dose therapy intensification compared with continued standard chemotherapy in multiple myeloma patients responding to the initial chemotherapy: long-term results from a prospective randomized trial from the Spanish cooperative group PETHEMA. *Blood*. Dec 01 2005; 106(12): 3755-9. PMID 16105975

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

24. Child JA, Morgan GJ, Davies FE, et al. High-dose chemotherapy with hematopoietic stem-cell rescue for multiple myeloma. *N Engl J Med.* May 08 2003; 348(19): 1875-83. PMID 12736280
25. Femand JP, Ravaud P, Chevret S, et al. High-dose therapy and autologous peripheral blood stem cell transplantation in multiple myeloma: up-front or rescue treatment? Results of a multicenter sequential randomized clinical trial. *Blood.* Nov 01 1998; 92(9): 3131-6. PMID 9787148
26. Palumbo A, Bringhen S, Petrucci MT, et al. Intermediate-dose melphalan improves survival of myeloma patients aged 50 to 70: results of a randomized controlled trial. *Blood.* Nov 15 2004; 104(10): 3052-7. PMID 15265788
27. Mian H, Mian OS, Rochweg B, et al. Autologous stem cell transplant in older patients (age65) with newly diagnosed multiple myeloma: A systematic review and meta-analysis. *J Geriatr Oncol.* Jan 2020; 11(1): 93-99. PMID 31153809
28. Koreth J, Cutler CS, Djulbegovic B, et al. High-dose therapy with single autologous transplantation versus chemotherapy for newly diagnosed multiple myeloma: A systematic review and meta-analysis of randomized controlled trials. *Biol Blood Marrow Transplant.* Feb 2007; 13(2): 183-96. PMID 17241924
29. Marini C, Maia T, Bergantim R, et al. Real-life data on safety and efficacy of autologous stem cell transplantation in elderly patients with multiple myeloma. *Ann Hematol.* Feb 2019; 98(2): 369-379. PMID 30368589
30. Attal M, Harousseau JL, Facon T, et al. Single versus double autologous stem-cell transplantation for multiple myeloma. *N Engl J Med.* Dec 25 2003; 349(26): 2495-502. PMID 14695409
31. Stadtmauer EA. Multiple myeloma, 2004--one or two transplants?. *N Engl J Med.* Dec 25 2003; 349(26): 2551-3. PMID 14695416
32. Cavo M, Tosi P, Zamagni E, et al. Prospective, randomized study of single compared with double autologous stem-cell transplantation for multiple myeloma: Bologna 96 clinical study. *J Clin Oncol.* Jun 10 2007; 25(17): 2434-41. PMID 17485707
33. Stadtmauer EA, Pasquini MC, Blackwell B, et al. Autologous Transplantation, Consolidation, and Maintenance Therapy in Multiple Myeloma: Results of the BMT CTN 0702 Trial. *J Clin Oncol.* Mar 01 2019; 37(7): 589-597. PMID 30653422
34. Garban F, Attal M, Michallet M, et al. Prospective comparison of autologous stem cell transplantation followed by dose-reduced allograft (IFM99-03 trial) with tandem autologous stem cell transplantation (IFM99-04 trial) in high-risk de novo multiple myeloma. *Blood.* May 01 2006; 107(9): 3474-80. PMID 16397129

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

35. Moreau P, Garban F, Attal M, et al. Long-term follow-up results of IFM99-03 and IFM99-04 trials comparing nonmyeloablative allotransplantation with autologous transplantation in high-risk de novo multiple myeloma. *Blood*. Nov 01 2008; 112(9): 3914-5. PMID 18948589
36. Bruno B, Rotta M, Patriarca F, et al. A comparison of allografting with autografting for newly diagnosed myeloma. *N Engl J Med*. Mar 15 2007; 356(11): 1110-20. PMID 17360989
37. Rosinol L, Perez-Simon JA, Sureda A, et al. A prospective PETHEMA study of tandem autologous transplantation versus autograft followed by reduced-intensity conditioning allogeneic transplantation in newly diagnosed multiple myeloma. *Blood*. Nov 01 2008; 112(9): 3591-3. PMID 18612103
38. Bjorkstrand B, Iacobelli S, Hegenbart U, et al. Autologous stem cell transplantation (ASCT) versus ASCT followed by reduced-intensity conditioning allogeneic SCT with identical sibling donor in previously untreated multiple myeloma: preliminary analysis of a prospective controlled trial by the EBMT [abstract]. *Bone Marrow Transplant*. 2008;41:S38.
39. Gahrton G, Iacobelli S, Bjorkstrand B, et al. Autologous/reduced-intensity allogeneic stem cell transplantation vs autologous transplantation in multiple myeloma: long-term results of the EBMT-NMAM2000 study. *Blood*. Jun 20 2013; 121(25): 5055-63. PMID 23482933
40. Krishnan A, Pasquini MC, Logan B, et al. Autologous haemopoietic stem-cell transplantation followed by allogeneic or autologous haemopoietic stem-cell transplantation in patients with multiple myeloma (BMT CTN 0102): a phase 3 biological assignment trial. *Lancet Oncol*. Dec 2011; 12(13): 1195-203. PMID 21962393
41. Maffini E, Storer BE, Sandmaier BM, et al. Long-term follow up of tandem autologous-allogeneic hematopoietic cell transplantation for multiple myeloma. *Haematologica*. Feb 2019; 104(2): 380-391. PMID 30262560
42. Giralt S, Koehne G. Allogeneic hematopoietic stem cell transplantation for multiple myeloma: what place, if any?. *Curr Hematol Malig Rep*. Dec 2013; 8(4): 284-90. PMID 24146203
43. Giralt S, Costa L, Schriber J, et al. Optimizing autologous stem cell mobilization strategies to improve patient outcomes: consensus guidelines and recommendations. *Biol Blood Marrow Transplant*. Mar 2014; 20(3): 295-308. PMID 24141007
44. Harousseau JL. The allogeneic dilemma. *Bone Marrow Transplant*. Dec 2007; 40(12): 1123-8. PMID 17680016
45. Crawley C, Iacobelli S, Bjorkstrand B, et al. Reduced-intensity conditioning for myeloma: lower nonrelapse mortality but higher relapse rates compared with myeloablative conditioning. *Blood*. Apr 15 2007; 109(8): 3588-94. PMID 17158231

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

46. Gahrton G, Bjorkstrand B. Allogeneic transplantation in multiple myeloma. *Haematologica*. Sep 2008; 93(9): 1295-300. PMID 18757850
47. Goldschmidt H, Baertsch MA, Schlenzka J, et al. Salvage autologous transplant and lenalidomide maintenance vs. lenalidomide/dexamethasone for relapsed multiple myeloma: the randomized GMMG phase III trial ReLApsE. *Leukemia*. Jul 21 2020. PMID 32694619
48. Cook G, Williams C, Brown JM, et al. High-dose chemotherapy plus autologous stem-cell transplantation as consolidation therapy in patients with relapsed multiple myeloma after previous autologous stem-cell transplantation (NCRI Myeloma X Relapse [Intensive trial]): a randomised, open-label, phase 3 trial. *Lancet Oncol*. Jul 2014; 15(8): 874-85. PMID 24948586
49. Cook G, Ashcroft AJ, Cairns DA, et al. The effect of salvage autologous stem-cell transplantation on overall survival in patients with relapsed multiple myeloma (final results from BSBMT/UKMF Myeloma X Relapse [Intensive]): a randomised, open-label, phase 3 trial. *Lancet Haematol*. Jul 2016; 3(7): e340-51. PMID 27374467
50. Ikeda T, Mori K, Kawamura K, et al. Comparison between autologous and allogeneic stem cell transplantation as salvage therapy for multiple myeloma relapsing/progressing after autologous stem cell transplantation. *Hematol Oncol*. Dec 2019; 37(5): 586-594. PMID 31674032
51. Michaelis LC, Saad A, Zhong X, et al. Salvage second hematopoietic cell transplantation in myeloma. *Biol Blood Marrow Transplant*. May 2013; 19(5): 760-6. PMID 23298856
52. Qazilbash MH, Saliba R, De Lima M, et al. Second autologous or allogeneic transplantation after the failure of first autograft in patients with multiple myeloma. *Cancer*. Mar 01 2006; 106(5): 1084-9. PMID 16456814
53. Ziogas DC, Terpos E, Dimopoulos MA. When to recommend a second autograft in patients with relapsed myeloma?. *Leuk Lymphoma*. Apr 2017; 58(4): 781-787. PMID 27894207
54. Garderet L, Cook G, Auner HW, et al. Treatment options for relapse after autograft in multiple myeloma - report from an EBMT educational meeting. *Leuk Lymphoma*. Apr 2017; 58(4): 797-808. PMID 27650125
55. Hahn T, Wingard JR, Anderson KC, et al. The role of cytotoxic therapy with hematopoietic stem cell transplantation in the therapy of multiple myeloma: an evidence-based review. *Biol Blood Marrow Transplant*. Jan 2003; 9(1): 4-37. PMID 12533739
56. Olin RL, Vogl DT, Porter DL, et al. Second auto-SCT is safe and effective salvage therapy for relapsed multiple myeloma. *Bone Marrow Transplant*. Mar 2009; 43(5): 417-22. PMID 18850013

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

57. McCarthy PL, Holstein SA. Role of stem cell transplant and maintenance therapy in plasma cell disorders. *Hematology Am Soc Hematol Educ Program*. Dec 02 2016; 2016(1): 504-511. PMID 27913522
58. Kuwabara S, Dispenzieri A, Arimura K, et al. Treatment for POEMS (polyneuropathy, organomegaly, endocrinopathy, M-protein, and skin changes) syndrome. *Cochrane Database Syst Rev*. Jun 13 2012; (6): CD006828. PMID 22696361
59. Autore F, Innocenti I, Luigetti M, et al. Autologous peripheral blood stem cell transplantation and the role of lenalidomide in patients affected by poems syndrome. *Hematol Oncol*. Apr 2018; 36(2): 392-398. PMID 28913957
60. D'Souza A, Lacy M, Gertz M, et al. Long-term outcomes after autologous stem cell transplantation for patients with POEMS syndrome (osteosclerotic myeloma): a single-center experience. *Blood*. Jul 05 2012; 120(1): 56-62. PMID 22611150
61. Jang IY, Yoon DH, Kim S, et al. Advanced POEMS syndrome treated with high-dose melphalan followed by autologous blood stem cell transplantation: a single-center experience. *Blood Res*. Mar 2014; 49(1): 42-8. PMID 24724066
62. Cook G, Iacobelli S, van Biezen A, et al. High-dose therapy and autologous stem cell transplantation in patients with POEMS syndrome: a retrospective study of the Plasma Cell Disorder sub-committee of the Chronic Malignancy Working Party of the European Society for Blood Marrow Transplantation. *Haematologica*. Jan 2017; 102(1): 160-167. PMID 27634201
63. Mikhael J, Ismaila N, Cheung MC, et al. Treatment of Multiple Myeloma: ASCO and CCO Joint Clinical Practice Guideline. *J Clin Oncol*. May 10 2019; 37(14): 1228-1263. PMID 30932732
64. Shah N, Callander N, Ganguly S, et al. Hematopoietic Stem Cell Transplantation for Multiple Myeloma: Guidelines from the American Society for Blood and Marrow Transplantation. *Biol Blood Marrow Transplant*. Jul 2015; 21(7): 1155-66. PMID 25769794
65. Giralt S, Garderet L, Durie B, et al. American Society of Blood and Marrow Transplantation, European Society of Blood and Marrow Transplantation, Blood and Marrow Transplant Clinical Trials Network, and International Myeloma Working Group Consensus Conference on Salvage Hematopoietic Cell Transplantation in Patients with Relapsed Multiple Myeloma. *Biol Blood Marrow Transplant*. Dec 2015; 21(12): 2039-2051. PMID 26428082
66. Lokhorst H, Einsele H, Vesole D, et al. International Myeloma Working Group consensus statement regarding the current status of allogeneic stem-cell transplantation for multiple myeloma. *J Clin Oncol*. Oct 10 2010; 28(29): 4521-30. PMID 20697091

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

- 67. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Multiple Myeloma. Version 3.2021. https://www.nccn.org/professionals/physician_gls/pdf/myeloma.pdf.
- 68. Centers for Medicare & Medicaid Services. National Coverage Determination (NCD) for Stem Cell Transplantation (Formerly 110.8.1) (110.23). 2016; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=366&ncdver=1&bc=AAAAIAAAAA&>.

Policy History

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

- 12/06/2001 Medical Policy Committee review
- 01/28/2002 Managed Care Advisory Council approval
- 12/06/2006 Medical Director review
- 12/20/2006 Medical Policy Committee approval. Coverage eligibility unchanged.
- 09/05/2007 Medical Director review
- 09/19/2007 Medical Policy Committee approval. Policy statement language regarding tandem transplants in newly diagnosed or responsive multiple myeloma clarified.
- 09/09/2008 Medical Director review
- 09/17/2008 Medical Policy Committee approval. No change to coverage eligibility.
- 09/03/2009 Medical Policy Committee approval
- 09/16/2009 Medical Policy Implementation Committee approval. No change to coverage eligibility.
- 09/09/2010 Medical Policy Committee review
- 09/15/2010 Medical Policy Implementation Committee approval. Policy title changed from “High-Dose Chemotherapy with Stem-Cell Support for Multiple Myeloma” to “Hematopoietic Stem-Cell Transplantation for Multiple Myeloma”. Policy language and statements extensively updated to reflect current practice.
- 09/01/2011 Medical Policy Committee review
- 09/14/2011 Medical Policy Implementation Committee approval. No changes to coverage.
- 09/06/2012 Medical Policy Committee review

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

09/19/2012 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

03/04/2013 Coding updated

10/03/2013 Medical Policy Committee review

10/16/2013 Medical Policy Implementation Committee approval. Title changed. Coverage for POEMS syndrome added.

11/06/2014 Medical Policy Committee review

11/21/2014 Medical Policy Implementation Committee approval. No change to coverage.

03/05/2015 Medical Policy Committee review

03/19/2015 Medical Policy Implementation Committee approval. Tandem autologous-autologous hematopoietic stem-cell transplantation to treat multiple myeloma clarified.

03/03/2016 Medical Policy Committee review

03/16/2016 Medical Policy Implementation Committee approval. No change to coverage.

01/01/2017 Coding update: Removing ICD-9 Diagnosis Codes

03/02/2017 Medical Policy Committee review

03/15/2017 Medical Policy Implementation Committee approval. No change to coverage.

03/01/2018 Medical Policy Committee review

03/21/2018 Medical Policy Implementation Committee approval. No change to coverage.

03/07/2019 Medical Policy Committee review

03/20/2019 Medical Policy Implementation Committee approval. No change to coverage.

03/05/2020 Medical Policy Committee review

03/11/2020 Medical Policy Implementation Committee approval. No change to coverage.

03/04/2021 Medical Policy Committee review

03/10/2021 Medical Policy Implementation Committee approval. No change to coverage.

Next Scheduled Review Date: 03/2022

Coding

The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)†, copyright 2020 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	38204, 38205, 38206, 38207, 38208, 38209, 38210, 38211, 38212, 38213, 38214, 38215, 38230, 38232, 38240, 38241, 38242, 38243
HCPCS	S2140, S2142, S2150
ICD-10 Diagnosis	C90.00-C90.02, E88.09

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or

- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
1. Consultation with the Blue Cross and Blue Shield Association technology assessment program (TEC) or other nonaffiliated technology evaluation center(s);
 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
 3. Reference to federal regulations.

**Medically Necessary (or “Medical Necessity”) - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. In accordance with nationally accepted standards of medical practice;
- B. Clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and
- C. Not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.

For these purposes, “nationally accepted standards of medical practice” means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

‡ Indicated trademarks are the registered trademarks of their respective owners.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Policy # 00060

Original Effective Date: 01/28/2002

Current Effective Date: 04/12/2021

NOTICE: If the Patient's health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.