



# Louisiana

## Leadless Cardiac Pacemakers

**Policy #** 00688

**Original Effective Date:** 01/01/2020

**Current Effective Date:** 11/09/2020

*Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.*

### Services Are Considered Investigational

*Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.*

Based on review of available data, the Company considers **leadless cardiac pacemakers**, including the Micra transcatheter pacing system, in all situations to be **investigational**.\*

### Background/Overview

#### **Conventional Pacemakers**

Pacemakers are intended to be used as a substitute for the heart's intrinsic pacing system to correct cardiac rhythm disorders. By providing an appropriate heart rate and heart rate response, cardiac pacemakers can reestablish effective circulation and more normal hemodynamics that are compromised by a slow heart rate. Pacemakers vary in system complexity and can have multiple functions as a result of the ability to sense and/or stimulate both the atria and the ventricles.

Transvenous pacemakers or pacemakers with leads (hereinafter referred to as conventional pacemakers) consist of two components: a pulse generator (ie, battery component) and electrodes (ie, leads). The pulse generator consists of a power supply and electronics that can provide periodic electrical pulses to stimulate the heart. The generator is commonly implanted in the infraclavicular region of the anterior chest wall and placed in a pre-pectoral position; in some cases, a subpectoral position is advantageous. The unit generates an electrical impulse, which is transmitted to the myocardium via the electrodes affixed to the myocardium to sense and pace the heart as needed.

Conventional pacemakers are also referred to as single-chamber or dual-chamber systems. In single-chamber systems, only one lead is placed, typically in the right ventricle. In dual-chamber pacemakers, two leads are placed one in the right atrium and the other in the right ventricle. Single-chamber ventricular pacemakers are more common.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Leadless Cardiac Pacemakers

Policy # 00688

Original Effective Date: 01/01/2020

Current Effective Date: 11/09/2020

Annually, approximately 200,000 pacemakers are implanted in the U. S. and 1 million worldwide. Implantable pacemakers are considered life-sustaining, life-supporting class III devices for patients with a variety of brady arrhythmias. Pacemaker systems have matured over the years with well-established, acceptable performance standards. As per the Food and Drug Administration (FDA), the early performance of conventional pacemaker systems from implantation through 60 to 90 days have usually demonstrated acceptable pacing capture thresholds and sensing. Intermediate performance (90 days through more than 5 years) has usually demonstrated the reliability of the pulse generator and lead technology. Chronic performance (5-10 years) includes a predictable decline in battery life and mechanical reliability but a vast majority of patients receive excellent pacing and sensing free of operative or mechanical reliability failures.

Even though the safety profile of conventional pacemakers is excellent, they are associated with complications particularly related to leads. Most safety data on the use of conventional pacemakers come from registries from Europe, particularly from Denmark where all pacemaker implants are recorded in a national registry. These data are summarized in Table 1. It is important to recognize that valid comparison of complication rates is limited by differences in definitions of complications, which results in a wide variance of outcomes, as well as by the large variance in follow-up times, use of single-chamber or dual-chamber systems, and data reported over more than two decades. As such, the following data are contemporary and limited to single-chamber systems when reported separately.

In many cases when a conventional pectoral approach is not possible, alternative approaches such as epicardial pacemaker implantation and trans-iliac approaches have been used. Cohen et al (2001) reported outcomes from a retrospective analysis of 123 patients who underwent 207 epicardial lead implantations. Congenital heart disease was present in 103 (84%) of the patients. Epicardial leads were followed for 29 months (range 1 to 207 months). Lead failure was defined as the need for replacement or abandonment due to pacing or sensing problems, lead fracture, or phrenic/muscle stimulation. The 1-, 2-, and 5-year lead survival was 96%, 90%, and 74%, respectively. Epicardial lead survival in those placed by a subxiphoid approach was 100% at 1 year and at 10 years, by the sternotomy approach (93.9% at 1 year and 75.9% at 10 years) and lateral thoracotomy approach (94.1% at 1 year and 62.4% at 10 years).

Doll et al (2008) reported results of a randomized controlled trial comparing epicardial implantation vs conventional pacemaker implantation in 80 patients with indications for cardiac resynchronization therapy. The authors reported that the conventional pacemaker group had a significantly shorter

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Leadless Cardiac Pacemakers

Policy # 00688

Original Effective Date: 01/01/2020

Current Effective Date: 11/09/2020

intensive care unit stay, less blood loss, and shorter ventilation times while the epicardial group had less exposure to radiation and less use of contrast medium. The left ventricular pacing threshold was similar in the two groups at discharge but longer in the epicardial group during follow-up. Adverse events were also similar in the two groups. The following events were experienced by one (3%) patient each in the epicardial group: pleural puncture, pneumothorax, wound infection, Acute Respiratory Distress Syndrome, and hospital mortality.

As a less invasive alternative to the epicardial approach, the trans-iliac approach has also been utilized. Data using trans-iliac approach is limited. Multiple other studies with smaller sample size report a wide range of lead longevity.

Harakeet al (2018) reported a retrospective analysis of 5 patients who underwent a transvenous iliac approach (median age 26.9 years). Pacing indications included AV block in three patients and sinus node dysfunction in two. After a median follow-up of 4.1 years (range 1.0-16.7 years), outcomes were reported for 4 patients. One patient underwent device revision for lead position-related groin discomfort; a second patient developed atrial lead failure following a Maze operation and underwent lead replacement by the iliac approach. One patient underwent heart transplantation six months after implant with only partial resolution of pacing-induced cardiomyopathy. Tsutsumi et al (2010) reported a case series of 4 patients from Japan in whom conventional pectoral approach was precluded due to recurrent lead infections (n=1), superior vena cava obstruction following cardiac surgery (n=2) and a postoperative dermal scar (n=1). The mean follow-up was 24 months and the authors concluded the iliac vein approach was satisfactory and less invasive alternative to epicardial lead implantation. However, the authors reported that the incidence of atrial lead dislodgement using this approach in the literature ranged from 7 to 21%. Experts who provided clinical input reported that trans-iliac or surgical epicardial approach requires special expertise and long-term performance is suboptimal.

Table 1. Reported Complication Rates with Conventional Pacemakers

Complications	Rates, % <sup>a</sup>
Traumatic complications	

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Leadless Cardiac Pacemakers

Policy # 00688

Original Effective Date: 01/01/2020

Current Effective Date: 11/09/2020

RV perforation	0.2-0.8
RV perforation with tamponade	0.07-0.4
Pneumo (hemo) thorax	0.7-2.2
<b>Pocket complications</b>	
Including all hematomas, difficult to control bleeding, infection, discomfort, skin erosion	4.75
Including only those requiring invasive correction or reoperation	0.66-1.0
<b>Lead-related complications</b>	
Including lead fracture, dislodgement, insulation problem, infection, stimulation threshold problem, diaphragm or pocket stimulation, other	1.6-3.8
All system-related infections requiring reoperation or extraction	0.5-0.7

Adapted from FDA executive summary memorandum (2016).

a Rates are for new implants only and ventricular single-chamber devices when data were available. Some rates listed in this column are for single- and dual-chamber devices when data were not separated in the publication. Note that Micra transcatheter pacing system is a single-chamber device.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Leadless Cardiac Pacemakers

Policy # 00688

Original Effective Date: 01/01/2020

Current Effective Date: 11/09/2020

### **Potential Advantages of Leadless Cardiac Pacemakers Over Conventional Pacemakers**

The potential advantages of leadless pacemakers fall into three categories: avoidance of risks associated with intravascular leads in conventional pacemakers, avoidance of risks associated with pocket creation for placement of conventional pacemakers, and an additional option for patients who require a single-chamber pacer.

Lead complications include lead failure, lead fracture, insulation defect, pneumothorax, infections requiring lead extractions and replacements that can result in a torn subclavian vein or the tricuspid valve. In addition, there are risks of venous thrombosis and occlusion of the subclavian system from the leads. Use of a leadless system eliminates such risks with the added advantage that a patient has vascular access preserved for other medical conditions (eg, dialysis, chemotherapy).

Pocket complications include infections, erosions, and pain that can be eliminated with leadless pacemakers. Further, a leadless cardiac pacemaker may be more comfortable and appealing because unlike conventional pacemakers, patients are unable to see or feel the device or have an implant scar on the chest wall.

Leadless pacemakers may also be a better option than surgical endocardial pacemakers for patients with no vascular access due to renal failure or congenital heart disease.

### **Leadless Cardiac Pacemakers in Clinical Development**

Leadless pacemakers are self-contained in a hermetically sealed capsule. The capsule houses a battery and electronics to operate the system. Similar to most pacing leads, the tip of the capsule includes a fixation mechanism and a monolithic controlled-release device. The controlled-release device elutes gluco-corticosteroid to reduce acute inflammation at the implantation site. Leadless pacemakers have rate-responsive functionality, and current device longevity estimates are based on bench data. Estimates have suggested that these devices may last over ten years, depending on the programmed parameters.

Three systems are currently being evaluated in clinical trials: (1) the Micra Transcatheter Pacing System (Medtronic), (2) the Nanostim leadless pacemaker (St. Jude Medical); and (3) the WiCS Wireless Cardiac Stimulation System (EBR Systems). The first two devices are free-standing capsule-sized devices that are delivered via femoral venous access using a steerable delivery sheath. However, the fixing mechanism differs between the two devices. In the Micra Transcatheter Pacing System, the fixation system consists of four self-expanding nitinol tines, which anchor into the myocardium; for the

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Leadless Cardiac Pacemakers

Policy # 00688

Original Effective Date: 01/01/2020

Current Effective Date: 11/09/2020

Nanostim device, there is a screw-in helix that penetrates about 1 mm into the myocardium, with nylon tines that provide secondary fixation. In both devices, the cathode is steroid eluting and delivers pacing current; the anode is located in a titanium case. The third device, WiCS system differs from the other devices; this system requires implanting a pulse generator subcutaneously near the heart, which then wirelessly transmits ultrasound energy to a receiver electrode implanted in the left ventricle. The receiver electrode converts the ultrasound energy and delivers electrical stimulation to the heart sufficient to pace the left ventricle synchronously with the right.

Of these three, only the Micra transcatheter pacing system is approved by the FDA and commercially available in the U. S. Multiple clinical studies of Nanostim have been published, but trials have been halted due to the migration of the docking button in the device. Evidence on Nanostim is not reviewed further because the device is not yet FDA approved.

The Micra is about 26 mm in length and introduced using a 23 French catheter via the femoral vein to the right ventricle. It weighs about 2 grams and has an accelerometer-based rate response.

Nanostim is about 40 mm in length and introduced using an 18 French catheter to the right ventricle. It also weighs about 2 grams and uses a temperature-based rate response sensor.

## **FDA or Other Governmental Regulatory Approval**

### **U.S. Food and Drug Administration (FDA)**

In April 2016, the Micra™<sup>†</sup> transcatheter pacing system (Medtronic) was approved by the FDA through the premarket approval process for use in patients who have experienced one or more of the following conditions:

- Symptomatic paroxysmal or permanent high-grade arteriovenous block in the presence of atrial fibrillation
- Paroxysmal or permanent high-grade arteriovenous block in the absence of atrial fibrillation, as an alternative to dual-chamber pacing, when atrial lead placement is considered difficult, high-risk, or not deemed necessary for effective therapy
- Symptomatic bradycardia-tachycardia syndrome or sinus node dysfunction (sinus bradycardia or sinus pauses), as an alternative to atrial or dual-chamber pacing, when atrial lead placement is considered difficult, high-risk, or not deemed necessary for effective therapy.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Leadless Cardiac Pacemakers

Policy # 00688

Original Effective Date: 01/01/2020

Current Effective Date: 11/09/2020

In January 2020, the Micra AV Transcatheter Pacing System Model MC1AVR1 and Application Software Model SW044 were approved as a PMA supplement (S061) to the Micra system described above. The Micra AV includes an enhanced algorithm to provide AV synchronous pacing.

As per the FDA label, the Micra Model MC1VR01 pacemaker is contraindicated for patients who have the following types of devices implanted:

- An implanted device that would interfere with the implant of the Micra device in the judgment of the implanting physician
- An implanted inferior vena cava filter
- A mechanical tricuspid valve
- An implanted cardiac device providing active cardiac therapy which may interfere with the sensing performance of the Micra device

As per the FDA label, the Micra Model MC1VR01 pacemaker is also contraindicated for patients who have the following conditions:

- Femoral venous anatomy unable to accommodate a 7.8 mm (23 French) introducer sheath or implant on the right side of the heart (for example, due to obstructions or severe tortuosity)
- Morbid obesity that prevents the implanted device to obtain telemetry communication within <12.5 cm (4.9 in)
- Known intolerance to titanium, titanium nitride, parylene C, primer for parylene C, polyether ether ketone, siloxane, nitinol, platinum, iridium, liquid silicone rubber, silicone medical adhesive, and heparin or sensitivity to contrast medical which cannot be adequately premedicated.

As per the FDA label, the Micra Model MC1VR01 pacemaker should not be used in patients for whom a single dose of 1.0 mg dexamethasone acetate cannot be tolerated because the device contains a molded and cured mixture of dexamethasone acetate with the target dosage of 272 µg dexamethasone acetate. It is intended to deliver the steroid to reduce inflammation and fibrosis.

For the MRI contraindications for patients with a Micra MRI device, refer to the Medtronic MRI Technical Manual.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Leadless Cardiac Pacemakers

Policy # 00688

Original Effective Date: 01/01/2020

Current Effective Date: 11/09/2020

Inclusion criteria for clinical trial included patients who were at least 18 years of age or older and had a Class I or II indication for implantation of a single chamber ventricular pacemaker according to ACC/AHA/ HRS 2008 guidelines.

Exclusion criteria included patients who had neurostimulator or any other chronically implanted device that used current, to avoid any possible electrical interference with Micra device. Patients with a mechanical tricuspid valve, implanted vena cava filter, or left ventricular assist device (LVAD) were excluded to avoid electrical or mechanical interference with placing Micra device. Morbidly obese patients were excluded (when telemetry communication of 5 inches or less could not be obtained) to ensure ability to communicate with programmer.

### **Rationale/Source**

Pacemakers are intended to be used as a substitute for the heart's intrinsic pacing system to correct cardiac rhythm disorders. Conventional pacemakers consist of two components: a pulse generator and electrodes (or leads). Pacemakers are considered life-sustaining, life-supporting class III devices for patients with a variety of brady arrhythmias. Even though the efficacy and safety profile of conventional pacemakers are excellent, in a small proportion of patients, they may result in lead complications and the requirement for a surgical pocket. Further, some patients are medically ineligible for conventional pacemakers due to lack of venous access and recurrent infection. Leadless pacemakers are single-unit devices that are implanted in the heart via femoral access, thereby eliminating the potential for complications as a result of leads and surgical pocket. The Micra transcatheter pacing system is the only commercially available leadless pacemaker in the U. S. approved by the FDA.

For individuals with a guidelines-based indication for a ventricular pacing system who are medically eligible for a conventional pacing system who receive a Micra transcatheter pacing system, the evidence includes a pivotal prospective cohort study and a post approval prospective cohort study. The relevant outcomes are overall survival, disease-specific survival, and treatment-related mortality and morbidity. Results at 6 months and 1 year for the pivotal study reported high procedural success (>99%) and device effectiveness (pacing capture threshold met in 98% patients). Most of the system- or procedural-related complications occurred within 30 days. At one year, the incidence of major complication did not increase substantially from six months (3.5% at six months vs 4% at one year). Results of the post approval study were consistent with a pivotal study and showed a lower incidence of major complications up to 30 days post implantation as well as 1 year (1.5% and 2.7%, respectively). In both studies, the point estimates

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.





# Louisiana

## Leadless Cardiac Pacemakers

Policy # 00688

Original Effective Date: 01/01/2020

Current Effective Date: 11/09/2020

of major complications were lower than the pooled estimates from six studies of conventional pacemakers used as a historical comparator. While Micra device eliminates lead- and surgical pocket-related complications, its use can result in potentially more serious complications related to implantation and release of the device (traumatic cardiac injury) and less serious complications related to the femoral access site (groin hematomas, access site bleeding). Considerable uncertainties and unknowns remain in terms of the durability of device and device end-of-life issues. Early and limited experience has suggested that retrieval of these devices is unlikely because in due course, the devices will be encapsulated. There are limited data on device-device interactions (both electrical and mechanical), which may occur when there is a deactivated Micra device alongside another leadless pacemaker or when a leadless pacemaker and trans venous device are both present. While the current evidence is encouraging, overall benefit with the broad use of Micra transcatheter pacing system compared with conventional pacemakers has not been shown. The evidence is insufficient to determine the effects of technology on health outcomes.

For individuals with a guidelines-based indication for a ventricular pacing system who are medically ineligible for a conventional pacing system who receive a Micra transcatheter pacing system, the evidence includes subgroup analysis of a pivotal prospective cohort study and a post approval prospective cohort study. The relevant outcomes are overall survival, disease-specific survival, and treatment-related mortality and morbidity. Information on the outcomes in the subgroup of patients from the post approval study showed that the Micra device was successfully implanted in 98% of cases and safety outcomes were similar to the original cohort. Even though the evidence is limited and long-term effectiveness and safety are unknown, the short-term benefits outweigh the risks because the complex trade-off of adverse events for these devices needs to be assessed in the context of the life-saving potential of pacing systems for patients, ineligible for conventional pacing systems. There are little data available regarding outcomes associated with other alternatives to conventional pacemaker systems such as epicardial leads or trans iliac placement. Epicardial leads are most relevant for the patient who is already going to have a thoracotomy for treatment of their underlying condition (e.g., congenital heart disease). Epicardial leads are associated with a longer intensive care unit stay, more blood loss, and longer ventilation times compared to conventional pacemaker systems. The evidence for trans iliac placement is limited to small case series and the incidence of atrial lead dislodgement using this approach in the literature ranged from 7 to 21%. Limitations in the published evidence preclude determining the effects of the technology on net health outcome.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

Leadless Cardiac Pacemakers

Policy # 00688

Original Effective Date: 01/01/2020

Current Effective Date: 11/09/2020

## **Supplemental Information**

### **Clinical Input from Physician Specialty Societies and Academic Medical Centers**

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

### **2019**

Clinical input was sought to help determine whether the use of leadless cardiac pacemakers for individuals with a guidelines-based indication for a ventricular pacing system would provide a clinically meaningful improvement in net health outcome and whether the use is consistent with generally accepted medical practice. In response to requests, clinical input was received from 2 respondents, including 1 specialty society-level response and 1 physician-level response identified through specialty societies including physicians with academic medical center affiliations.

For individuals with a guidelines-based indication for a ventricular pacing system who are medically ineligible for a conventional pacing system who receive a Micra transcatheter pacing system, clinical input supports this use provides a clinically meaningful improvement in net health outcomes and indicates this use is consistent with generally accepted medical practice in a subgroup of appropriately selected patients when both conditions below are met:

- The patient has symptomatic paroxysmal or permanent high-grade arteriovenous block or symptomatic bradycardia-tachycardia syndrome or sinus node dysfunction (sinus bradycardia or sinus pauses).
- The patient has a significant contraindication precluding placement of conventional single-chamber ventricular pacemaker leads such as any of the following:
  - History of an endovascular or CIED infection or who are very high-risk for infection
  - Limited access for transvenous pacing given venous anomaly, occlusion of axillary veins or planned use of such veins for a semi-permanent catheter or current or planned use of an AV fistula for hemodialysis
  - Presence of a bioprosthetic tricuspid valve

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Leadless Cardiac Pacemakers

Policy # 00688

Original Effective Date: 01/01/2020

Current Effective Date: 11/09/2020

### **Practice Guidelines and Position Statements**

#### **Heart Rhythm Society**

In 2020, the Heart Rhythm Society (HRS), along with the International Society for Cardiovascular Infectious Diseases (ISCVID) and several other Asian, European and Latin American societies, endorsed the European Heart Rhythm Association (EHRA) international consensus document on how to prevent, diagnose, and treat cardiac implantable electronic device infections. The consensus states that for patients at high risk of device-related infections, avoiding a transvenous system, and implanting an epicardial system, may be preferential. It makes the following statements regarding leadless pacemakers:

- 'There is hope that 'leadless' pacemakers will be less prone to infection and can be used in a similar manner [as epicardial systems] in high-risk patients.'
- 'In selected high-risk patients, the risk of infection with leadless pacemakers appears low. The device also seems safe and feasible in patients with pre-existing CIED infection and after extraction of infected leads.'

#### **U.S. Preventive Services Task Force Recommendations**

Not applicable.

#### **Medicare National Coverage**

The Centers for Medicare & Medicaid (CMS) cover leadless pacemakers under coverage with evidence development criteria when procedures are performed in “prospective longitudinal studies” approved by the U.S. FDA using “leadless pacemakers ... in accordance with the FDA approved label for devices that have either:

- An associated ongoing FDA approved post-approval study; or
- Completed an FDA post-approval study.

Each study must be approved by CMS and as a fully-described, written part of its protocol, must address the following research questions:

- What are the peri-procedural and post-procedural complications of leadless pacemakers?
- What are the long term outcomes of leadless pacemakers?
- What are the effects of patient characteristics (age, gender, comorbidities) on the use and health effects of leadless pacemakers?”

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Leadless Cardiac Pacemakers

Policy # 00688

Original Effective Date: 01/01/2020

Current Effective Date: 11/09/2020

The following 2 studies are currently approved by CMS: (1) The Micra CED Study (NCT03039712); CMS approval date: 03/09/17; and (2) Micra Transcatheter Pacing System Post-Approval Registry (NCT02536118); CMS approval date: 02/09/17.

### Ongoing and Unpublished Clinical Trials

Some currently ongoing and unpublished trials that might influence this review are listed in Table 2.

**Table 2. Summary of Key Trials**

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT03039712	Longitudinal Coverage With Evidence Development Study on Micra Leadless Pacemakers (Micra CED)	37,000	Jun 2021
NCT02610673 <sup>a</sup>	WiCS-LV Post Market Surveillance Registry	100	Nov 2021
NCT02051972 <sup>a</sup>	Nanostim Study for a Leadless Cardiac Pacemaker System	1000	Mar 2024
NCT02536118 <sup>a</sup>	Micra Transcatheter Pacing System Post-Approval Registry	3100	Aug 2026

NCT: national clinical trial.

<sup>a</sup> Denotes industry-sponsored or cosponsored trial.

## References

1. Blue Cross and Blue Shield Association, Medical Policy Reference Manual, “Leadless Cardiac Pacemakers”, 2.02.32, August 2020.
2. Reddy VY, Exner DV, Cantillon DJ, et al. Percutaneous implantation of an entirely intracardiac leadless pacemaker. N Engl J Med. Sep 17 2015;373(12):1125-1135. PMID 26321198.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Leadless Cardiac Pacemakers

Policy # 00688

Original Effective Date: 01/01/2020

Current Effective Date: 11/09/2020

3. Udo EO, Zuithoff NP, van Hemel NM, et al. Incidence and predictors of short- and long-term complications in pacemaker therapy: the FOLLOWPACE study. *Heart Rhythm*. May 2012;9(5):728-735. PMID 22182495.
4. Haight, PP, Stewart, RR, Saarel, EE, Pettersson, GG, Najm, HH, Aziz, PP. Lateral thoracotomy for epicardial pacemaker placement in patients with congenital heart disease. *Interact Cardiovasc Thorac Surg*, 2018 Jan 5;26(5). PMID 29300890.
5. Cohen, MM, Bush, DD, Vetter, VV, Tanel, RR, Wieand, TT, Gaynor, JJ, Rhodes, LL. Permanent epicardial pacing in pediatric patients: seventeen years of experience and 1200 outpatient visits. *Circulation*, 2001 May 31;103(21). PMID 11382728.
6. Doll, NN, Piorkowski, CC, Czesla, MM, Kallenbach, MM, Rastan, AA, Arya, AA, Mohr, FF. Epicardial versus transvenous left ventricular lead placement in patients receiving cardiac resynchronization therapy: results from a randomized prospective study. *Thorac Cardiovasc Surg*, 2008 Jul 11;56(5). PMID 18615370.
7. Harake, DD, Shannon, KK, Aboulhosn, JJ, Moore, JJ. Transvenous pacemaker implantation after the bidirectional Glenn operation for patients with complex congenital disease. *J. Cardiovasc. Electrophysiol.*, 2017 Dec 15;29(3). PMID 29240293.
8. Tsutsumi, K. , Hashizume, K. , Kimura, N. , Taguchi, S. , Inoue, Y. , Kashima, I. and Takahashi, R. (2010), Permanent Pacemaker Implantation via the Iliac Vein: An Alternative in 4 Cases with Contraindications to the Pectoral Approach. *Journal of Arrhythmia*, 26: 55-61. doi:10.1016/S1880-4276(10)80037-7.
9. Healey JS, Toff WD, Lamas GA, et al. Cardiovascular outcomes with atrial-based pacing compared with ventricular pacing: meta-analysis of randomized trials, using individual patient data. *Circulation*. Jul 4 2006;114(1):11-17. PMID 16801463.
10. Kirkfeldt RE, Johansen JB, Nohr EA, et al. Risk factors for lead complications in cardiac pacing: a population-based cohort study of 28,860 Danish patients. *Heart Rhythm*. Oct 2011;8(10):1622-1628. PMID 21699827.
11. Kirkfeldt RE, Johansen JB, Nohr EA, et al. Complications after cardiac implantable electronic device implantations: an analysis of a complete, nationwide cohort in Denmark. *Eur Heart J*. May 2014;35(18):1186-1194. PMID 24347317.
12. Food and Drug Administration. FDA Executive Summary Memorandum. General Issues: Leadless Pacemaker Devices Prepared for the February 18, 2016 meeting of the Circulatory System Devices Advisory Panel Gaithersburg Hilton; Gaithersburg, MD. 2016; <https://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/MedicalDevices/MedicalDevicesAdvisoryCommittee/CirculatorySystemDevicesPanel/UCM485093.pdf>.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Leadless Cardiac Pacemakers

Policy # 00688

Original Effective Date: 01/01/2020

Current Effective Date: 11/09/2020

13. American Heart Association. Statement of the American Heart Association to the Food and Drug Administration Circulatory System Devices Panel February 18, 2016: Leadless Cardiac Pacemaker Devices. 2016; <https://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/MedicalDevices/MedicalDevicesAdvisoryCommittee/CirculatorySystemDevicesPanel/UCM486235.pdf>.
14. Reddy VY, Miller MA, Knops RE, et al. Retrieval of the leadless cardiac pacemaker: a multicenter experience. *Circ Arrhythm Electrophysiol.* Dec 2016;9(12). PMID 27932427.
15. Reddy VY, Knops RE, Sperzel J, et al. Permanent leadless cardiac pacing: results of the LEADLESS trial. *Circulation.* Apr 8 2014;129(14):1466-1471. PMID 24664277.
16. Knops RE, Tjong FV, Neuzil P, et al. Chronic performance of a leadless cardiac pacemaker: 1-year follow-up of the LEADLESS trial. *J Am Coll Cardiol.* Apr 21 2015;65(15):1497-1504. PMID 25881930.
17. Lakkireddy D, Knops R, Atwater B, et al. A worldwide experience of the management of battery failures and chronic device retrieval of the Nanostim leadless pacemaker. *Heart Rhythm.* Dec 2017;14(12):1756-1763. PMID 28705736.
18. Tjong FVY, Knops RE, Neuzil P, et al. Midterm safety and performance of a leadless cardiac pacemaker: 3-year follow-up to the LEADLESS Trial (Nanostim Safety and Performance Trial for a Leadless Cardiac Pacemaker System). *Circulation.* Feb 6 2018;137(6):633-635. PMID 28705736.
19. Sperzel, JJ, Defaye, PP, Delnoy, PP, Garcia Guerrero, JJ, Knops, RR, Tondo, CC, Deharo, JJ, Wong, TT, Neuzil, PP. Primary safety results from the LEADLESS Observational Study. *Europace.* 2018 Jan 25;20(9). PMID 29365073.
20. Zuckerman B, Shein M, Paulsen J, et al. Circulatory System Devices Panel Meeting: Leadless Pacemakers. FDA Presentation. 2016; <https://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/MedicalDevices/MedicalDevicesAdvisoryCommittee/CirculatorySystemDevicesPanel/UCM486733.pdf>.
21. Ritter P, Duray GZ, Zhang S, et al. The rationale and design of the Micra Transcatheter Pacing Study: safety and efficacy of a novel miniaturized pacemaker. *Europace.* May 2015;17(5):807-813. PMID 25855677.
22. Ritter P, Duray GZ, Steinwender C, et al. Early performance of a miniaturized leadless cardiac pacemaker: the Micra Transcatheter Pacing Study. *Eur Heart J.* Oct 1 2015;36(37):2510-2519. PMID 26045305.
23. Tjong, FF, Beurskens, NN, de Groot, JJ, Waweru, CC, Liu, SS, Ritter, PP, Reynolds, DD, Wilde, AA, Knops, RR. Health-related quality of life impact of a transcatheter pacing system. *J. Cardiovasc. Electrophysiol.*, 2018 Sep 1;29(12). PMID 30168233.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Leadless Cardiac Pacemakers

Policy # 00688

Original Effective Date: 01/01/2020

Current Effective Date: 11/09/2020

24. Reynolds D, Duray GZ, Omar R, et al. A leadless intracardiac transcatheter pacing system. *N Engl J Med*. Feb 11 2016;374(6):533-541. PMID 26551877.
25. Lloyd M, Reynolds D, Sheldon T, et al. Rate adaptive pacing in an intracardiac pacemaker. *Heart Rhythm*. Feb 2017;14(2):200-205. PMID 27871854.
26. Food and Drug Administration. Summary of Safety and Effectiveness Data (SSED): Micra Transcatheter Pacing System (PMS P150033). 2016; [https://www.accessdata.fda.gov/cdrh\\_docs/pdf15/P150033B.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf15/P150033B.pdf).
27. Transcript of the United States of America Department of Health and Human Services Food and Drug Administration Center for Devices and Radiological Health Medical Devices Advisory Committee: Circulatory System Devices Panel Meeting. February 18, 2016. <https://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/MedicalDevices/MedicalDevicesAdvisoryCommittee/CirculatorySystemDevicesPanel/UCM489547.pdf>.
28. Medtronic. Meet Micra (brochure). n.d.; [http://www.medtronic.com/content/dam/medtronic-com/01\\_crhf/brady/pdfs/medtronic-micra-transcatheter-pacing-system-hcp-brochure.pdf](http://www.medtronic.com/content/dam/medtronic-com/01_crhf/brady/pdfs/medtronic-micra-transcatheter-pacing-system-hcp-brochure.pdf).
29. Grubman, EE, Ritter, PP, Ellis, CC, Giocondo, MM, Augostini, RR, Neuzil, PP, Ravindran, BB, Patel, AA, Omdahl, PP, Pieper, KK, Stromberg, KK, Hudnall, JJ, Reynolds, DD. To retrieve, or not to retrieve: System revisions with the Micra transcatheter pacemaker. *Heart Rhythm*, 2017 Jul 18;14(12). PMID 28713024.
30. Roberts PR, Clementy N, Al Samadi F, et al. A leadless pacemaker in the real-world setting: The Micra Transcatheter Pacing System Post-Approval Registry. *Heart Rhythm*. Sep 2017;14(9):1375-1379. PMID 28502871.
31. El-Chami MF, Brock Johansen J, Zaidi A, et al. Leadless Pacemaker Implant in Patients with Pre-Existing Infections: Results from the Micra Post-Approval Registry. Paper presented at: Heart Rhythm Scientific Sessions. 2018 May 10; Boston, MA.
32. El-Chami, MM, Al-Samadi, FF, Clementy, NN, Garweg, CC, Martinez-Sande, JJ, Piccini, JJ, Iacopino, SS, Lloyd, MM, Viñolas Prat, XX, Jacobsen, MM, Ritter, PP, Johansen, JJ, Tondo, CC, Liu, FF, Fagan, DD, Eakley, AA, Roberts, PP. Updated performance of the Micra transcatheter pacemaker in the real-world setting: A comparison to the investigational study and a transvenous historical control. *Heart Rhythm*, 2018 Aug 14;15(12). PMID 30103071.
33. Duray GZ, Ritter P, El-Chami M, et al. Long-term performance of a transcatheter pacing system: 12-Month results from the Micra Transcatheter Pacing Study. *Heart Rhythm*. May 2017;14(5):702-709. PMID 28192207.
34. El-Chami, MM, Johansen, JJ, Zaidi, AA, Faerstrand, SS, Reynolds, DD, Garcia-Seara, JJ, Mansourati, JJ, Pasquie, JJ, McElderry, HH, Roberts, PP, Soejima, KK, Stromberg, KK, Piccini, JJ.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Leadless Cardiac Pacemakers

Policy # 00688

Original Effective Date: 01/01/2020

Current Effective Date: 11/09/2020

- Leadless pacemaker implant in patients with pre-existing infections: Results from the Micra postapproval registry. *J. Cardiovasc. Electrophysiol.*, 2019 Jan 21;30(4). PMID 30661279.
35. Epstein AE, DiMarco JP, Ellenbogen KA, et al. 2012 ACCF/AHA/HRS focused update incorporated into the ACCF/AHA/HRS 2008 guidelines for device-based therapy of cardiac rhythm abnormalities: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines and the Heart Rhythm Society. *J Am Coll Cardiol.* Jan 22 2013;61(3):e6-75. PMID 23265327.
  36. Gillis AM, Russo AM, Ellenbogen KA, et al. HRS/ACCF expert consensus statement on pacemaker device and mode selection. *J Am Coll Cardiol.* Aug 14 2012;60(7):682-703. PMID 22854177.
  37. Centers for Medicare & Medicaid Services. Decision Memo for Leadless Pacemakers (CAG-00448N). 2017; <https://www.cms.gov/medicare-coverage-database/details/nca-decision-memo.aspx?NCAId=285&bc=ACAAAAAAQAAA&>.

### Policy History

Original Effective Date: 01/01/2020

Current Effective Date: 11/09/2020

10/03/2019 Medical Policy Committee review

10/09/2019 Medical Policy Implementation Committee approval. New policy.

10/01/2020 Medical Policy Committee review

10/07/2020 Medical Policy Implementation Committee approval. No change to coverage.

Next Scheduled Review Date: 10/2021

### Coding

*The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)‡, copyright 2019 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.*

*The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Blue Cross*

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.





# Louisiana

## Leadless Cardiac Pacemakers

Policy # 00688

Original Effective Date: 01/01/2020

Current Effective Date: 11/09/2020

*Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.*

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	33274, 33275
HCPCS	No codes
ICD-10 Diagnosis	All related diagnoses

\*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
  - 1. Consultation with the Blue Cross and Blue Shield Association technology assessment program (TEC) or other nonaffiliated technology evaluation center(s);

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Leadless Cardiac Pacemakers

Policy # 00688

Original Effective Date: 01/01/2020

Current Effective Date: 11/09/2020

2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
3. Reference to federal regulations.

‡ Indicated trademarks are the registered trademarks of their respective owners.

**NOTICE:** If the Patient's health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

**NOTICE:** Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.