



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

*Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.*

### When Services Are Eligible for Coverage

*Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:*

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Based on review of available data, the Company may consider a liver transplant using a cadaver or living donor, for carefully selected patients with end-stage liver failure due to irreversibly damaged livers to be **eligible for coverage**.\*\*

Etiologies of end-stage liver disease include, but are not limited to, the following:

*A. Hepatocellular diseases*

- Alcoholic liver disease
- Viral hepatitis (either A, B, C, or non-A, non-B)
- Autoimmune hepatitis
- Alpha-1 antitrypsin deficiency
- Hemochromatosis
- Nonalcoholic steatohepatitis (NASH)
- Protoporphyrria
- Wilson's disease

*B. Cholestatic liver diseases*

- Primary biliary cirrhosis
- Primary sclerosing cholangitis with development of secondary biliary cirrhosis
- Biliary atresia

*C. Vascular disease*

- Budd-Chiari syndrome

*D. Primary hepatocellular carcinoma (HCC)*

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

(See Policy Guidelines section for patient selection criteria)

E. *Inborn errors of metabolism*

F. *Trauma and toxic reactions*

G. *Miscellaneous*

- Familial amyloid polyneuropathy

Based on review of available data, the Company may consider liver transplantation in patients with polycystic disease of the liver who have massive hepatomegaly causing obstruction or functional impairment to be **eligible for coverage**.\*\*

Based on review of available data, the Company may consider liver transplantation in patients with unresectable hilar cholangiocarcinoma (CCA) to be **eligible for coverage**\*\* (See Policy Guidelines section for patient selection criteria).

Based on review of available data, the Company may consider liver transplantation in pediatric patients with nonmetastatic hepatoblastoma to be **eligible for coverage**.\*\*

Based on review of available data, the Company may consider liver *retransplantation* to be **eligible for coverage**\*\* in patients with:

- Primary graft nonfunction
- Hepatic artery thrombosis
- Chronic rejection
- Ischemic type biliary lesions after donation after cardiac death
- Recurrent nonneoplastic disease causing late graft failure

Based on review of available data, the Company may consider combined liver-kidney transplantation (CLKT) in patients who qualify for liver transplantation and have advanced irreversible kidney disease to be **eligible for coverage**.\*\*

## When Services Are Considered Not Medically Necessary

Based on review of available data, the Company may consider combined liver-kidney transplantation (CLKT) in patients who qualify for liver transplantation and have advanced irreversible kidney disease to be **not medically necessary**\*\* in the following patients:

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

- Patients with HCC that has extended beyond the liver (See Policy Guidelines section for patient selection criteria).
- Patients with ongoing alcohol and/or drug abuse. (Evidence for abstinence may vary among liver transplant programs, but generally a minimum of 3 months is required.)

## When Services Are Considered Investigational

*Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.*

Based on review of available data, the Company considers liver transplantation to be **investigational\*** in the following situations:

- Patients with intrahepatic cholangiocarcinoma (CCA)
- Patients with neuroendocrine tumors (NETs) metastatic to the liver

Based on review of available data, the Company considers liver transplantation in all other situations not described above to be **investigational.\***

## Policy Guidelines

### **Contraindications**

Potential contraindications for solid organ transplant are subject to the judgment of the transplant center and include the following:

- Known current malignancy, including metastatic cancer
- Recent malignancy with high risk of recurrence
- Untreated systemic infection making immunosuppression unsafe, including chronic infection
- Other irreversible end-stage diseases not attributed to liver disease
- History of cancer with a moderate risk of recurrence
- Systemic disease that could be exacerbated by immunosuppression
- Psychosocial conditions or chemical dependency affecting ability to adhere to therapy.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

### **Liver-Specific Criteria**

The Model for End-stage Liver Disease (MELD) and Pediatric End-stage Liver Disease (PELD) scores range from 6 (less ill) to 40 (gravely ill). The MELD and PELD scores will change during a patient's tenure on the waiting list.

Patients with liver disease related to alcohol or drug abuse must be actively involved in a substance abuse treatment program.

Tobacco consumption is a contraindication. Documentation should include confirmation of patient enrollment into smoking cessation program and treatment plan if patient continues to smoke despite interventions.

Smoking remains an independent and significant risk factor for post-transplant morbidity and mortality. Moreover, long-term morbidity and mortality are also linked to continued smoking, which accelerates cardiovascular and cancer risk.

Some authorities have advocated removing patients from the transplant waiting list who continue to smoke despite these interventions. Roughly 20 percent of transplant centers report that they will refuse to list patients due to smoking. One report found that up to 40 percent of patients who had undergone transplantation for alcoholic liver disease resumed smoking early in the post-transplant course, underscoring the need for continued counseling and monitoring after transplantation.

Patients with polycystic disease of the liver do not develop liver failure but may require transplantation due to the anatomic complications of a hugely enlarged liver. The MELD and PELD score may not apply to these cases. One of the following complications should be present:

- Enlargement of liver impinging on respiratory function
- Extremely painful enlargement of liver
- Enlargement of liver significantly compressing and interfering with function of other abdominal organs.

Patients with familial amyloid polyneuropathy do not experience liver disease per se, but develop polyneuropathy and cardiac amyloidosis due to the production of a variant transthyretin molecule by the liver. MELD and PELD exception criteria and scores may apply to these cases. Candidacy

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

for liver transplant is an individual consideration based on the morbidity of the polyneuropathy. Many patients may not be candidates for liver transplant alone due to coexisting cardiac disease.

### **Hepatocellular Carcinoma**

Criteria used for patient selection of hepatocellular carcinoma (HCC) patients eligible for liver transplant include the Milan criteria, which is considered the criterion standard, the University of California, San Francisco expanded criteria, and United Network of Organ Sharing (UNOS) criteria.

#### **Milan Criteria**

A single tumor 5 cm or less or 2 to 3 tumors 3 cm or less.

#### **University of California, San Francisco Expanded Criteria**

A single tumor 6.5 cm or less or up to 3 tumors 4.5 cm or less, and a total tumor size of 8 cm or less.

#### **UNOS Stage T2 Criteria**

A single tumor 2 cm or greater and up to 5 cm or less or 2 to 3 tumors 1 cm or greater and up to 3 cm or less and without extrahepatic spread or macrovascular invasion. UNOS criteria were updated in 2018 ([https://optn.transplant.hrsa.gov/media/1200/optn\\_policies.pdf#nameddest=Policy\\_09](https://optn.transplant.hrsa.gov/media/1200/optn_policies.pdf#nameddest=Policy_09))

Patients with HCC are appropriate candidates for liver transplant only if the disease remains confined to the liver. Therefore, the patient should be periodically monitored while on the waiting list, and if metastatic disease develops, the patient should be removed from the transplant waiting list. Also, at the time of transplant, a backup candidate should be scheduled. If locally extensive or metastatic cancer is discovered at the time of exploration before hepatectomy, the transplant should be aborted, and the backup candidate scheduled for transplant.

Note that liver transplantation for those with T3 HCC is not prohibited by UNOS guidelines, but such patients do not receive any priority on the waiting list. All patients with HCC awaiting transplantation are reassessed at 3-month intervals. Those whose tumors have progressed and are no longer stage T2 will lose the additional allocation points.

Additionally, nodules identified through imaging of cirrhotic livers are given a class 5 designation. Class 5B and 5T nodules are eligible for automatic priority. Class 5B criteria consist of a single nodule 2 cm or larger and up to 5 cm (T2 stage) that meets specified imaging criteria. Class 5T

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

nodules have undergone subsequent locoregional treatment after being automatically approved on initial application or extension. A single class 5A nodule (>1 cm and <2 cm) corresponds to T1 HCC and does not qualify for automatic priority. However, combinations of class 5A nodules are eligible for automatic priority if they meet stage T2 criteria. Class 5X lesions are outside of stage T2 and ineligible for automatic exception points. Nodules less than 1 cm are considered indeterminate and are not considered for additional priority. Therefore, the UNOS allocation system provides strong incentives to use locoregional therapies to downsize tumors to T2 status and to prevent progression while on the waiting list.

### **Cholangiocarcinoma**

According to the Organ Procurement and Transplantation Network (OPTN) policy on liver allocation, candidates with cholangiocarcinoma meeting the following criteria will be eligible for a MELD or PELD exception with a 10% mortality equivalent increase every 3 months:

- Centers must submit a written protocol for patient care to the OPTN and UNOS Liver and Intestinal Organ Transplantation Committee before requesting a MELD score exception for a candidate with cholangiocarcinoma. This protocol should include selection criteria, administration of neoadjuvant therapy before transplantation, and operative staging to exclude patients with regional hepatic lymph node metastases, intrahepatic metastases, and/or extrahepatic disease. The protocol should include data collection as deemed necessary by the OPTN and UNOS Liver and Intestinal Organ Transplantation Committee.
- Candidates must satisfy diagnostic criteria for hilar cholangiocarcinoma: malignant-appearing stricture on cholangiography and one of the following: carbohydrate antigen 19-9 100 U/mL, or and biopsy or cytology results demonstrating malignancy, or aneuploidy. The tumor should be considered unresectable on the basis of technical considerations or underlying liver disease (eg, primary sclerosing cholangitis).
- If cross-sectional imaging studies (computed tomography scan, ultrasound, magnetic resonance imaging) demonstrate a mass, the mass should be 3 cm or less.
- Intra- and extrahepatic metastases should be excluded by cross-sectional imaging studies of the chest and abdomen at the time of initial exception and every 3 months before score increases.
- Regional hepatic lymph node involvement and peritoneal metastases should be assessed by operative staging after completion of neoadjuvant therapy and before liver transplantation.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

Endoscopic ultrasound-guided aspiration of regional hepatic lymph nodes may be advisable to exclude patients with obvious metastases before neoadjuvant therapy is initiated.

- Transperitoneal aspiration or biopsy of the primary tumor (either by endoscopic ultrasound, operative, or percutaneous approaches) should be avoided because of the high risk of tumor seeding associated with these procedures.

### **Living Donor Criteria**

Donor morbidity and mortality are prime concerns in donors undergoing right lobe, left lobe, or left lateral segment donor partial hepatectomy as part of living donor liver transplantation. Partial hepatectomy is a technically demanding surgery, the success of which may be related to the availability of an experienced surgical team. The American Society of Transplant Surgeons proposed the following guidelines for living donors (American Society of Transplant Surgeons: Ethics Committee. American Society of Transplant Surgeons' position paper on adult-to-adult living donor liver transplantation. *Liver Transplant*. 2000;6(6):815-817. PMID 11084076):

- They should be healthy individuals who are carefully evaluated and approved by a multidisciplinary team including hepatologists and surgeons to assure that they can tolerate the procedure
- They should undergo evaluation to ensure that they fully understand the procedure and associated risks
- They should be of legal age and have sufficient intellectual ability to understand the procedures and give informed consent
- They should be emotionally related to the recipients
- They must be excluded if the donor is felt or known to be coerced
- They need to have the ability and willingness to comply with long-term follow-up.

### **Background/Overview**

Solid organ transplantation offers a treatment option for patients with different types of end stage organ failure that can be lifesaving or provide significant improvements to a patient's quality of life. Many advances have been made in the last several decades to reduce perioperative complications. Available data supports improvement in long-term survival as well as improved quality of life particularly for liver, kidney, pancreas, heart, and lung transplants. Allograft rejection remains a key early and late complication risk for any organ transplantation. Transplant recipients require life-long immunosuppression to prevent rejection. Patients are prioritized for transplant by mortality risk and

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

severity of illness criteria developed by Organ Procurement and Transplantation Network and United Network of Organ Sharing.

### **Liver transplantation**

Liver transplantation is routinely performed as a treatment of last resort for patients with end-stage liver disease. Liver transplantation may be performed with liver donation after a brain or cardiac death or with a liver segment donation from a living donor. Certain populations are prioritized as Status 1A (eg, acute liver failure with a life expectancy of fewer than 7 days without a liver transplant) or Status 1B (pediatric patients with chronic liver disease). Following Status 1, donor livers are prioritized to those with the highest scores on the Model for End-stage Liver Disease (MELD) and Pediatric End-stage Liver Disease (PELD) scales. Due to the scarcity of donor livers, a variety of strategies have been developed to expand the donor pool. For example, a split graft refers to dividing a donor liver into 2 segments that can be used for 2 recipients. Living donor liver transplantation (LDLT) is now commonly performed for adults and children from a related or unrelated donor. Depending on the graft size needed for the recipient, either the right lobe, left lobe, or the left lateral segment can be used for LDLT. In addition to addressing the problem of donor organ scarcity, LDLT allows the procedure to be scheduled electively before the recipient's condition deteriorates or serious complications develop. LDLT also shortens the preservation time for the donor liver and decreases disease transmission from donor to recipient.

## **FDA or Other Governmental Regulatory Approval**

### **U.S. Food and Drug Administration (FDA)**

Solid organ transplants are a surgical procedure and, as such, are not subject to regulation by the U.S. Food and Drug Administration (FDA).

The FDA regulates human cells and tissues intended for implantation, transplantation, or infusion through the Center for Biologics Evaluation and Research, under Code of Federal Regulation Title 21, parts 1270 and 1271. Solid organs used for transplantation are subject to these regulations.

## **Rationale/Source**

Liver transplantation is currently the treatment of last resort for patients with end-stage liver disease. Liver transplantation may be performed with a liver donation after a brain or cardiac death or with a liver segment donation from a living donor. Patients are prioritized for transplant by mortality risk

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.





## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

and severity of illness criteria developed by the Organ Procurement and Transplantation Network and the United Network of Organ Sharing. The severity of illness is determined by the Model for End-stage Liver Disease and Pediatric End-stage Liver Disease scores.

For individuals who have a hepatocellular disease who receive a liver transplant, the evidence includes case series, registry studies, and systematic reviews. Relevant outcomes include overall survival (OS), morbid events, and treatment-related morbidity and mortality. Studies on liver transplantation for viral hepatitis have found that survival is lower than for other liver diseases. Although these statistics raise questions about the most appropriate use of a scarce resource (donor livers), the long-term survival rates are significant in a group of patients who have no other treatment options. Also, survival can be improved by the eradication of the hepatitis virus before transplantation. For patients with nonalcoholic steatohepatitis, OS rates have been shown to be similar to other indications for liver transplantation. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have primary hepatocellular carcinoma who receive a liver transplant, the evidence includes systematic reviews of observational studies. Relevant outcomes include OS, morbid events, and treatment-related morbidity and mortality. In the past, long-term outcomes in patients with primary hepatocellular malignancies had been poor (19%) compared with the OS of liver transplant recipients. However, the recent use of standardized patient selection criteria (eg, the Milan criteria diameter) has dramatically improved OS rates. In the appropriately selected patients, a liver transplant has been shown to result in higher survival rates than resection. In patients who present with unresectable organ-confined disease, transplant represents the only curative approach. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have extrahepatic cholangiocarcinoma who receive a liver transplant, the evidence includes systematic reviews of observational studies. Relevant outcomes include OS, morbid events, and treatment-related morbidity and mortality. For patients with extrahepatic (hilar or perihilar) cholangiocarcinoma who are treated with adjuvant chemotherapy, survival rates have been reported as high as 76%. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

For individuals who have intrahepatic cholangiocarcinoma who receive a liver transplant, the evidence includes registry studies. Relevant outcomes include OS, morbid events, and treatment-related morbidity and mortality. Five-year survival rates after liver transplantation in patients with cholangiocarcinoma are less than 30%. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have metastatic neuroendocrine tumors who receive a liver transplant, the evidence includes systematic reviews of case series. Relevant outcomes include OS, morbid events, and treatment-related morbidity and mortality. In select patients with nonresectable, hormonally active liver metastases refractory to medical therapy, liver transplantation has been considered as an option to extend survival and minimize endocrine symptoms. While some centers may perform liver transplants on select patients with neuroendocrine tumors, the available studies are limited by their heterogeneous populations. Further studies are needed to determine the appropriate selection criteria. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have pediatric hepatoblastoma who receive a liver transplant, the evidence includes case series. Relevant outcomes include OS, morbid events, and treatment-related morbidity and mortality. The literature on liver transplantation for pediatric hepatoblastoma is limited but case series have demonstrated good outcomes and high rates of long-term survival. Additionally, nonmetastatic pediatric hepatoblastoma is among in United Network for Organ Sharing criteria for patients eligible for liver transplantation. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have a failed liver transplant who receive a liver retransplant, the evidence includes observational studies. Relevant outcomes include OS, morbid events, and treatment-related morbidity and mortality. Case series have demonstrated favorable outcomes with liver retransplantation in certain populations, such as when criteria for original liver transplantation are met for retransplantation. While some evidence has suggested outcomes after retransplantation may be less favorable than for initial transplantation in some patients, long-term survival benefits have been demonstrated. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals with indications for liver and kidney transplant who receive a combined liver-kidney transplant, the evidence includes registry studies. Relevant outcomes include OS, morbid events,

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

and treatment-related morbidity and mortality. Most of the evidence involves adults with cirrhosis and kidney failure. Indications for combined liver-kidney transplant in children are rare and often congenital and include liver-based metabolic abnormalities affecting the kidney, along with structural diseases affecting both the liver and kidney. In both adults and children, comparisons with either liver or kidney transplantation alone would suggest that combined liver-kidney transplant is no worse, and possibly better, for graft and patient survival. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

Liver transplant is an accepted treatment of end-stage liver disease that provides a survival benefit in appropriately selected patients and may be considered medically necessary for the indications listed in the Coverage section and inpatients otherwise meeting United Network of Organ Sharing criteria. Liver transplantation is investigational in patients in whom the procedure is expected to be futile due to comorbid disease or in whom posttransplantation care is expected to worsen comorbid conditions significantly. Based on survival data, transplantation in patients with hilar cholangiocarcinoma who meet strict eligibility criteria may be considered medically necessary; transplantation for neuroendocrine tumors metastatic to the liver is considered investigational. Clinical vetting supported retransplantation following primary graft nonfunction, hepatic artery thrombosis, ischemic biliary injury after donation after cardiac death, chronic rejection, or certain recurrent non-neoplastic diseases resulting in end-stage liver failure in a primary transplant. As a result, retransplantation after initially failed liver transplant may be considered medically necessary in these situations.

## **Supplemental Information**

### **Clinical Input From Physician Specialty Societies and Academic Medical Centers**

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

In response to requests, input was received from 3 physician specialty societies and 5 academic medical centers while this policy was under review. There was a consensus among reviewers that liver transplantation may be medically necessary for end-stage liver failure due to irreversibly damaged livers from various disease states such as those considered during the report update. There

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

was also a consensus among reviewers that liver retransplantation is appropriate in patients with acute or chronic liver failure such as primary graft nonfunction, ischemic-type biliary injury after donation after cardiac death, hepatic artery thrombosis, chronic rejection or recurrent diseases such as primary sclerosing cholangitis, autoimmune hepatitis, and hepatitis C resulting in end-stage liver failure. There was general support for the use of liver transplantation as a treatment for cholangiocarcinoma in patients who meet strict eligibility criteria. In general, there was no support for the use of liver transplantation for a neuroendocrine tumor metastatic to the liver.

### **Practice Guidelines and Position Statements**

#### **International Consensus Conference**

The Milan criteria were recommended for use as the benchmark for patient selection, although it was suggested that the Milan criteria might be modestly expanded based on data from expansion studies that demonstrated outcomes are comparable with outcomes from studies using the Milan criteria. Candidates for liver transplantation should also have a predicted survival of 5 years or more. The consensus criteria indicate alpha-fetoprotein concentrations may be used with imaging to assist in determining patient prognosis.

Regarding liver retransplantation, the consensus criteria issued a weak recommendation for retransplantation after graft failure of a living donor transplant for hepatocellular carcinoma (HCC) in patients meeting regional criteria for a deceased donor liver transplant. A strong recommendation was issued against liver retransplantation with a deceased donor for graft failure for patients exceeding regional criteria. Also, the consensus criteria issued a strong recommendation that liver retransplantation for recurrent HCC would not be appropriate. However, a de novo case of HCC may be treated as a new tumor, and retransplantation may be considered even though data to support this is limited.

#### **American Association for the Study of Liver Diseases and American Society of Transplantation**

In 2013, the American Association for the Study of Liver Diseases and the American Society of Transplantation issued joint guidelines on evaluating patients for a liver transplant. These guidelines indicated liver transplantation for severe acute or advanced chronic liver disease after all effective medical treatments have been attempted. The formal evaluation should confirm the irreversible nature of the liver disease and lack of effective alternative medical therapy.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

The guidelines also stated that liver transplant is indicated for the following conditions:

- Acute liver failure complications of cirrhosis
- Liver-based metabolic condition with systemic manifestations
  - $\alpha_1$ -Antitrypsin deficiency
  - Familial amyloidosis
  - Glycogen storage disease
  - Hemochromatosis
  - Primary oxaluria
  - Wilson disease
- Systemic complications of chronic liver disease.

The guidelines also included 1-A recommendations (strong recommendation with high-quality evidence) for a liver transplant that:

- "Tobacco consumption should be prohibited in LT [liver transplant] candidates."
- "Patients with HIV infection are candidates for LT if immune function is adequate and the virus is expected to be undetectable by the time of LT."
- "LT candidates with HCV [hepatitis C virus] have the same indications for LT as for other etiologies of cirrhosis."

Contraindications to liver transplant included:

- "MELD [Model for End-stage Liver Disease] score < 15
- Severe cardiac or pulmonary disease
- AIDS
- Ongoing alcohol or illicit substance abuse
- Hepatocellular carcinoma with metastatic spread
- Uncontrolled sepsis
- Anatomic abnormality that precludes liver transplantation
- Intrahepatic cholangiocarcinoma\
- Extrahepatic malignancy
- Fulminant hepatic failure
- Hemangiosarcoma
- Persistent noncompliance
- Lack of adequate social support system."

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

In 2014, the American Association for the Study of Liver Diseases, the American Society of Transplantation, and the North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition issued joint guidelines on the evaluation of the pediatric patients for liver transplant. The guidelines stated that "disease categories suitable for referral to a pediatric LT program are similar to adults: acute liver failure, autoimmune, cholestasis, metabolic or genetic, oncologic, vascular, and infectious. However, specific etiologies and outcomes differ widely from adult patients, justifying independent pediatric guidelines." The indications listed for liver transplantation included biliary atresia, Alagille syndrome, pediatric acute liver failure, hepatic tumors, HCC, hemangioendothelioma, cystic fibrosis-associated liver disease, urea cycle disorders, immune-mediated liver disease, along with other metabolic or genetic disorders.

In 2019, the American Association for the Study of Liver Diseases guideline on alcohol-associated liver disease provides recommendations on the timing of referral and selection of candidates for liver transplant. The guidance notes that the patient's history of addiction to alcohol is a primary driver in selecting appropriate candidates for liver transplantation. Clinical characteristics that should trigger an evaluation and consideration for liver transplant include decompensated alcohol-associated cirrhosis, Child-Pugh-Turcotte class C cirrhosis, or a MELD-Na score  $\geq 21$ . Additionally, the guideline notes that candidate selection "should not be based solely on a fixed interval of abstinence" and instead a formal psychological evaluation can help stratify patients into higher- or lesser-risk strata for relapse.

### **National Comprehensive Cancer Network**

The NCCN guidelines on hepatobiliary cancers (v.4.2020) recommend referral to a liver transplant center or bridge therapy for patients with HCC meeting United Network of Organ Sharing criteria of a single tumor measuring 2 to 5 cm, or 2 to 3 tumors 3 cm or less with no macrovascular involvement or extrahepatic disease. Patients should be referred to the transplant center. Patients should be referred to the transplant center before the biopsy. In patients who are ineligible for transplant and in select patients with Child-Pugh class A or B liver function with tumors that are resectable, the NCCN indicates resection is the preferred treatment option; locoregional therapy may also be considered. Patients with unresectable HCC should be evaluated for liver transplantation; if the patient is a transplant candidate, then referral to a transplant center should be given or bridge therapy should be considered. The NCCN guidelines on hepatobiliary cancers also indicate that. These are level 2A recommendations based on lower-level evidence and uniform consensus.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

The NCCN guidelines on neuroendocrine tumors (v.1.2019) indicate that liver transplantation for neuroendocrine liver metastases is considered investigational despite "encouraging" 5-year survival rates.

### **U.S. Preventive Services Task Force Recommendations**

Not applicable.

### **Medicare National Coverage**

Medicare covers adult liver transplantation for end-stage liver disease and HCC when performed in a facility approved by the Centers for Medicare & Medicaid Services as meeting institutional coverage criteria for liver transplants. The following conditions must be met for coverage of HCC:

- "The patient is not a candidate for subtotal liver resection;
- The patient's tumor(s) is less than or equal to 5 cm in diameter;
- There is no macrovascular involvement; and
- There is no identifiable extrahepatic spread of tumor to surrounding lymph nodes, lungs, abdominal organs or bone; and
- The transplant is furnished in a facility that is approved by CMS [Centers for Medicare & Medicaid Services]..."

Beginning in June 2012, on review of this national coverage decision for new evidence, Medicare began covering adult liver transplantation, at Medicare administrative contractor discretion, for extrahepatic unresectable cholangiocarcinoma, liver metastases due to a neuroendocrine tumor, and hemangioendothelioma. Adult liver transplantation is excluded from other malignancies.

Pediatric liver transplantation is covered for children (<18 years of age) when performed at pediatric hospitals approved by the Centers for Medicare & Medicaid Services. Coverage includes extrahepatic biliary atresia or any other form of end-stage liver disease, except for children with a malignancy extending beyond the margins of the liver or those with persistent viremia.

### **Ongoing and Unpublished Clinical Trials**

Some currently ongoing and unpublished trials that might influence this review are listed in Table 1.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

**Table 1. Summary of Key Trials**

<b>NCT No.</b>	<b>Trial Name</b>	<b>Planned Enrollment</b>	<b>Completion Date</b>
<i>Ongoing</i>			
NCT03500315	HOPE in Action Prospective Multicenter, Clinical Trial of Deceased HIVD+ Kidney Transplants for HIV+ Recipients	160	Aug 2022
NCT02878473	Liver Transplantation for the Treatment of Early Stages of Intrahepatic Cholangiocarcinoma in Cirrhotics	30	Jan 2029

NCT: national clinical trial.

## **References**

1. Blue Cross and Blue Shield Association, Medical Policy Reference Manual, “Liver Transplant and Combined Liver-Kidney Transplant”, 7.03.06, September 2020.
2. Lucey MR. Liver transplantation in patients with alcoholic liver disease. *Liver Transpl.* 2011;17(7):751.
3. Muñoz SJ. Tobacco use by liver transplant recipients: grappling with a smoking gun. *Liver Transpl.* 2005;11(6):606.
4. Bataller R. Time to ban smoking in patients with chronic liver diseases. *Hepatology.* 2006;44(6):1394
5. Ehlers SL, Rodrigue JR, Widows MR, Reed AI, Nelson DR. Tobacco use before and after liver transplantation: a single center survey and implications for clinical practice and research. *Liver Transpl.* 2004;10(3):412.
6. Black CK, Termanini KM, Aguirre O, et al. Solid organ transplantation in the 21 st century. *Ann Transl Med.* Oct 2018; 6(20): 409. PMID 30498736
7. Belle SH, Beringer KC, Detre KM. An update on liver transplantation in the United States: recipient characteristics and outcome. *Clin Transpl.* 1995: 19-33. PMID 8794252
8. Sheiner P, Rochon C. Recurrent hepatitis C after liver transplantation. *Mt Sinai J Med.* Mar-Apr 2012; 79(2): 190-8. PMID 22499490

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.





# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

9. Gadiparthi C, Cholankeril G, Perumpail BJ, et al. Use of direct-acting antiviral agents in hepatitis C virus-infected liver transplant candidates. *World J Gastroenterol*. Jan 21 2018; 24(3): 315-322. PMID 29391754
10. Wang X, Li J, Riaz DR, et al. Outcomes of liver transplantation for nonalcoholic steatohepatitis: a systematic review and meta-analysis. *Clin Gastroenterol Hepatol*. Mar 2014; 12(3): 394-402.e1. PMID 24076414
11. Cholankeril G, Wong RJ, Hu M, et al. Liver Transplantation for Nonalcoholic Steatohepatitis in the US: Temporal Trends and Outcomes. *Dig Dis Sci*. Oct 2017; 62(10): 2915-2922. PMID 28744836
12. Schoenberg MB, Bucher JN, Vater A, et al. Resection or Transplant in Early Hepatocellular Carcinoma. *Dtsch Arztebl Int*. Aug 07 2017; 114(31-32): 519-526. PMID 28835324
13. Zheng Z, Liang W, Milgrom DP, et al. Liver transplantation versus liver resection in the treatment of hepatocellular carcinoma: a meta-analysis of observational studies. *Transplantation*. Jan 27 2014; 97(2): 227-34. PMID 24142034
14. Guiteau JJ, Cotton RT, Washburn WK, et al. An early regional experience with expansion of Milan Criteria for liver transplant recipients. *Am J Transplant*. Sep 2010; 10(9): 2092-8. PMID 20883543
15. Pomfret EA, Washburn K, Wald C, et al. Report of a national conference on liver allocation in patients with hepatocellular carcinoma in the United States. *Liver Transpl*. Mar 2010; 16(3): 262-78. PMID 20209641
16. Ioannou GN, Perkins JD, Carithers RL. Liver transplantation for hepatocellular carcinoma: impact of the MELD allocation system and predictors of survival. *Gastroenterology*. May 2008; 134(5): 1342-51. PMID 18471511
17. Chan EY, Larson AM, Fix OK, et al. Identifying risk for recurrent hepatocellular carcinoma after liver transplantation: implications for surveillance studies and new adjuvant therapies. *Liver Transpl*. Jul 2008; 14(7): 956-65. PMID 18581511
18. Mazzaferro V, Regalia E, Doci R, et al. Liver transplantation for the treatment of small hepatocellular carcinomas in patients with cirrhosis. *N Engl J Med*. Mar 14 1996; 334(11): 693-9. PMID 8594428
19. Firl DJ, Kimura S, McVey J, et al. Reframing the approach to patients with hepatocellular carcinoma: Longitudinal assessment with hazard associated with liver transplantation for HCC (HALTHCC) improves ablate and wait strategy. *Hepatology*. Oct 2018; 68(4): 1448-1458. PMID 29604231

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

20. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Hepatobiliary Cancers. Version 4.2020.
21. Yadav DK, Chen W, Bai X, et al. Salvage Liver Transplant versus Primary Liver Transplant for Patients with Hepatocellular Carcinoma. *Ann Transplant*. Aug 03 2018; 23: 524-545. PMID 30072683
22. Murali AR, Patil S, Phillips KT, et al. Locoregional Therapy With Curative Intent Versus Primary Liver Transplant for Hepatocellular Carcinoma: Systematic Review and Meta-Analysis. *Transplantation*. Aug 2017; 101(8): e249-e257. PMID 28282359
23. Maggs JR, Suddle AR, Aluvihare V, et al. Systematic review: the role of liver transplantation in the management of hepatocellular carcinoma. *Aliment Pharmacol Ther*. May 2012; 35(10): 1113-34. PMID 22432733
24. Chan DL, Alzahrani NA, Morris DL, et al. Systematic review of efficacy and outcomes of salvage liver transplantation after primary hepatic resection for hepatocellular carcinoma. *J Gastroenterol Hepatol*. Jan 2014; 29(1): 31-41. PMID 24117517
25. Zhu Y, Dong J, Wang WL, et al. Short- and long-term outcomes after salvage liver transplantation versus primary liver transplantation for hepatocellular carcinoma: a meta-analysis. *Transplant Proc*. Nov 2013; 45(9): 3329-42. PMID 24182812
26. Gu J, Bai J, Shi X, et al. Efficacy and safety of liver transplantation in patients with cholangiocarcinoma: a systematic review and meta-analysis. *Int J Cancer*. May 01 2012; 130(9): 2155-63. PMID 21387295
27. Heimbach JK. Successful liver transplantation for hilar cholangiocarcinoma. *Curr Opin Gastroenterol*. May 2008; 24(3): 384-8. PMID 18408469
28. Darwish Murad S, Kim WR, Harnois DM, et al. Efficacy of neoadjuvant chemoradiation, followed by liver transplantation, for perihilar cholangiocarcinoma at 12 US centers. *Gastroenterology*. Jul 2012; 143(1): 88-98.e3; quiz e14. PMID 22504095
29. Heimbach JK, Gores GJ, Haddock MG, et al. Predictors of disease recurrence following neoadjuvant chemoradiotherapy and liver transplantation for unresectable perihilar cholangiocarcinoma. *Transplantation*. Dec 27 2006; 82(12): 1703-7. PMID 17198263
30. Rea DJ, Heimbach JK, Rosen CB, et al. Liver transplantation with neoadjuvant chemoradiation is more effective than resection for hilar cholangiocarcinoma. *Ann Surg*. Sep 2005; 242(3): 451-8; discussion 458-61. PMID 16135931
31. Pascher A, Jonas S, Neuhaus P. Intrahepatic cholangiocarcinoma: indication for transplantation. *J Hepatobiliary Pancreat Surg*. 2003; 10(4): 282-7. PMID 14598146

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

32. Friman S, Foss A, Isoniemi H, et al. Liver transplantation for cholangiocarcinoma: selection is essential for acceptable results. *Scand J Gastroenterol*. Mar 2011; 46(3): 370-5. PMID 21073376
33. Meyer CG, Penn I, James L. Liver transplantation for cholangiocarcinoma: results in 207 patients. *Transplantation*. Apr 27 2000; 69(8): 1633-7. PMID 10836374
34. Robles R, Figueras J, Turrion VS, et al. Spanish experience in liver transplantation for hilar and peripheral cholangiocarcinoma. *Ann Surg*. Feb 2004; 239(2): 265-71. PMID 14745336
35. Casavilla FA, Marsh JW, Iwatsuki S, et al. Hepatic resection and transplantation for peripheral cholangiocarcinoma. *J Am Coll Surg*. Nov 1997; 185(5): 429-36. PMID 9358085
36. Fan ST, Le Treut YP, Mazzaferro V, et al. Liver transplantation for neuroendocrine tumour liver metastases. *HPB (Oxford)*. Jan 2015; 17(1): 23-8. PMID 24992381
37. Mathe Z, Tagkalos E, Paul A, et al. Liver transplantation for hepatic metastases of neuroendocrine pancreatic tumors: a survival-based analysis. *Transplantation*. Mar 15 2011; 91(5): 575-82. PMID 21200365
38. Hamilton EC, Balogh J, Nguyen DT, et al. Liver transplantation for primary hepatic malignancies of childhood: The UNOS experience. *J Pediatr Surg*. Oct 12 2017. PMID 29108844
39. Barrena S, Hernandez F, Miguel M, et al. High-risk hepatoblastoma: results in a pediatric liver transplantation center. *Eur J Pediatr Surg*. Jan 2011; 21(1): 18-20. PMID 20938901
40. Malek MM, Shah SR, Atri P, et al. Review of outcomes of primary liver cancers in children: our institutional experience with resection and transplantation. *Surgery*. Oct 2010; 148(4): 778-82; discussion 782-4. PMID 20728194
41. Browne M, Sher D, Grant D, et al. Survival after liver transplantation for hepatoblastoma: a 2-center experience. *J Pediatr Surg*. Nov 2008; 43(11): 1973-81. PMID 18970927
42. Czauderna P, Otte JB, Aronson DC, et al. Guidelines for surgical treatment of hepatoblastoma in the modern era--recommendations from the Childhood Liver Tumour Strategy Group of the International Society of Paediatric Oncology (SIOPEL). *Eur J Cancer*. May 2005; 41(7): 1031-6. PMID 15862752
43. Organ Procurement and Transplantation Network (OPTN). Policy 9: Allocation of Livers and Liver-Intestines. 2018; [https://optn.transplant.hrsa.gov/media/1200/optn\\_policies.pdf#nameddest=Policy\\_01](https://optn.transplant.hrsa.gov/media/1200/optn_policies.pdf#nameddest=Policy_01). Accessed July 6, 2020.
44. Bellido CB, Martinez JM, Artacho GS, et al. Have we changed the liver retransplantation survival?. *Transplant Proc*. Jul-Aug 2012; 44(6): 1526-9. PMID 22841203
45. Remiszewski P, Kalinowski P, Dudek K, et al. Influence of selected factors on survival after liver retransplantation. *Transplant Proc*. Oct 2011; 43(8): 3025-8. PMID 21996216

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

46. Hong JC, Kaldas FM, Kositamongkol P, et al. Predictive index for long-term survival after retransplantation of the liver in adult recipients: analysis of a 26-year experience in a single center. *Ann Surg.* Sep 2011; 254(3): 444-8; discussion 448-9. PMID 21817890
47. Lunsford KE, Bodzin AS, Markovic D, et al. Avoiding Futility in Simultaneous Liver-kidney Transplantation: Analysis of 331 Consecutive Patients Listed for Dual Organ Replacement. *Ann Surg.* May 2017; 265(5): 1016-1024. PMID 27232249
48. Fong TL, Khemichian S, Shah T, et al. Combined liver-kidney transplantation is preferable to liver transplant alone for cirrhotic patients with renal failure. *Transplantation.* Aug 27 2012; 94(4): 411-6. PMID 22805440
49. Ruiz R, Jennings LW, Kim P, et al. Indications for combined liver and kidney transplantation: propositions after a 23-yr experience. *Clin Transplant.* Nov-Dec 2010; 24(6): 807-11. PMID 20002463
50. Calinescu AM, Wildhaber BE, Poncet A, et al. Outcomes of combined liver-kidney transplantation in children: analysis of the scientific registry of transplant recipients. *Am J Transplant.* Dec 2014; 14(12): 2861-8. PMID 25274400
51. de la Cerda F, Jimenez WA, Gjertson DW, et al. Renal graft outcome after combined liver and kidney transplantation in children: UCLA and UNOS experience. *Pediatr Transplant.* Jun 2010; 14(4): 459-64. PMID 20070563
52. Marcos A, Ham JM, Fisher RA, et al. Single-center analysis of the first 40 adult-to-adult living donor liver transplants using the right lobe. *Liver Transpl.* May 2000; 6(3): 296-301. PMID 10827229
53. Malago M, Testa G, Marcos A, et al. Ethical considerations and rationale of adult-to-adult living donor liver transplantation. *Liver Transpl.* Oct 2001; 7(10): 921-7. PMID 11679994
54. Renz JF, Busuttil RW. Adult-to-adult living-donor liver transplantation: a critical analysis. *Semin Liver Dis.* 2000; 20(4): 411-24. PMID 11200412
55. Bak T, Wachs M, Trotter J, et al. Adult-to-adult living donor liver transplantation using right-lobe grafts: results and lessons learned from a single-center experience. *Liver Transpl.* Aug 2001; 7(8): 680-6. PMID 11510011
56. Shiffman ML, Brown RS, Olthoff KM, et al. Living donor liver transplantation: summary of a conference at The National Institutes of Health. *Liver Transpl.* Feb 2002; 8(2): 174-88. PMID 11862598
57. Grant RC, Sandhu L, Dixon PR, et al. Living vs. deceased donor liver transplantation for hepatocellular carcinoma: a systematic review and meta-analysis. *Clin Transplant.* Jan-Feb 2013; 27(1): 140-7. PMID 23157398

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

58. Cooper C, Kanters S, Klein M, et al. Liver transplant outcomes in HIV-infected patients: a systematic review and meta-analysis with synthetic cohort. *AIDS*. Mar 27 2011; 25(6): 777-86. PMID 21412058
59. Terrault NA, Roland ME, Schiano T, et al. Outcomes of liver transplant recipients with hepatitis C and human immunodeficiency virus coinfection. *Liver Transpl*. Jun 2012; 18(6): 716-26. PMID 22328294
60. Organ Procurement and Transplantation Network (OPTN). Organ Procurement and Transplantation Network Policies. 2018; [https://optn.transplant.hrsa.gov/media/1200/optn\\_policies.pdf](https://optn.transplant.hrsa.gov/media/1200/optn_policies.pdf).
61. Blumberg EA, Rogers CC. Solid organ transplantation in the HIV-infected patient: Guidelines from the American Society of Transplantation Infectious Diseases Community of Practice. *Clin Transplant*. Sep 2019; 33(9): e13499. PMID 30773688
62. American Association for the Study of Liver Diseases, American Society of Transplantation. Liver transplantation, evaluation of the adult patient. 2013; <https://www.aasld.org/publications/practice-guidelines-0>.
63. Squires RH, Ng V, Romero R, et al. Evaluation of the pediatric patient for liver transplantation: 2014 practice guideline by the American Association for the Study of Liver Diseases, American Society of Transplantation and the North American Society for Pediatric Gastroenterology, Hepatology and Nutrition. *Hepatology*. Jul 2014; 60(1): 362-98. PMID 24782219
64. American Association for the Study of Liver Diseases. Diagnosis and Treatment of Alcohol-Associated Liver Diseases: 2019 Practice Guidance From the American Association for the Study of Liver Diseases. <https://aasldpubs.onlinelibrary.wiley.com/doi/full/10.1002/hep.30866>.
65. National Comprehensive Cancer Network. Hepatobiliary Cancers. Version 4.2020. [https://www.nccn.org/professionals/physician\\_gls/PDF/hepatobiliary.pdf](https://www.nccn.org/professionals/physician_gls/PDF/hepatobiliary.pdf).
66. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Neuroendocrine and Adrenal Tumors. Version 1.2019. [https://www.nccn.org/professionals/physician\\_gls/pdf/neuroendocrine.pdf](https://www.nccn.org/professionals/physician_gls/pdf/neuroendocrine.pdf).
67. Centers for Medicare & Medicaid Services. National Coverage Determination (NCD) for Adult Liver Transplantation (260.1). 2012; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCAId=259&NcaName=Liver+Transplantation+for+Malignancies&ExpandComments=n&CommentPeriod=0&NCDId=70&ncdver=3&id=186&bc=gABAAAAAEEAAAA%3D%3D&>.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

68. Centers for Medicare & Medicaid Services. National Coverage Determination (NCD) for Pediatric Liver Transplantation (260.2). 1991; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?MCDId=17&ExpandComments=n&McdName=Clinical+Pharmacology+Compendium+Revision+Request+-+CAG-00392&mcdtypename=Compendia&MCDIndexType=6&NCDId=71&ncdver=1&bc=AgAEA AAAAgAA&>.

### **Policy History**

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

05/01/2014	Medical Policy Committee review
05/21/2014	Medical Policy Implementation Committee approval. New policy.
03/05/2015	Medical Policy Committee review
03/20/2015	Medical Policy Implementation Committee approval. Removed requirement for “Patients with ongoing alcohol and/or drug abuse. (Evidence for abstinence may vary among liver transplant programs, but generally a minimum of 3 months is required.)”.
08/03/2015	Coding update: ICD10 Diagnosis code section added; ICD9 Procedure code section removed.
01/01/2016	Coding update
03/03/2016	Medical Policy Committee review
03/16/2016	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
01/01/2017	Coding update: Removing ICD-9 Diagnosis Codes
03/02/2017	Medical Policy Committee review
03/15/2017	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
03/01/2018	Medical Policy Committee review
03/21/2018	Medical Policy Implementation Committee approval. Policy title changed from “Liver Transplant” to “Liver Transplant and Combined Liver-Kidney Transplant.” Combined liver-kidney transplantation added to policy as eligible for coverage. Policy Guidelines moved from the Background/Overview to a new Policy

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

- 01/01/2019 Guidelines section. Contraindication for smoking and HIV criteria added to Policy Guidelines.
- 01/01/2019 Coding update
- 03/07/2019 Medical Policy Committee review
- 03/20/2019 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
- 12/10/2019 Coding update
- 03/05/2020 Medical Policy Committee review
- 03/11/2020 Medical Policy Implementation Committee approval. Added a reference to see the patient selection criteria in the Policy Guidelines section for liver transplantation in patients with unresectable hilar cholangiocarcinoma. Added a not medically necessary criteria bullet as follows:
  - Patients with ongoing alcohol and/or drug abuse. (Evidence for abstinence may vary among liver transplant programs, but generally a minimum of 3 months is required.)
- 09/16/2020 Coding update
- 03/04/2021 Medical Policy Committee review
- 03/10/2021 Medical Policy Implementation Committee approval. Coverage eligibility unchanged. Added information on tobacco consumption under Liver-Specific Criteria in the Policy Guidelines.

Next Scheduled Review Date: 03/2022

### Coding

*The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)‡, copyright 2020 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.*

*The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Blue Cross*

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

*Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.*

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	47135, 47140, 47141, 47142, 47143, 47144, 47145, 47146, 47147, 47399
HCPCS	No codes
ICD-10 Diagnosis	A52.15, B15.0B15.9, B16.0-B16.9, B17.10-B17.11, B17.8-B17.9, B18.0-B18.9, B19.10-B19.11, B19.9, B25.1, B66.1, B66.3, D64.0-D64.3, D81.810, D84.1, E70.0-E70.1, E70.21-E70.29, E70.30, E70.310-E70.319, E70.320-E70.329, E70.330-E70.339, E70.40-E70.41, E70.49, E70.5, E70.8-E70.9, E71.0, E71.110-E71.111, E71.118, E71.120-E71.121, E71.128, E71.19, E71.2, E71.30, E72.00-E72.09, E72.10-E72.19, E72.20-E72.29, E72.3-E72.4, E72.50-E72.59, E74.8-E74.9, E75.21-E75.22, E75.240-E75.249, E75.3, E75.5-E75.6, E77.0-E77.9, E78.0-E78.6, E78.70-E78.79, E78.81-E78.89, E78.9, E80.0-E80.1, E80.20-E80.29, E83.00-E83.19, E85.0-E85.9 E88.1-E88.2, E88.89, E88.9, G63, G65.0-G65.2, G70.00-G70.01, I74.8, I82.0, I87.9, I99.9, K71.0-K71.9, K74.3-K74.5, K75.2-K75.3, K75.81, K75.89, K75.9, K76.0, K76.4, K76.89, K77, K80.30-K80.37, K83.01-K83.1, K83.5, K83.8, K87, M34.83, Q43.4-Q43.9, S31.609A, S36.112A-S36.119A, T86.40-T86.49, Z52.6

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.





# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

Added codes eff 10/1/2020: D84.81, D84.821, D84.822, D84.89, E70.81, E70.89, E74.810, E74.818, E74.819, E74.89, K74.00-K74.02, T86.8401-T86.8409, T86.8411-T86.8419, T86.8421-T86.8429, T86.8481-T86.8489, T86.8491-T86.8499
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

**\*Investigational** – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
  - 1. Consultation with the Blue Cross and Blue Shield Association technology assessment program (TEC) or other nonaffiliated technology evaluation center(s);
  - 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
  - 3. Reference to federal regulations.

**\*\*Medically Necessary** (or “Medical Necessity”) - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. In accordance with nationally accepted standards of medical practice;
- B. Clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and
- C. Not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Liver Transplant and Combined Liver-Kidney Transplant

Policy # 00411

Original Effective Date: 05/21/2014

Current Effective Date: 04/12/2021

at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.

For these purposes, "nationally accepted standards of medical practice" means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

‡ Indicated trademarks are the registered trademarks of their respective owners.

**NOTICE:** If the Patient's health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

**NOTICE:** Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.