



# Louisiana

## **Sacral Nerve Neuromodulation/Stimulation**

**Policy #** 00108

Original Effective Date: 03/25/2002

Current Effective Date: 07/11/2022

*Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.*

*Note: Transanal Radiofrequency Treatment of Fecal Incontinence is addressed separately in medical policy 00571.*

*Note: Percutaneous Tibial Nerve Stimulation is addressed separately in medical policy 00415.*

## **Urinary Incontinence and Non-obstructive Retention**

### **When Services May Be Eligible for Coverage**

*Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:*

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Based on review of available data, the Company may consider a trial period of sacral nerve neuromodulation (SNM) with either percutaneous nerve stimulation or a temporarily implanted lead to be **eligible for coverage\*\***.

#### **Patient Selection Criteria**

Coverage eligibility may be considered for a trial period of SNM with either percutaneous nerve stimulation or a temporarily implanted lead in patients who meet ALL of the following criteria:

- There is a diagnosis of at least ONE of the following:
  - Urge incontinence; OR
  - Urgency-frequency syndrome; OR
  - Non-obstructive urinary retention; OR
  - Overactive bladder;

AND

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Sacral Nerve Neuromodulation/Stimulation

Policy # 00108

Original Effective Date: 03/25/2002

Current Effective Date: 07/11/2022

- There is documented failure or intolerance to at least two conventional conservative therapies (e.g., behavioral training such as bladder training, prompted voiding, or pelvic muscle exercise training, pharmacologic treatment for at least a sufficient duration to fully assess its efficacy, and/or surgical corrective therapy); AND
- The patient is an appropriate surgical candidate; AND
- Incontinence or retention is not related to a neurologic condition (e.g., Parkinson's disease, multiple sclerosis, spinal cord injury).

Based on review of available data, the Company may consider permanent implantation of a sacral nerve neuromodulation (SNM) device to be **eligible for coverage\*\***.

### Patient Selection Criteria

Coverage eligibility may be considered for permanent implantation of a SNM device in patients who meet ALL of the following criteria:

- All of the above criteria in the Urinary Incontinence and Non-obstructive Retention section are met, AND
- A trial stimulation period demonstrates at least 50% improvement in symptoms over a period of at least 48 hours.

## **When Services Are Considered Investigational**

*Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.*

Based on review of available data, the Company considers other urinary/voiding applications of sacral nerve neuromodulation (SNM), including but not limited to treatment of stress incontinence or urge incontinence due to a neurologic condition (eg, detrusor hyperreflexia, multiple sclerosis, spinal cord injury, other types of chronic voiding dysfunction) to be **investigational.\***

The use of sacral nerve neuromodulation (SNM) is considered to be **investigational\*** when patient selection criteria are not met for the Urinary Incontinence and Non-obstructive Retention coverage section.

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

Sacral Nerve Neuromodulation/Stimulation

Policy # 00108

Original Effective Date: 03/25/2002

Current Effective Date: 07/11/2022

## **Fecal Incontinence**

### **When Services May Be Eligible for Coverage**

*Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:*

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Based on review of available data, the Company may consider a trial period of sacral nerve neuromodulation (SNM) with either percutaneous nerve stimulation or a temporarily implanted lead to be **eligible for coverage\*\***.

#### **Patient Selection Criteria**

Coverage eligibility may be considered for a trial period of SNM with either percutaneous nerve stimulation or a temporarily implanted lead in patients who meet ALL of the following criteria:

- There is a diagnosis of chronic fecal incontinence of more than 2 incontinent episodes on average per week for more than 6 months or for more than 12 months after vaginal childbirth; AND
- There is documented failure or intolerance to conventional conservative therapy (eg, dietary modification, the addition of bulking and pharmacologic treatment) for at least a sufficient duration to fully assess its efficacy; AND
- The patient is an appropriate surgical candidate; AND
- The condition is not related to an anorectal malformation (eg, congenital anorectal malformation; defects of the external anal sphincter over 60°; visible sequelae of pelvic radiation; active anal abscesses and fistulae) or chronic inflammatory bowel disease; AND
- Incontinence is not related to a neurologic condition; AND
- The patient has not had rectal surgery in the previous 12 months or, in the case of cancer, the patient has not had rectal surgery in the past 24 months.

Based on review of available data, the Company may consider permanent implantation of a sacral nerve neuromodulation (SNM) device to be **eligible for coverage\*\***.

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Sacral Nerve Neuromodulation/Stimulation

Policy # 00108

Original Effective Date: 03/25/2002

Current Effective Date: 07/11/2022

### Patient Selection Criteria

Coverage eligibility may be considered for permanent implantation of a SNM device in patients who meet ALL of the following criteria:

- All of the above criteria in the Fecal Incontinence section are met; AND
- A trial stimulation period demonstrates at least 50% improvement in symptoms over a period of at least 48 hours.

## **When Services Are Considered Investigational**

*Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.*

Based on review of available data, the Company considers sacral nerve neuromodulation (SNM) in the treatment of chronic constipation or chronic pelvic pain to be **investigational**.\*

The use of sacral nerve neuromodulation (SNM) is considered to be **investigational**\* when patient selection criteria are not met for the Fecal Incontinence coverage section.

## **Policy Guidelines**

The International Continence Society has defined overactive bladder syndrome as “urinary urgency, usually accompanied by increased daytime frequency and/or nocturia, with urinary incontinence (OAB-wet) or without (OAB-dry), in the absence of urinary tract infection or other detectable disease”

(available at <https://www.ics.org/glossary/symptom/overactivebladderoaburgencysyndrome>).

## **Background/Overview**

### **Treatment**

Treatment using sacral nerve neuromodulation, also known as indirect sacral nerve stimulation, is 1 of several alternative modalities for patients with urinary or fecal incontinence (urge incontinence, significant symptoms of urgency-frequency, nonobstructive urinary retention) who have failed behavioral (eg, prompted voiding) and/or pharmacologic therapies.

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Sacral Nerve Neuromodulation/Stimulation

Policy # 00108

Original Effective Date: 03/25/2002

Current Effective Date: 07/11/2022

The sacral nerve neuromodulation device consists of an implantable pulse generator that delivers controlled electrical impulses. This pulse generator is attached to wire leads that connect to the sacral nerves, most commonly the S3 nerve root. Two external components of the system help control the electrical stimulation. A control magnet, kept by the patient, is used to turn the device on or off. A console programmer is kept by the physician and used to adjust the settings of the pulse generator.

Before implantation of the permanent device, patients undergo an initial testing phase to estimate potential response to treatment. The first type of testing developed was percutaneous nerve evaluation. This procedure is done with the patient under local anesthesia, using a test needle to identify the appropriate sacral nerve(s). Once identified, a temporary wire lead is inserted through the test needle and left in place for 4 to 7 days. This lead is connected to an external stimulator, which is carried by patients in their pocket or on their belt. The results of this test phase are used to determine whether patients are appropriate candidates for the permanent device. If patients show a 50% or greater reduction in symptom frequency, they are deemed eligible for the permanent device.

The second type of testing is a 2 stage surgical procedure. In the first stage, a quadripolar-tined lead is implanted (stage 1). The testing phase can last as long as several weeks, and if patients show a 50% or greater reduction in symptom frequency, they can proceed to stage 2 of the surgery, which is permanent implantation of the neuromodulation device. The 2 stage surgical procedure has been used in various ways. They include its use instead of percutaneous nerve evaluation, for patients who failed percutaneous nerve evaluation, for patients with an inconclusive percutaneous nerve evaluation, or for patients who had a successful percutaneous nerve evaluation to refine patient selection further.

The permanent device is implanted with the patient under general anesthesia. The electrical leads are placed in contact with the sacral nerve root(s) via an incision in the lower back, and the wire leads are extended through a second incision underneath the skin, across the flank to the lower abdomen. Finally, a third incision is made in the lower abdomen where the pulse generator is inserted and connected to the wire leads. Following implantation, the physician programs the pulse generator to the optimal settings for that patient. The patient can switch the pulse generator on and off by placing the control magnet over the area of the pulse generator for 1 to 2 seconds.

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

Sacral Nerve Neuromodulation/Stimulation

Policy # 00108

Original Effective Date: 03/25/2002

Current Effective Date: 07/11/2022

This evidence review does not address pelvic floor stimulation, which refers to electrical stimulation of the pudendal nerve. Also, this review does not address devices that provide direct sacral nerve stimulation in patients with spinal cord injuries.

## **FDA or Other Governmental Regulatory Approval**

### **U.S. Food and Drug Administration (FDA)**

In 1997, the InterStim<sup>®</sup> Sacral Nerve Stimulation system (Medtronic) was approved by the U.S. FDA through the premarket approval process for the indication of urinary urge incontinence in patients who have failed or could not tolerate more conservative treatments. In 1999, the device received the FDA approval for the additional indications of urgency-frequency and urinary retention in patients without mechanical obstruction. In 2006, the InterStim II<sup>®</sup> System (Medtronic) was approved by the FDA through the premarket approval process for the treatment of intractable cases of overactive bladder and urinary retention. The new device is smaller and lighter than the original and is reported to be suited for those with lower energy requirements or small stature. The device also includes updated software and programming options.

In 2011, the InterStim System was approved by the FDA through the premarket approval process for the indication of chronic fecal incontinence in patients who have failed or could not tolerate more conservative treatments.

The InterStim device has not been specifically approved by the FDA for the treatment of chronic pelvic pain.

FDA product code: EZW.

## **Rationale/Source**

This medical policy was developed through consideration of peer-reviewed medical literature generally recognized by the relevant medical community, U.S. Food and Drug Administration approval status, nationally accepted standards of medical practice and accepted standards of medical practice in this community, technology evaluation centers, reference to federal regulations, other plan medical policies, and accredited national guidelines.

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Sacral Nerve Neuromodulation/Stimulation

Policy # 00108

Original Effective Date: 03/25/2002

Current Effective Date: 07/11/2022

Sacral nerve neuromodulation, also known as sacral nerve stimulation, involves the implantation of a permanent device that modulates the neural pathways controlling bladder or rectal function. This evidence review addresses the use of sacral nerve neuromodulation to treat urinary or fecal incontinence, fecal nonobstructive retention, and chronic pelvic pain in patients with intact neural innervation of the bladder and/or rectum.

### **Summary of Evidence**

For individuals with urinary incontinence who have failed conservative treatment who receive sacral nerve neuromodulation, the evidence includes randomized controlled trials (RCTs) and case series. Relevant outcomes are symptoms, morbid events, and treatment, related morbidity. Results from the RCTs and case series with long-term follow-up have suggested that sacral nerve neuromodulation reduces symptoms of urge incontinence, urgency-frequency syndrome, nonobstructive urinary retention, and overactive bladder in selected patients. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with fecal incontinence who have failed conservative treatment who receive sacral nerve neuromodulation, the evidence includes RCTs, systematic reviews, and observational studies including several with long-term follow-up. Relevant outcomes are symptoms, morbid events, and treatment, related morbidity. Although relatively small, the available trials had a low risk of bias and demonstrated improvements in incontinence relative to alternatives in selected patients. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with constipation who have failed conservative treatment who receive sacral nerve neuromodulation, the evidence includes RCTs, systematic reviews, and case series including several with long-term follow-up. Relevant outcomes are symptoms, morbid events, and treatment, related morbidity. The available trials have not consistently reported improvements in outcomes with sacral nerve neuromodulation. Additional studies are needed to demonstrate the health benefits of this technology. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with chronic pelvic pain who receive sacral nerve neuromodulation, the evidence is limited to case series. Relevant outcomes are symptoms, morbid events, and treatment-related

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

Sacral Nerve Neuromodulation/Stimulation

Policy # 00108

Original Effective Date: 03/25/2002

Current Effective Date: 07/11/2022

morbidity. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

## **Supplemental Information**

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

### **Clinical Input From Physician Specialty Societies and Academic Medical Centers**

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

In response to requests, input was received from 4 physician specialty societies and 2 academic medical centers while this policy was under review in 2012. Reviewers from 2 specialty societies and 2 academic medical centers provided opinions on the possible medical necessity of implantable leads for test stimulation, as part of a 2-stage process for device implantation. All 4 respondents supported the use of implantable leads for test stimulation as an alternative to percutaneous test stimulation, for patients who had failed percutaneous test stimulation and/or for patients with inconclusive percutaneous test stimulation. Reasons for support included a longer period of interrupted treatment with stage-1 stimulation due to less lead migration and a higher rate of positive tests compared with percutaneous test stimulation.

### **Practice Guidelines and Position Statements**

Guidelines or position statements will be considered for inclusion in ‘Supplemental Information’ if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

## **Urinary Disorders**

### **American Urological Association**

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.





# Louisiana

## Sacral Nerve Neuromodulation/Stimulation

Policy # 00108

Original Effective Date: 03/25/2002

Current Effective Date: 07/11/2022

In 2019, the American Urological Association updated its guidelines on the diagnosis and treatment of overactive bladder. The guidelines stated that sacral neuromodulation may be offered as a third-line treatment in carefully selected patients with severe refractory symptoms or into those who are not candidates for second-line therapy (eg, oral antimuscarinics, oral  $\beta$ 3-adrenoceptor agonists, transdermal oxybutynin) and are willing to undergo surgery (recommendation, evidence strength Grade C).

### **American College of Obstetricians and Gynecologists**

A 2015 practice bulletin on urinary incontinence (replaced practice bulletin number 63, 2005; reaffirmed in 2019) from the College stated, "sacral neuromodulation may be considered for patients with recalcitrant urinary urge incontinence who have failed other conservative measures, including bladder training, pelvic floor physical therapy with biofeedback, and pharmacologic treatment."

### **Fecal Disorders**

#### **National Institute for Health and Care Excellence**

In 2007, the National Institute for Health and Care Excellence issued guidance on the management of fecal incontinence. The guidance was reviewed in 2014 and 2018, and no changes were made. The guidance has recommended:

"a trial of temporary sacral nerve stimulation should be considered for people with fecal incontinence in whom sphincter surgery is deemed inappropriate.... All individuals should be informed of the potential benefits and limitations of this procedure and should undergo a trial stimulation period of at least 2 weeks to determine if they are likely to benefit. People with fecal incontinence should be offered sacral nerve stimulation on the basis of their response to percutaneous nerve evaluation during specialist assessment, which is predictive of therapy success."

#### **American College of Gastroenterology**

In its 2014 clinical guideline on the management of benign anorectal disorders, including fecal incontinence, the American College of Gastroenterology (ACG) found that "sacral nerve stimulation should be considered in [fecal incontinence] who do not respond to conservative therapy (strong recommendation, moderate quality of evidence)."<sup>42</sup> The 2021 update of these guidelines keep the recommendation for sacral nerve stimulation in patients with fecal incontinence refractory to medical therapy the same as in the 2014 version.<sup>43</sup> Additionally, due to a lack of evidence supporting efficacy and the risk of adverse events and complications, the 2021 ACG Panel makes a statement

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Sacral Nerve Neuromodulation/Stimulation

Policy # 00108

Original Effective Date: 03/25/2002

Current Effective Date: 07/11/2022

stating that sacral nerve stimulation "cannot be recommended in patients with constipation of any type."

### **American College of Obstetricians and Gynecologists**

A 2020 practice bulletin on fecal incontinence from the College stated, "Sacral nerve stimulation can be considered as a surgical treatment option for women with fecal incontinence with or without anal sphincter disruption who have failed conservative treatments."

### **American Society of Colon and Rectal Surgeons**

In 2015, the American Society of Colon and Rectal Surgeons released a clinical practice guideline for the treatment of fecal incontinence. They stated that "sacral neuromodulation may be considered as a first-line surgical option for incontinent patients with and without sphincter defects (Grade of Recommendation: Strong, based on moderate-quality evidence, 1B)."

In 2016, the Society released a clinical practice guideline for the management of constipation. In this guideline, they stated "sacral neuromodulation may be an effective treatment for patients with chronic constipation and successful peripheral nerve evaluation test when conservative measures have failed; however, it is not currently approved by the US Food and Drug Administration for this condition in the United States (Grade of Recommendation: Weak, based on moderate quality evidence, 2B)."

### **U.S. Preventive Services Task Force Recommendations**

Not applicable.

### **Medicare National Coverage**

In 2002, the Centers for Medicare & Medicaid Services covers sacral nerve stimulation for the "treatment of urinary urge incontinence, urgency-frequency syndrome, and urinary retention." Sacral nerve stimulation "involves both a temporary test stimulation to determine if an implantable stimulator would be effective and a permanent implantation in appropriate candidates. Both the test and the permanent implantation are covered."

"The following limitations for coverage apply to all three indications:

- Patients must be refractory to conventional therapy ... and be appropriate surgical candidates such that implantation with anesthesia can occur.

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Sacral Nerve Neuromodulation/Stimulation

Policy # 00108

Original Effective Date: 03/25/2002

Current Effective Date: 07/11/2022

- Patients with stress incontinence, urinary obstruction, and specific neurologic diseases ... that are associated with secondary manifestations ... are excluded.
- Patients must have had successful test stimulation in order to support subsequent implantation. Before patients are eligible for permanent implantation, they must demonstrate a 50% or greater improvement through test stimulation. Improvement is measured through voiding diaries.”

### Ongoing and Unpublished Clinical Trials

Some currently ongoing trials that might influence this review are listed in Table 1.

**Table 1. Summary of Key Trials**

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT03139734	Sacral Neuromodulation for Pelvic Pain Associated with Endometriosis	50	May 2022
NCT03811821	Comparative Effects of Biofeedback, Sacral Nerve Stimulation, and Injectable Bulking Agents for Treatment of Fecal Incontinence: The Fecal Incontinence Treatment Study (FIT) Study	285	Dec 2025
NCT04710433	Non-invasive Sacral Nerve Stimulation in Children and Adolescents With Chronic Constipation: a case-control study on external neuromodulatory treatment	50	Dec 2021
NCT04506866	Evaluation of InterStim Micro System Performance and Safety (ELITE) to Confirm Long-Term Outcomes - Post Market Clinical Follow-Up Study	160	Apr 2024
NCT04713085	Sacral Nerve Stimulation in Children and Adolescents With Chronic Constipation: a Case-control Study on Invasive and Non-invasive Neuromodulatory Treatment	30	June 2022

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Sacral Nerve Neuromodulation/Stimulation

Policy # 00108

Original Effective Date: 03/25/2002

Current Effective Date: 07/11/2022

NCT04549818	Does Sacral Neuromodulation Relieve Chronic Pelvic Cancer Pain, Compared With Medical Treatment?	44	Dec 2021
NCT04232696 <sup>a</sup>	Clinical Study of Neuraspera's Implantable Sacral Nerve Stimulation (SNS) System in Patients With Symptoms of Overactive Bladder (OAB)	310	Dec 2023
NCT02577302 <sup>a</sup>	Multi-center, Prospective, Randomized, Controlled, Non-Inferiority, Clinical Trial of Chronic Afferent Nerve Stimulation (CAN-Stim) of the Tibial Nerve Versus Sacral Nerve Stimulation (SNS) in the Treatment of Urinary Urgency Incontinence Resulting From Refractory Overactive Bladder (OAB)	200	Oct 2021

<sup>a</sup> denotes an industry-sponsored trial  
NCT: national clinical trial.

## References

1. Food and Drug Administration (FDA). Summary of Safety and Effectiveness: Medtronic Interstim System for Urinary Control. [http://www.accessdata.fda.gov/cdrh\\_docs/pdf/P970004S004b.pdf](http://www.accessdata.fda.gov/cdrh_docs/pdf/P970004S004b.pdf).
2. Weil EH, Ruiz-Cerda JL, Eerdmans PH, et al. Sacral root neuromodulation in the treatment of refractory urinary urge incontinence: a prospective randomized clinical trial. *Eur Urol.* Feb 2000; 37(2): 161-71. PMID 10705194
3. Siegel S, Noblett K, Mangel J, et al. Results of a prospective, randomized, multicenter study evaluating sacral neuromodulation with InterStim therapy compared to standard medical therapy at 6-months in subjects with mild symptoms of overactive bladder. *Neurourol Urodyn.* Mar 2015; 34(3): 224-30. PMID 24415559
4. Noblett K, Siegel S, Mangel J, et al. Results of a prospective, multicenter study evaluating quality of life, safety, and efficacy of sacral neuromodulation at twelve months in subjects with symptoms of overactive bladder. *Neurourol Urodyn.* Feb 2016; 35(2): 246-51. PMID 25546568
5. Amundsen CL, Richter HE, Menefee SA, et al. OnabotulinumtoxinA vs Sacral Neuromodulation on Refractory Urgency Urinary Incontinence in Women: A Randomized Clinical Trial. *JAMA.* Oct 04 2016; 316(13): 1366-1374. PMID 27701661

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Sacral Nerve Neuromodulation/Stimulation

Policy # 00108

Original Effective Date: 03/25/2002

Current Effective Date: 07/11/2022

6. Groen J, Blok BF, Bosch JL. Sacral neuromodulation as treatment for refractory idiopathic urge urinary incontinence: 5-year results of a longitudinal study in 60 women. *J Urol.* Sep 2011; 186(3): 954-9. PMID 21791355
7. White WM, Mobley JD, Doggweiler R, et al. Incidence and predictors of complications with sacral neuromodulation. *Urology.* Apr 2009; 73(4): 731-5. PMID 19193415
8. Thaha MA, Abukar AA, Thin NN, et al. Sacral nerve stimulation for faecal incontinence and constipation in adults. *Cochrane Database Syst Rev.* Aug 24 2015; (8): CD004464. PMID 26299888
9. Thin NN, Horrocks EJ, Hotouras A, et al. Systematic review of the clinical effectiveness of neuromodulation in the treatment of faecal incontinence. *Br J Surg.* Oct 2013; 100(11): 1430-47. PMID 24037562
10. Tan E, Ngo NT, Darzi A, et al. Meta-analysis: sacral nerve stimulation versus conservative therapy in the treatment of faecal incontinence. *Int J Colorectal Dis.* Mar 2011; 26(3): 275-94. PMID 21279370
11. Maeda Y, Matzel K, Lundby L, et al. Postoperative issues of sacral nerve stimulation for fecal incontinence and constipation: a systematic literature review and treatment guideline. *Dis Colon Rectum.* Nov 2011; 54(11): 1443-60. PMID 21979192
12. Tjandra JJ, Chan MK, Yeh CH, et al. Sacral nerve stimulation is more effective than optimal medical therapy for severe fecal incontinence: a randomized, controlled study. *Dis Colon Rectum.* May 2008; 51(5): 494-502. PMID 18278532
13. Leroi AM, Parc Y, Lehur PA, et al. Efficacy of sacral nerve stimulation for fecal incontinence: results of a multicenter double-blind crossover study. *Ann Surg.* Nov 2005; 242(5): 662-9. PMID 16244539
14. Wexner SD, Collier JA, Devroede G, et al. Sacral nerve stimulation for fecal incontinence: results of a 120-patient prospective multicenter study. *Ann Surg.* Mar 2010; 251(3): 441-9. PMID 20160636
15. Mellgren A, Wexner SD, Collier JA, et al. Long-term efficacy and safety of sacral nerve stimulation for fecal incontinence. *Dis Colon Rectum.* Sep 2011; 54(9): 1065-75. PMID 21825885
16. Hull T, Giese C, Wexner SD, et al. Long-term durability of sacral nerve stimulation therapy for chronic fecal incontinence. *Dis Colon Rectum.* Feb 2013; 56(2): 234-45. PMID 23303153
17. Altomare DF, Giuratrabocchetta S, Knowles CH, et al. Long-term outcomes of sacral nerve stimulation for faecal incontinence. *Br J Surg.* Mar 2015; 102(4): 407-15. PMID 25644687

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Sacral Nerve Neuromodulation/Stimulation

Policy # 00108

Original Effective Date: 03/25/2002

Current Effective Date: 07/11/2022

18. Leo CA, Thomas GP, Bradshaw E, et al. Long-term outcome of sacral nerve stimulation for faecal incontinence. *Colorectal Dis.* Dec 2020; 22(12): 2191-2198. PMID 32954658
19. Desprez C, Damon H, Meurette G, et al. Ten-year Evaluation of a Large Retrospective Cohort Treated by Sacral Nerve Modulation for Fecal Incontinence: Results of a French Multicenter Study. *Ann Surg.* Apr 01 2022; 275(4): 735-742. PMID 32740249
20. De Meyere C, Nuytens F, Parmentier I, et al. Five-year single center experience of sacral neuromodulation for isolated fecal incontinence or fecal incontinence combined with low anterior resection syndrome. *Tech Coloproctol.* Sep 2020; 24(9): 947-958. PMID 32556866
21. Picciariello A, Rinaldi M, Dibra R, et al. Ageing with sacral nerve modulation for fecal incontinence: how many patients get benefit after more than 10 years?. *Updates Surg.* Feb 2022; 74(1): 185-191. PMID 34982410
22. Pauwels N, Willemsse C, Hellemans S, et al. The role of neuromodulation in chronic functional constipation: a systematic review. *Acta Gastroenterol Belg.* Jul-Sep 2021; 84(3): 467-476. PMID 34599572
23. Pilkington SA, Emmett C, Knowles CH, et al. Surgery for constipation: systematic review and practice recommendations: Results V: Sacral Nerve Stimulation. *Colorectal Dis.* Sep 2017; 19 Suppl 3: 92-100. PMID 28960926
24. Thomas GP, Dudding TC, Rahbour G, et al. Sacral nerve stimulation for constipation. *Br J Surg.* Jan 2013; 100(2): 174-81. PMID 23124687
25. Knowles CH, Thin N, Gill K, et al. Prospective randomized double-blind study of temporary sacral nerve stimulation in patients with rectal evacuatory dysfunction and rectal hyposensitivity. *Ann Surg.* Apr 2012; 255(4): 643-9. PMID 22418005
26. Zerbib F, Siproudhis L, Lehur PA, et al. Randomized clinical trial of sacral nerve stimulation for refractory constipation. *Br J Surg.* Feb 2017; 104(3): 205-213. PMID 27779312
27. Dinning PG, Hunt L, Patton V, et al. Treatment efficacy of sacral nerve stimulation in slow transit constipation: a two-phase, double-blind randomized controlled crossover study. *Am J Gastroenterol.* May 2015; 110(5): 733-40. PMID 25895520
28. Kamm MA, Dudding TC, Melenhorst J, et al. Sacral nerve stimulation for intractable constipation. *Gut.* Mar 2010; 59(3): 333-40. PMID 20207638
29. Maeda Y, Lundby L, Buntzen S, et al. Sacral nerve stimulation for constipation: suboptimal outcome and adverse events. *Dis Colon Rectum.* Jul 2010; 53(7): 995-9. PMID 20551750
30. Tirlapur SA, Vlismas A, Ball E, et al. Nerve stimulation for chronic pelvic pain and bladder pain syndrome: a systematic review. *Acta Obstet Gynecol Scand.* Aug 2013; 92(8): 881-7. PMID 23710833

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Sacral Nerve Neuromodulation/Stimulation

Policy # 00108

Original Effective Date: 03/25/2002

Current Effective Date: 07/11/2022

31. Martellucci J, Naldini G, Carriero A. Sacral nerve modulation in the treatment of chronic pelvic pain. *Int J Colorectal Dis.* Jul 2012; 27(7): 921-6. PMID 22203519
32. Siegel S, Paszkiewicz E, Kirkpatrick C, et al. Sacral nerve stimulation in patients with chronic intractable pelvic pain. *J Urol.* Nov 2001; 166(5): 1742-5. PMID 11586214
33. Baxter C, Kim JH. Contrasting the percutaneous nerve evaluation versus staged implantation in sacral neuromodulation. *Curr Urol Rep.* Sep 2010; 11(5): 310-4. PMID 20535593
34. Leong RK, De Wachter SG, Nieman FH, et al. PNE versus 1st stage tined lead procedure: a direct comparison to select the most sensitive test method to identify patients suitable for sacral neuromodulation therapy. *Neurourol Urodyn.* Sep 2011; 30(7): 1249-52. PMID 21404317
35. Scheepens WA, Van Koeveringe GA, De Bie RA, et al. Long-term efficacy and safety results of the two-stage implantation technique in sacral neuromodulation. *BJU Int.* Dec 2002; 90(9): 840-5. PMID 12460343
36. Marcelissen TA, Leong RK, de Bie RA, et al. Long-term results of sacral neuromodulation with the tined lead procedure. *J Urol.* Nov 2010; 184(5): 1997-2000. PMID 20850820
37. Borawski KM, Foster RT, Webster GD, et al. Predicting implantation with a neuromodulator using two different test stimulation techniques: A prospective randomized study in urge incontinent women. *Neurourol Urodyn.* 2007; 26(1): 14-8. PMID 17123297
38. Bannowsky A, Wefer B, Braun PM, et al. Urodynamic changes and response rates in patients treated with permanent electrodes compared to conventional wire electrodes in the peripheral nerve evaluation test. *World J Urol.* Dec 2008; 26(6): 623-6. PMID 18629503
39. Diagnosis and treatment of non-neurogenic overactive bladder (OAB) in adults: an AUA/SUFU Guideline (2019). [https://www.auanet.org/guidelines/overactive-bladder-\(oab\)-guideline](https://www.auanet.org/guidelines/overactive-bladder-(oab)-guideline).
40. ACOG Practice Bulletin No. 155: Urinary Incontinence in Women. *Obstet Gynecol.* Nov 2015; 126(5): e66-e81. PMID 26488524
41. National Institute for Health and Care Excellence (NICE). Faecal incontinence in adults: management [CG49]. 2007; <https://www.nice.org.uk/guidance/CG49>. Accessed March 5, 2022.
42. Wald A, Bharucha AE, Cosman BC, et al. ACG clinical guideline: management of benign anorectal disorders. *Am J Gastroenterol.* Aug 2014; 109(8): 1141-57; (Quiz) 1058. PMID 25022811
43. Wald A, Bharucha AE, Limketkai B, et al. ACG Clinical Guidelines: Management of Benign Anorectal Disorders. *Am J Gastroenterol.* Oct 01 2021; 116(10): 1987-2008. PMID 34618700
44. ACOG Practice Bulletin No. 210: Fecal Incontinence. *Obstet Gynecol.* Apr 2019; 133(4): e260-e273. PMID 30913197

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Sacral Nerve Neuromodulation/Stimulation

Policy # 00108

Original Effective Date: 03/25/2002

Current Effective Date: 07/11/2022

45. Paquette IM, Varma MG, Kaiser AM, et al. The American Society of Colon and Rectal Surgeons' Clinical Practice Guideline for the Treatment of Fecal Incontinence. *Dis Colon Rectum*. Jul 2015; 58(7): 623-36. PMID 26200676
46. Paquette IM, Varma M, Ternent C, et al. The American Society of Colon and Rectal Surgeons' Clinical Practice Guideline for the Evaluation and Management of Constipation. *Dis Colon Rectum*. Jun 2016; 59(6): 479-92. PMID 27145304
47. Centers for Medicare & Medicaid Services. National Coverage Determination (NCD) for SACRAL NERVE Stimulation For Urinary Incontinence (230.18). 2002; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=249&ncdver=1&CoverageSelection=National&KeyWord=sacral+nerve&KeyWordLookUp=T> title&KeyWordSearchType=And&clickon=search&bc=gAAAABAAAAAAAAAA%3d%3d&

### **Policy History**

Original Effective Date: 03/25/2002

Current Effective Date: 07/11/2022

- |            |   |
|------------|---|
| 03/21/2002 | Medical Policy Committee review   |
| 03/25/2002 | Managed Care Advisory Council approval  |
| 06/24/2002 | Format revision. No substance change to policy.   |
| 10/05/2004 | Medical Director review   |
| 11/16/2004 | Medical Policy Committee review. Format revision. Policy focus expanded to include other pelvic floor dysfunction conditions in addition to urinary incontinence. |
| 11/29/2004 | Managed Care Advisory Council approval  |
| 10/05/2005 | Medical Director review   |
| 10/18/2005 | Medical Policy Committee review. Format revision. FDA approval information added. Coverage eligibility unchanged.   |
| 10/27/2005 | Quality Care Advisory Council approval  |
| 10/04/2006 | Medical Director review   |
| 10/18/2006 | Medical Policy Committee approval. Format revision; updated with additional references. Coverage eligibility unchanged.   |
| 09/05/2007 | Medical Director review   |
| 09/19/2007 | Medical Policy Committee approval. Coverage eligibility unchanged.  |
| 09/09/2008 | Medical Director review   |

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.





# Louisiana

## Sacral Nerve Neuromodulation/Stimulation

Policy # 00108

Original Effective Date: 03/25/2002

Current Effective Date: 07/11/2022

- 09/17/2008 Medical Policy Committee approval. Coverage eligibility unchanged. A note stating that a successful trial response to a peripheral nerve stimulation test is required prior to permanent placement under general anesthesia.
- 06/04/2009 Medical Director review
- 06/17/2009 Medical Policy Committee approval. Coverage eligibility unchanged.
- 06/03/2010 Medical Policy Committee approval
- 06/16/2010 Medical Policy Implementation Committee approval. Added criteria for SNM for the treatment of patients with urge incontinence, urgency-frequency and non-obstructive urinary retention to be eligible for coverage as follows:
- The patient is an appropriate surgical candidate; and
  - A successful percutaneous test stimulation, defined as at least 50% improvement in symptoms, was performed.
- Change coverage for fecal incontinence from investigational to eligible for coverage with criteria.
- 06/02/2011 Medical Policy Committee review
- 06/15/2011 Medical Policy Implementation Committee approval. Format revision; including addition of FDA and or other governmental regulatory approval and rationale/source. Coverage eligibility unchanged.
- 06/14/2012 Medical Policy Committee review
- 06/20/2012 Medical Policy Implementation Committee approval. No change to coverage.
- 06/06/2013 Medical Policy Committee review
- 06/25/2013 Medical Policy Implementation Committee approval. Title changed. “For Pelvic Floor” dropped. Criteria revised.
- 06/05/2014 Medical Policy Committee review
- 06/18/2014 Medical Policy Implementation Committee approval. No change to coverage.
- 06/04/2015 Medical Policy Committee review
- 06/17/2015 Medical Policy Implementation Committee approval. Added overactive bladder to criteria section of eligibility statement.
- 08/03/2015 Coding update: ICD10 Diagnosis code section added; ICD9 Procedure code section removed.
- 06/02/2016 Medical Policy Committee review

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Sacral Nerve Neuromodulation/Stimulation

Policy # 00108

Original Effective Date: 03/25/2002

Current Effective Date: 07/11/2022

12 months, or in the case of cancer, the patient has not had rectal surgery in the past 24

- 06/01/2017 Medical Policy Committee review
  - 06/21/2017 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
  - 06/07/2018 Medical Policy Committee review
  - 06/20/2018 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
  - 06/06/2019 Medical Policy Committee review
  - 06/19/2019 Medical Policy Implementation Committee approval. Added “or retention” to differentiate incontinence “or” retention. Neurologic condition examples added (e.g., Parkinson’s disease, multiple sclerosis, spinal cord injury) to the Urinary Incontinence and Non-obstructive Retention section. Clarified coverage sections in the criteria bullets. Added two investigational statements to address when patient selection criteria are not met.
  - 06/04/2020 Medical Policy Committee review
  - 06/10/2020 Medical Policy Implementation Committee approval. No change to coverage. New FDA information added.
  - 06/03/2021 Medical Policy Committee review
  - 06/09/2021 Medical Policy Implementation Committee approval. No change to coverage.
  - 06/02/2022 Medical Policy Committee review
  - 06/08/2022 Medical Policy Implementation Committee approval. No change to coverage.
- Next Scheduled Review Date: 06/2023

### **Coding**

*The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)‡, copyright 2021 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.*

*The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability*

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Sacral Nerve Neuromodulation/Stimulation

Policy # 00108

Original Effective Date: 03/25/2002

Current Effective Date: 07/11/2022

*attributable or related to any use, nonuse or interpretation of information contained in Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.*

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	64561, 64581, 64585, 64590, 64595, 95970, 95971, 95972
HCPCS	A4290, C1767, C1778, C1816, C1820, C1883, C1897, E0745, L8680, L8685, L8686, L8687, L8688, L8689
ICD-10 Diagnosis	N39.3, N39.41-N39.46, N39.490, N39.498, R15.0-R15.9, R32, R33.0-R33.9, R39.14

\*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Sacral Nerve Neuromodulation/Stimulation

Policy # 00108

Original Effective Date: 03/25/2002

Current Effective Date: 07/11/2022

diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:

1. Consultation with technology evaluation center(s);
2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
3. Reference to federal regulations.

**\*\*Medically Necessary (or “Medical Necessity”)** - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. In accordance with nationally accepted standards of medical practice;
- B. Clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and
- C. Not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.

For these purposes, “nationally accepted standards of medical practice” means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

‡ Indicated trademarks are the registered trademarks of their respective owners.

**NOTICE:** If the Patient’s health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

**NOTICE:** Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.