



Louisiana

Select Drug Quantity Management

Policy # 00625

Original Effective Date: 01/01/2019

Current Effective Date: 11/09/2020

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Note: Topical Antipruritics coverage criteria (NON-QUANTITY related) are addressed separately in medical policy 00581.

Note: Topical Immunomodulators coverage criteria (NON-QUANTITY related) are addressed separately in medical policy 00524.

Note: Topical Actinic Keratosis Products coverage criteria (NON-QUANTITY related) are addressed separately in medical policy 00579.

Note: Topical Antifungals coverage criteria (NON-QUANTITY related) are addressed separately in medical policy 00527.

Note: Topical Corticosteroids coverage criteria (NON-QUANTITY related) are addressed separately in medical policy 00318.

Note: Topical Anesthetics coverage criteria (NON-QUANTITY related) are addressed separately in medical policy 00580.

When Services May Be Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Based on review of available data, the Company may consider quantity override requests for the drugs included in this policy to be **eligible for coverage**** when the requested drug's quantity override criteria are met.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Select Drug Quantity Management

Policy # 00625

Original Effective Date: 01/01/2019

Current Effective Date: 11/09/2020

Patient Selection Criteria

Quantity override requests for the drugs included in this policy will be considered when the selected drug's criteria are met:

- **Topical Antipruritics:**

Drugs	Quantity Override per 30 Days	Quantity Override per 90 Days	Timeframe
doxepin 5% cream	90 gm	N/A	30 days
prudoxin 5% cream	90 gm	N/A	30 days
Zonalon [®] ‡ (doxepin) 5% cream	90 gm	N/A	30 days

- Patient is treating greater than 9% of body surface area (BSA) OR patient requires two, 8 day treatment periods per 30 days.

- **Topical Atopic Dermatitis Products:**

Drugs	Quantity Override per 30 Days	Quantity Override per 90 Days	Timeframe
Elidel [®] ‡ (pimecrolimus) 1% cream	200 gm	600 gm	12 months
pimecrolimus 1% cream	200 gm	600 gm	12 months
Protopic [®] ‡ (tacrolimus) 0.03%, 0.1% ointment	200 gm	600 gm	12 months
tacrolimus 0.03%, 0.1% ointment	200 gm	600 gm	12 months
Eucrisa [®] ‡ (crisaborole) 2% ointment	240 gm	720 gm	12 months

- Patient is treating greater than 9% of BSA OR patient is applying the selected drug more frequently than twice per day.

- **Topical Actinic Keratosis Products:**

Drugs	Quantity Override per 28 Days	Quantity Override per 84 Days	Timeframe
diclofenac 3% gel	100 gm per EACH ADDITIONAL	100 gm per EACH ADDITIONAL	12 months

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Select Drug Quantity Management

Policy # 00625

Original Effective Date: 01/01/2019

Current Effective Date: 11/09/2020

	THREE 5 cm x 5 cm sites treated	THREE 5 cm x 5 cm sites treated	
Solaraze ^{®†} (diclofenac) 3% gel	100 gm per EACH ADDITIONAL THREE 5 cm x 5 cm sites treated	100 gm per EACH ADDITIONAL THREE 5 cm x 5 cm sites treated	12 months

- Patient is treating MORE than THREE 5 cm x 5 cm actinic keratosis lesions.

- **Topical Antifungals:**

Drugs	Quantity Override per 28 Days	Quantity Override per 84 Days	Timeframe
Mentax ^{®†} (butenafine) 1% cream	30 gm	N/A	28 days
Loprox ^{®†} (ciclopirox) 0.77% cream	90 gm	N/A	28 days
ciclopirox 0.77% cream	90 gm	N/A	28 days
ciclodan 0.77% cream	90 gm	N/A	28 days
ciclopirox 0.77% gel	45 gm	N/A	28 days
Loprox (ciclopirox) 0.77% suspension	60 mL	N/A	28 days
ciclopirox 0.77% suspension	60 mL	N/A	28 days
clotrimazole 1% cream	45 gm	N/A	28 days
clotrimazole 1% solution	30 mL	N/A	28 days
Lotrisone ^{®†} (clotrimazole-betamethasone) 1%/0.05% cream	45 gm	N/A	28 days
clotrimazole-betamethasone 1%/0.05% cream	45 gm	N/A	28 days
clotrimazole-betamethasone 1%/0.05% lotion	60 mL	N/A	28 days
econazole 1% cream	85 gm	N/A	28 days
Ecoza ^{™†} (econazole) 1% foam	70 gm	N/A	28 days

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Select Drug Quantity Management

Policy # 00625

Original Effective Date: 01/01/2019

Current Effective Date: 11/09/2020

ketoconazole 2% cream	60 gm	N/A	28 days
Extina [®] ‡ (ketoconazole) 2% foam	100 gm	N/A	28 days
ketoconazole 2% foam	100 gm	N/A	28 days
Xolegel [®] ‡ (ketoconazole) 2% gel	45 gm	N/A	28 days
Luzu [®] ‡ (luliconazole) 1% cream	60 gm	N/A	28 days
Naftin [®] ‡ (naftifine) 1%, 2% cream	60 gm	N/A	28 days
naftifine 1%, 2% cream	60 gm	N/A	28 days
Naftin (naftifine) 1%, 2% gel	60 gm	N/A	28 days
Oxistat [®] ‡ (oxiconazole) 1% cream	60 gm	N/A	28 days
oxiconazole 1% cream	60 gm	N/A	28 days
Oxistat (oxiconazole) 1% lotion	60 mL	N/A	28 days
Ertaczo [®] ‡ (sertaconazole) 2% cream	60 gm	N/A	28 days
Exelderm [®] ‡ (sulconazole) 1% cream	60 gm	N/A	28 days
Exelderm (sulconazole) 1% solution	60 mL	N/A	28 days
nystatin 100,000 units/gm powder	120 gm	N/A	28 days
nystop 100,000 units/gm powder	120 gm	N/A	28 days
nyamyc 100,000 units/gm powder	120 gm	N/A	28 days
nystatin 100,000 units/gm cream	30 gm	N/A	28 days
nystatin 100,000 units/gm ointment	30 gm	N/A	28 days

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Select Drug Quantity Management

Policy # 00625

Original Effective Date: 01/01/2019

Current Effective Date: 11/09/2020

nystatin-triamcinolone 100,000units/0.1% cream	60 gm	N/A	28 days
nystatin-triamcinolone 100,000units/0.1% ointment	60 gm	N/A	28 days
ANY other topical antifungal product	NO OVERRIDES ALLOWED	NO OVERRIDES ALLOWED	NO OVERRIDES ALLOWED

- Patient is treating greater than 9% of BSA OR patient is treating the condition for longer than 14 days.

- **Topical Anti-inflammatory Products:**

Drugs	Quantity Override	Quantity Override	Timeframe
Clobex ^{®‡} (clobetasol) 0.05% lotion	118 mL per 28 days	N/A	28 days
clobetasol 0.05% lotion	118 mL per 28 days	N/A	28 days
Clobex (clobetasol) 0.05% spray	125 mL per 28 days	N/A	28 days
clobetasol 0.05% spray	125 mL per 28 days	N/A	28 days
Vanos ^{®‡} (fluocinonide) cream 0.1%	180 gm per 30 days	540 gm per 90 days	12 months
fluocinonide 0.1% cream	180 gm per 30 days	540 gm per 90 days	12 months
ANY other topical clobetasol or fluocinonide product	NO OVERRIDES ALLOWED	NO OVERRIDES ALLOWED	NO OVERRIDES ALLOWED

- clobetasol products (lotion and spray ONLY):
 - Patient's condition has not sufficiently improved after the initial two weeks of therapy with the requested product.
- fluocinonide products (0.1% cream ONLY [Vanos, generic]):
 - Patient is treating greater than 8% of BSA OR patient is applying the selected drug more frequently than twice per day.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Select Drug Quantity Management

Policy # 00625

Original Effective Date: 01/01/2019

Current Effective Date: 11/09/2020

- **Topical Lidocaine Products:**

Drugs	Quantity Override per 30 Days	Quantity Override per 90 Days	Timeframe
lidocaine 5% ointment	150 gm	N/A	12 months
Xylocaine (lidocaine) 2% jelly	1800 mL	N/A	12 months
lidocaine 2% jelly	1800 mL	N/A	12 months
glydo (lidocaine) 2% jelly	1800 mL	N/A	12 months
EMLA (lidocaine/prilocaine) 2.5%/2.5% cream	30 gm for each instance of criteria	N/A	12 months
lidocaine/prilocaine 2.5%/2.5% cream	30 gm for each instance of criteria	N/A	12 months
ANY other lidocaine or prilocaine topical product	NO OVERRIDES ALLOWED	NO OVERRIDES ALLOWED	NO OVERRIDES ALLOWED

- lidocaine 5% ointment:
 - Patient is using to produce anesthesia of accessible mucous membranes of the oropharynx greater than 2% of BSA OR patient administers more frequently than two times a day.
- lidocaine 2% jelly, glydo, Xylocaine:
 - Patient performs self-catheterization on a routine basis.
- lidocaine 2.5%/prilocaine 2.5% cream, EMLA:
 - Patient needs topical anesthesia for greater than 12 separate dermal procedures (intravenous cannulation and venipuncture) utilizing 2.5 grams OR 6 separate dermal procedures (intravenous cannulation and venipuncture) utilizing 5 grams.

When Services Are Considered Not Medically Necessary

Based on review of available data, the Company considers quantity override requests for the drugs included in this policy when the patient selection criteria are not met to be **not medically necessary**.**

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Select Drug Quantity Management

Policy # 00625

Original Effective Date: 01/01/2019

Current Effective Date: 11/09/2020

Background/Overview

Topical Antipruritics/Topical Atopic Dermatitis Products

Doxepin topical cream is indicated for the short-term (up to 8 days) management of moderate pruritus in adult patients with atopic dermatitis or lichen simplex chronicus. A thin film of doxepin cream should be applied four times each day with at least a 3 to 4 hour interval between applications. There are no data to establish the safety and effectiveness of doxepin cream when used for greater than 8 days. Chronic use beyond eight days may result in higher systemic levels and should be avoided. Use of doxepin cream for longer than 8 days may result in an increased likelihood of contact sensitization.

Elidel (pimecrolimus cream) and Protopic (tacrolimus ointment) are indicated as second-line therapy for the short-term and non-continuous chronic treatment of moderate to severe atopic dermatitis in non-immunocompromised adults and children who have failed to respond adequately to other topical prescription treatments for atopic dermatitis, or when those treatments are not advisable. Eucrisa (crisaborole) is indicated for topical treatment of mild to moderate atopic dermatitis.

In clinical trials of patients with atopic dermatitis, 75-80% had disease affecting the face and/or neck region. The most common areas of the body affected by atopic dermatitis are the face, chest and back of scalp in infants and young children. In older children and adults, the front of elbows, behind the knees, face, palms of hands and soles of feet are most commonly affected. The head and neck region, upper or lower chest, each leg, or each arm comprise approximately 9% of BSA. References related to the quantity of topical creams and ointments needed to treat the involved BSA of various dermatoses estimate that between 85 - 135 grams of ointment or cream would be needed to cover a 9% BSA region when applying two times daily for one month.

Topical Actinic Keratosis Products

Solaraze Gel (diclofenac 3%) is applied to lesion areas twice daily. It is to be smoothed onto the affected skin gently. The amount needed depends upon the size of the lesion site. Normally 0.5 gm of gel is used on each 5 cm x 5 cm lesion site. The recommended duration of therapy is from 60 days to 90 days. Complete healing of the lesion(s) or optimal therapeutic effect may not be evident for up to 30 days following cessation of therapy. Lesions that do not respond to therapy should be carefully re-evaluated and management reconsidered.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Select Drug Quantity Management

Policy # 00625

Original Effective Date: 01/01/2019

Current Effective Date: 11/09/2020

Topical Antifungals

Topical antifungal products are used to treat a variety of superficial fungal infections (e.g., tinea, candida) diaper dermatitis, and seborrheic dermatitis. Frequency of administration is typically one to two times daily. Duration of treatment varies depending on the fungus being treated, but is most often used for an initial two week period or less. Treatment can last for up to four weeks in some cases if no clinical improvement is seen after two weeks of treatment.

The quantity limits for topical antifungal products supplies a sufficient quantity for each of the topical antifungal products to treat 9% of a patient's BSA when applied up to twice a day for 14 days. For patients treating a larger surface area or for a longer duration than 14 days, additional quantities listed are available through coverage review

Topical Anti-inflammatory Products

Clobetasol lotion and spray are indicated for the relief of the inflammatory and pruritic manifestations of corticosteroid-responsive dermatoses, including moderate-to-severe plaque psoriasis. Clobetasol is a super-high potency topical corticosteroid; therefore, treatment should be limited to 2 consecutive weeks and amounts greater than 50 grams or 50 ml per week should not be used.

Vanos (fluocinonide cream) is indicated for the relief of the inflammatory and pruritic manifestations of corticosteroid-responsive dermatoses, including moderate-to-severe plaque psoriasis. Moderate-to-severe psoriasis is typically defined as involvement of more than 5 to 10 percent of the BSA (the entire palmar surface, including fingers, of one hand is approximately 1 percent of the BSA) or involvement of the face, palm or sole, or disease that is otherwise disabling. Patients with more than 5 to 10 percent BSA affected are generally candidates for phototherapy or systemic therapy, since application of topical agents to a large area is not usually practical or acceptable for most patients.

Topical Lidocaine Products

Lidocaine ointment is indicated for production of anesthesia of accessible mucous membranes of the oropharynx. It is also useful as an anesthetic lubricant for intubation and for the temporary relief of pain associated with minor burns, including sunburn, abrasions of the skin, and insect bites. While there is no frequency of administration listed in the prescribing information, medical literature reports typical administration of two times daily. The initial quantity limit is enough drug to cover 2% of the BSA when applying two times daily for one month.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Select Drug Quantity Management

Policy # 00625

Original Effective Date: 01/01/2019

Current Effective Date: 11/09/2020

Xylocaine (lidocaine HCl) 2% Jelly is indicated for 1) prevention and control of pain in procedures involving the male and female urethra, 2) for topical treatment of painful urethritis, 3) and as an anesthetic lubricant for endotracheal intubation (oral and nasal). Prior to catheterization, small volumes of 5 to 10 mL (100 to 200 mg) are usually adequate for lubrication. For surface anesthesia of the female adult urethra, 3 to 5 ml (60 to 100 mg) is instilled into the urethra. No more than 600 mg (30 ml) of lidocaine 2% should be administered in any 12 hour period for any of the listed indications.

For lidocaine jelly and ointment, no overrides are recommended for patients with peripheral or post-herpetic neuralgia, post-traumatic peripheral neuropathy, or peripheral diabetic neuropathy. Cochrane reviewed three trials that utilized lidocaine 8% spray or 5% gel in patients with peripheral herpetic neuralgia (PHN) or post-traumatic peripheral neuropathy, and a fourth trial where lidocaine 5% cream had been applied twice daily for 1 week in 30 patients who had PHN, peripheral diabetic neuropathy, or post-traumatic neuropathy. Based on this review, none of the non-patch lidocaine alternatives can be recommended as therapeutic options for treatment of peripheral neuropathic pain due to the relative absence of data.

EMLA, lidocaine and prilocaine cream (a eutectic mixture of lidocaine 2.5% and prilocaine 2.5%), is indicated as a topical anesthetic for use on normal intact skin for local analgesia of minor procedures such as intravenous cannulation and venipuncture, major dermal procedures such as split thickness skin graft harvesting, and genital mucous membranes for superficial minor surgery and as pretreatment for infiltration anesthesia. The amount of cream required for topical anesthesia during a minor dermal procedure, such as intravenous cannulation and venipuncture is 2.5 grams according to prescribing information. The initial covered quantity is enough drug to allow for 6 separate dermal procedures utilizing 5 grams or 12 separate dermal procedures utilizing 2.5 grams.

Rationale/Source

The intent of this policy is to prevent stockpiling, misuse, and/or overuse while also allowing appropriate quantities for clinically acceptable and relevant use.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Select Drug Quantity Management

Policy # 00625

Original Effective Date: 01/01/2019

Current Effective Date: 11/09/2020

References

1. Elidel 1% cream [prescribing information]. Bridgewater, NJ: Valeant Pharmaceuticals North America, LLC, January 2017.
2. Protopic 0.03% and 0.1% ointment [prescribing information]. Madison, NJ: Leo Pharma, Inc., November 2016.
3. Zonalon 5% cream [prescribing information]. Newtown, PA: Presthium Pharma, Inc., October 2014.
4. Schneider L, Tilles S, Lio P, et al. Atopic Dermatitis: A Practice Parameter Update. J Allergy Clin Immunol 2013; 131:295-9. Available at: <https://www.aaaai.org/Aaaai/media/MediaLibrary/PDF%20Documents/Practice%20and%20Parameters/Atopic-dermatitis-2013.pdf>. Accessed May 10, 2017.
5. Dermatologic therapies – Basic Dermatology Curriculum. American Academy of Dermatology. Available at: <https://www.aad.org/File%20Library/Main%20navigation/Education/Basic%20Derm%20Curriculum/PDFs/Topical-Dermatologic-Therapies.pdf>. Last updated January 2016. Accessed May 10, 2017.
6. Nelson A, Miller A, Fleischer A, Balkrishnan R, Feldman S. How much of a topical agent should be prescribed for children of different sizes? J Derm Treat 2006; 17:224-228.
7. Prudoxin 5% cream [prescribing information]. Newtown, PA: Presthium Pharma, Inc., February 2015.
8. Long CC, Finlay AY. The finger-tip unit—a new practical measure. Clin Exp Dermatol 1991 Nov; 16(6):444-7.
9. McPhee SJ, Papadakis MA: Current Medical Diagnosis and Treatment 2010, 49th Edition. Figure 37.2: 3589-90. Available at: <http://sana.nahad.ir/uploads/sana/Download/Ketab/Pezeshki/CURRENT.pdf>. Accessed May 16, 2016.
10. Solaraze 3% gel [prescribing information]. Melville, NY: Fougera Pharmaceuticals, Inc. April 2016.
11. Mentax 1% Cream [prescribing information]. Morgantown, WV: Mylan Pharmaceuticals, Inc. November 2013.
12. Loprox 0.77% Cream [prescribing information]. West Fairfield, NJ: Medimetriks Pharmaceuticals, Inc. January 2016.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Select Drug Quantity Management

Policy # 00625

Original Effective Date: 01/01/2019

Current Effective Date: 11/09/2020

13. Ciclopirox 0.77% gel [prescribing information]. Mahwah, NJ: Glenmark Pharmaceuticals Ltd. September 2014.
14. Loprox 1% Shampoo [prescribing information]. Bridgewater, NJ: Valeant Pharmaceuticals North America LLC. February 2017.
15. Loprox 0.77% Suspension [prescribing information]. West Fairfield, NJ: Medimetriks Pharmaceuticals, Inc. March 2016.
16. Clotrimazole 1% cream [prescribing information]. Mahwah, NJ: Glenmark Pharmaceuticals Ltd. September 2014.
17. Clotrimazole 1% solution [prescribing information]. North Wales, PA: Teva Pharmaceuticals USA, Inc. December 2015.
18. Lotrisone cream [prescribing information]. Whitehouse Station, NJ: Merck & Co., Inc. August 2015.
19. Clotrimazole/betamethasone solution [prescribing information]. Melville, NY: Fougera Pharmaceuticals, Inc. December 2015.
20. Econazole 1% cream [prescribing information]. Pine Brook, NJ: Alvogen, Inc. January 2017.
21. Ecoza 1% foam [prescribing information]. Florham Park, NJ: Exeltis USA Dermatology, LLC. July 2016.
22. Ketoconazole 2% cream [prescribing information]. Melville, NY: Fougera Pharmaceuticals, Inc. May 2012.
23. Extina 2% foam [prescribing information]. Newtown, PA: Prestium Pharma, Inc. June 2013.
24. Xolegel 2% gel [prescribing information]. West Chester, PA: Aqua Pharmaceuticals, September 2013.
25. Nizoral 2% shampoo [prescribing information]. Titusville, NJ: Janssen Pharmaceuticals, Inc. October 2016.
26. Luzu 1% cream [prescribing information]. Bridgewater, NJ: Valeant Pharmaceuticals North America LLC. February 2017.
27. Vusion Ointment [prescribing information]. Newtown, PA: Prestium Pharma, Inc. October 2013.
28. Naftin 1% and 2% cream [prescribing information]. Greensboro, NC: Merz Pharmaceuticals. April 2011.
29. Naftin 1% and 2% gel [prescribing information]. Greensboro, NC: Merz Pharmaceuticals. April 2011.
30. Nystatin cream [prescribing information]. South Plainfield, NJ: G&W Laboratories, Inc. August 2015.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Select Drug Quantity Management

Policy # 00625

Original Effective Date: 01/01/2019

Current Effective Date: 11/09/2020

31. Nystatin ointment [prescribing information]. Melville, NY: Fougera Pharmaceuticals, Inc. August 2011.
32. Nystatin-triamcinolone cream [prescribing information]. Parsippany, NJ: ActavisPharma, Inc. May 2015.
33. Nystatin-triamcinolone ointment [prescribing information]. Parsippany, NJ: ActavisPharma, Inc. May 2015.
34. Oxistat 1% cream and lotion [prescribing information]. Melville, NY: Fougera Pharmaceuticals, Inc. January 2012.
35. Ertaczo 2% cream [prescribing information]. Bridgewater, NJ: Valeant Pharmaceuticals North America LLC. January 2014.
36. Exelderm 1% cream [prescribing information]. Jacksonville, FL: Ranbaxy Laboratories, Inc. December 2012.
37. Exelderm 1% solution [prescribing information].]. Jacksonville, FL: Ranbaxy Laboratories, Inc. March 2013.
38. Lidocaine Ointment [prescribing information]. Madisonville, LA: Solubiomix, January 2016.
39. Derry S, Wiffen PJ, Moore RA et al. Topical Lidocaine for Neuropathic Pain in Adults (Review). Cochrane Database of Systemic Reviews 2014: 7: 1-41.
40. Dermatologic therapies – Basic Dermatology Curriculum. American Academy of Dermatology. Available at:
<https://www.aad.org/File%20Library/Main%20navigation/Education/Basic%20Derm%20Curriculum/PDFs/Topical-Dermatologic-Therapies.pdf> . Last updated January 2016. Accessed May 16, 2016.
41. Long CC, Finlay AY. The finger-tip unit—a new practical measure. Clin Exp Dermatol 1991 Nov; 16(6):444-7.
42. Lidocaine and Prilocaine Cream [prescribing information]. Hayward, CA: Impax Generics, September 2015.
43. Xylocaine Jelly [prescribing information]. Schaumburg, IL: APP Pharmaceuticals, LLC, June 2008.
44. Fluocinonide cream, cream-emulsified base, gel, ointment, 0.05% [prescribing information]. Hawthorne, NY: Taro Pharmaceuticals, July 2011.
45. Fluocinonide solution 0.05% [prescribing information]. Hawthorne, NY: Taro Pharmaceuticals, Jan 2016.
46. Vanos 0.1% cream [prescribing information]. Scottsdale, AZ: Medicis, Aug 2015.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Select Drug Quantity Management

Policy # 00625

Original Effective Date: 01/01/2019

Current Effective Date: 11/09/2020

47. Temovate cream, ointment 0.05% [prescribing information]. Melville, NY: PharmaDerm, January 2012.
48. clobetasol gel 0.05% [prescribing information]. Alpharetta, GA: Direct RX, March 2008.
49. Temovate Scalp Application 0.05% [prescribing information]. Melville, NY: Pharmaderm, October 2008.
50. Clobetasol emollient cream 0.05% [prescribing information]. Melville, NY: Fougera Pharmaceuticals Inc., September 2016.
51. Clobex lotion 0.05% [prescribing information]. Fort Worth, TX: Galderma Laboratories, L.P., July 2014.
52. Clobex shampoo 0.05% [prescribing information]. Fort Worth, TX: Galderma Laboratories, L.P., November 2012.
53. Clobex spray 0.05% [prescribing information]. Fort Worth, TX: Galderma Laboratories, L.P., March 2013.
54. Olux foam 0.05% [prescribing information]. Newton, PA: Prestium Pharma, Inc., October 2014.
55. Olux E foam 0.05% [prescribing information]. Newton, PA: Prestium Pharma, Inc., March 2014.
56. Cormax Scalp Application 0.05% [prescribing information]. Richmond, VA: ECR Pharmaceuticals, April 2012.
57. Clodan shampoo 0.05% [prescribing information]. Fairfield, NJ: Medimetriks, April 2014.
58. Impoyz cream 0.025% [prescribing information]. Princeton, NJ: Promius Pharma LLC.; November 2017.
59. Clobetasol Topical Products Duration Limits. Express Scripts. Updated 1/30/2018.
60. Fluocinonide Topical Products Duration Limit. Express Scripts. Updated 6/2017.
61. Lidocaine Topical Products Duration Limit. Express Scripts. Updated 10/2017.
62. Topical Antifungal Products Duration Limit. Express Scripts. Updated 5/2017.
63. Topical Diclofenac Duration Limit. Express Scripts. Updated 6/2017.
64. Topical Doxepin Cream Duration Limit. Express Scripts. Updated 5/2017.
65. Topical Products for Atopic Dermatitis Duration Limit. Express Scripts. Updated 4/2017.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Select Drug Quantity Management

Policy # 00625

Original Effective Date: 01/01/2019

Current Effective Date: 11/09/2020

Policy History

Original Effective Date: 01/01/2019

Current Effective Date: 11/09/2020

10/04/2018 Medical Policy Committee review

10/17/2018 Medical Policy Implementation Committee approval. New policy.

10/03/2019 Medical Policy Committee review

10/09/2019 Medical Policy Implementation Committee approval. Added the generic for Elidel (pimecrolimus) to the policy for completeness.

10/01/2020 Medical Policy Committee review

10/07/2020 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

Next Scheduled Review Date: 10/2021

****Medically Necessary (or “Medical Necessity”)** - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. In accordance with nationally accepted standards of medical practice;
- B. Clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and
- C. Not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.

For these purposes, “nationally accepted standards of medical practice” means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

‡ Indicated trademarks are the registered trademarks of their respective owners.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Select Drug Quantity Management

Policy # 00625

Original Effective Date: 01/01/2019

Current Effective Date: 11/09/2020

NOTICE: If the Patient's health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.