



# Louisiana

## **Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease**

**Policy #** 00123

**Original Effective Date:** 06/24/2002

**Current Effective Date:** 03/08/2021

*Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the “Company”), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.*

*Note: Injectable Bulking Agents for the Treatment of Urinary and Fecal Incontinence is addressed separately in medical policy 00095.*

*Note: Endoscopic Radiofrequency Ablation or Cryoablation for Barrett’s Esophagus is addressed separately in medical policy 00261.*

### **Services Are Considered Investigational**

*Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.*

Based on review of available data, the Company considers endoscopic submucosal implantation of a prosthesis or injection of a bulking agent (polymethylmethacrylate [PMMA] beads, zirconium oxide spheres) as a treatment of gastroesophageal reflux disease (GERD) to be **investigational**.\*

Based on review of available data, the Company considers transoral incisionless fundoplication (ie, EsophyX) as a treatment of gastroesophageal reflux disease to be **investigational**.\*

Based on review of available data, the Company considers transesophageal radiofrequency to create submucosal thermal lesions of the gastroesophageal junction (ie, Stretta procedure) as a treatment of gastroesophageal reflux disease to be **investigational**.\*

### **Background/Overview**

#### **Gastroesophageal Reflux Disease**

GERD is a common disorder characterized by heartburn and other symptoms related to reflux of stomach acid into the esophagus. Nearly all individuals experience such symptoms at some point in their lives; a smaller number have chronic symptoms and are at risk for complications of GERD.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease

Policy # 00123

Original Effective Date: 06/24/2002

Current Effective Date: 03/08/2021

The prevalence of GERD has been estimated to be 10% to 20% in the Western world, with a lower prevalence in Asia.

### **Pathophysiology**

The pathophysiology of GERD involves excessive exposure to stomach acid, which occurs for several reasons. There can be an incompetent barrier between the esophagus and stomach, either due to dysfunction of the lower esophageal sphincter or incompetence of the diaphragm. Another mechanism is an abnormally slow clearance of stomach acid. In this situation, delayed clearance leads to an increased reservoir of stomach acid and a greater tendency to reflux.

In addition to troubling symptoms, some patients will have a more serious disease, which results in complications such as erosive esophagitis, dysphagia, Barrett esophagus, and esophageal carcinoma. Pulmonary complications may result from aspiration of stomach acid into the lungs and can include asthma, pulmonary fibrosis, and bronchitis, or symptoms of chronic hoarseness, cough, and sore throat.

### **Treatment**

Guidelines on the management of GERD emphasize initial medical management. Weight loss, smoking cessation, head of the bed elevation, and elimination of food triggers are all recommended in recent practice guidelines. Proton pump inhibitors have been shown to be the most effective medical treatment. In a Cochrane systematic review, van Pinxteren et al (2010) reported that proton pump inhibitors demonstrated superiority to H<sub>2</sub>-receptor antagonists and prokinetics in both network meta-analyses and direct comparisons.

### **Surgical Treatment**

The most common surgical procedure used for GERD is laparoscopic Nissen fundoplication. Fundoplication involves wrapping a portion of the gastric fundus around the distal esophagus to increase lower esophageal sphincter pressure. If a hiatal hernia is present, the procedure also restores the position of the lower esophageal sphincter to the correct location. Laparoscopic fundoplication was introduced in 1991 and has been rapidly adopted because it avoids complications associated with an open procedure.

Although fundoplication results in a high proportion of patients reporting symptom relief, complications can occur, and sometimes require conversion to an open procedure. Patients who have

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease

Policy # 00123

Original Effective Date: 06/24/2002

Current Effective Date: 03/08/2021

relief of symptoms of GERD after fundoplication may have dysphagia or gas-bloat syndrome (excessive gastrointestinal gas).

### Other Treatment Options

Due in part to the high prevalence of GERD, there has been interest in creating a minimally invasive transesophageal therapeutic alternative to open or laparoscopic fundoplication or chronic medical therapy. This type of procedure may be considered natural orifice transluminal surgery. Three types of procedures have been investigated.

1. Transesophageal endoscopic gastroplasty (gastroplication, transoral incisionless fundoplication) can be performed as an outpatient procedure. During this procedure, the fundus of the stomach is folded and then held in place with staples or fasteners that are deployed by the device. The endoscopic procedure is designed to recreate a valve and barrier to reflux.
2. Radiofrequency energy has been used to produce submucosal thermal lesions at the gastroesophageal junction. (This technique has also been referred to as the Stretta procedure.) Specifically, radiofrequency energy is applied through 4 electrodes inserted into the esophageal wall at multiple sites both above and below the squamocolumnar junction. The mechanism of action of the thermal lesions is not precisely known but may be related to the ablation of the nerve pathways responsible for sphincter relaxation or may induce a tissue-tightening effect related to heat-induced collagen contraction and fibrosis.
3. Submucosal injection or implantation of a prosthetic or bulking agent to enhance the volume of the lower esophageal sphincter has also been investigated. One bulking agent, pyrolytic carbon-coated zirconium oxide spheres (Durasphere), is being evaluated. The Gatekeeper<sup>TM</sup> Reflux Repair System (Medtronic) uses a soft, pliable, expandable prosthesis made of a polyacrylonitrile-based hydrogel. The prosthesis is implanted into the esophageal submucosa, and with time, the prosthesis absorbs water and expands, creating bulk in the region of implantation. However, as the only identified RCT was terminated early due to lack of efficacy and no information is available on the 'Digestive and Gastrointestinal' Products section of Medtronic's website, it is suspected that The Gatekeeper Reflux Repair System is not commercially available. U.S. Food and Drug Administration (FDA) product code: DQX. Endoscopic submucosal implantation of polymethylmethacrylate beads into the lower esophageal folds has also been investigated.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease

Policy # 00123

Original Effective Date: 06/24/2002

Current Effective Date: 03/08/2021

The Agency for Healthcare Research and Quality published a systematic review of management strategies for GERD in 2005, which was updated by Ip et al (2011). The 2005 comparative effectiveness review evaluated studies on the EndoCinch Suturing System, Stretta, Enteryx, and the NDO Plicator. The 2011 update excluded Enteryx and the NDO Plicator, because they were no longer available in the US, and added the EsophyX procedure (endoscopic fundoplication), which was commercialized after the 2005 review. The 2011 report concluded that, for the 3 available endoscopic procedures (EndoCinch, Stretta, EsophyX), effectiveness remained substantially uncertain for the long-term management of GERD. All procedures have been associated with complications, including dysphagia, infection/fever, and bloating, although bloating and dysphagia are also adverse events of laparoscopic fundoplication. A review of endoscopic treatment of GERD by Hummel and Richards (2015) noted that EndoCinch is no longer manufactured.

## **FDA or Other Governmental Regulatory Approval**

### **U.S. Food and Drug Administration (FDA)**

The EsophyX<sup>®†</sup> (EndoGastric Solutions) is a TIF device that was originally cleared for marketing by the FDA through the 510(k) process in 2007 and has subsequently undergone 2 evolutions: Generation 2=EsophyX2 iterations (E2-Plus, HD) and Generation 3=Z iterations (EZ/ZR, Z+). Some of the key Regulatory Status changes are summarized herein. In 2007, EsophyX (EndoGastric Solutions) was cleared for marketing by the FDA through the 510(k) process for full-thickness plication. In 2016, EsophyX<sup>®†</sup> Z Device with SerosaFuse Fasteners was cleared for marketing by the FDA through the 510(k) process (K160960) for use in transoral tissue approximation, full-thickness plication, ligation in the gastrointestinal tract, narrowing the gastroesophageal junction, and reduction of hiatal hernias of 2 cm or less in patients with symptomatic chronic GERD. In June 2017, EsophyX2 HD and the third-generation EsophyX Z Devices with SerosaFuse fasteners and accessories were cleared for marketing by the FDA through the 510(k) process (K171307) for expanded indications, including patients who require and respond to pharmacologic therapy and patients with hiatal hernias larger than 2 cm when a laparoscopic hiatal hernia repair reduces a hernia to 2 cm or less. The most recent FDA 510(k) clearance (K172811) occurred in October 2017 for new product specification iterations of EsophyX2 HD and EsophyX Z Devices. This clearance allows for "a moderate increase in the upper limit of the temporary Tissue Mold clamping pressure occurring during each fastener deployment." FDA product code: ODE.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease

Policy # 00123

Original Effective Date: 06/24/2002

Current Effective Date: 03/08/2021

The Medigus SRS Endoscopic Stapling System (MUSE, Medigus) was cleared for marketing by the FDA through the 510(k) process in 2012 (K120299) and 2014 (K132151). MUSE is intended for endoscopic placement of surgical staples in the soft tissue of the esophagus and stomach to create anterior partial fundoplication for the treatment of symptomatic chronic GERD in patients who require and respond to pharmacologic therapy. FDA product code: ODE.

In 2000, the CSM Stretta<sup>®‡</sup> System was cleared for marketing by the FDA through the 510(k) process for general use in the electrosurgical coagulation of tissue and was specifically intended for use in the treatment of GERD. Stretta is currently manufactured by Mederi Therapeutics. FDA product code: GEI.

Durasphere<sup>®‡</sup> is a bulking agent approved for the treatment of urinary and fecal incontinence (see medical policy 00095). Use of this product for esophageal reflux would be considered off-label use. The website of Carbon Medical Technologies states that the Durasphere<sup>®‡</sup> GR product is “intended to treat problems associated with GERD” but is considered an investigational device in the U.S.

### **Rationale/Source**

Transesophageal endoscopic therapies are being developed for the treatment of gastroesophageal reflux disease (GERD). A variety of procedures are being evaluated, including transesophageal (or transoral) incisionless fundoplication (TIF), application of radiofrequency energy, and injection/implantation of prosthetic devices or bulking agents.

For individuals who have GERD and a hiatal hernia of 2 cm or less that is not controlled by proton pump inhibitors (PPIs) who receive TIF (eg, EsophyX), the evidence includes 2 randomized controlled trials (RCTs) comparing TIF with PPI therapy, nonrandomized studies comparing TIF with fundoplication, and case series with longer-term follow-up. Relevant outcomes are symptoms, change in disease status, quality of life (QOL), medication use, and treatment-related morbidity. The highest quality RCT (RESPECT) was sham-controlled that compared TIF with PPI therapy while the other RCT (TEMPO) compared TIF with maximum PPI therapy. Both trials found a significant benefit of TIF on the primary outcome measure in about 65% of patients. The sham-controlled trial reported improvement in 45% of the sham-controlled group and no benefit on secondary subjective outcome measures. The nonblinded RCT found significant improvements in subjective measures but no difference in objective outcome measures compared with PPI therapy. Together, these trial results

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease

Policy # 00123

Original Effective Date: 06/24/2002

Current Effective Date: 03/08/2021

would suggest a strong placebo effect of the surgery and a modest benefit of TIF in patients whose symptoms were not controlled by PPIs. For these patients, the most appropriate comparator would be laparoscopic fundoplication. Studies comparing TIF with fundoplication have limitations that include earlier TIF procedures and unbalanced groups at baseline and are inadequate to determine relative efficacy. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have GERD and a hiatal hernia of 2 cm or less that is controlled by PPIs who receive TIF (eg, EsophyX), the evidence includes 2 RCTs and observational studies with longer-term follow-up. Relevant outcomes are symptoms, change in disease status, QOL, medication use, and treatment-related morbidity. A sham-controlled trial found that the time to resume PPI therapy was longer following TIF and the remission rate was higher, indicating that TIF is more effective than no therapy. The nonblinded RCT found a benefit of TIF compared with continued PPI therapy for subjective measures, but not for the objective measures of pH normalization and esophagitis. These results raise questions about a possible placebo effect for the procedure. Also, observational studies have indicated a loss of treatment effectiveness over time. Adverse events associated with the procedure (eg, perforation) may be severe. At present, the available evidence does not support the use of this intervention in patients whose symptoms are adequately controlled by medical therapy. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have GERD who receive endoscopic radiofrequency energy (eg, Stretta), the evidence includes 2 meta-analyses, 6 small RCTs, 2 nonrandomized comparative studies, and observational studies with longer-term follow-up. Relevant outcomes are symptoms, change in disease status, QOL, medication use, and treatment-related morbidity. The RCTs reported some improvements in symptoms and QOL following treatment with radiofrequency energy compared with sham controls. However, objective measures of GERD and a meta-analysis of 4 RCTs found no significant improvements in outcomes, raising questions about the mechanism of the symptom relief. Symptom relief and clinical success is reported to be lower than after fundoplication, and reoperations and other severe adverse events greater. Larger RCTs with longer follow-up, preferably compared with fundoplication, are needed to define the risks and benefits of this procedure better. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have GERD who receive esophageal bulking agents, the evidence includes an RCT and case series. Relevant outcomes are symptoms, change in disease status, QOL, medication

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease

Policy # 00123

Original Effective Date: 06/24/2002

Current Effective Date: 03/08/2021

use, and treatment-related morbidity. The RCT for a single product was terminated early due to lack of efficacy, while other products have only case series to support use. High-quality data from large RCTs are needed to compare bulking procedures with both sham controls and with the currently accepted treatments for GERD (ie, drug therapy, laparoscopic fundoplication). Well-designed trials should use standardized outcome measures to examine whether subjective improvement (eg, discontinuation of medication therapy, Gastroesophageal Reflux Disease Health-Related Quality of Life scores) is supported by objective improvement (eg, esophageal acid exposure). The evidence is insufficient to determine the effects of the technology on health outcomes.

## **Supplemental Information**

### **Clinical Input From Physician Specialty Societies and Academic Medical Centers**

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

### **2015 Input**

In response to requests for clinical input on transesophageal radiofrequency (Stretta) as a treatment of gastroesophageal reflux disease (GERD), input was received from 1 physician specialty society (2 reviewers) and 3 academic medical centers while this policy was under review in 2015. Input was mixed on the treatment of GERD with transesophageal radiofrequency to create submucosal thermal lesions of the gastroesophageal junction (ie, Stretta). Potential conflicts of interest were noted by 2 reviewers.

### **2011 Input**

In response to requests for clinical input on transoral incisionless fundoplication (TIF) using EsophyX, input was received from 2 physician specialty societies and 4 academic medical centers while this policy was under review in 2011. Reviewers agreed that TIF differed sufficiently from laparoscopic Nissen fundoplication to warrant evaluation as a separate procedure. Reviewers considered TIF (ie, EsophyX) to be investigational for the treatment of GERD.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease

Policy # 00123

Original Effective Date: 06/24/2002

Current Effective Date: 03/08/2021

### **Practice Guidelines and Position Statements**

#### **American Society for Gastrointestinal Endoscopy**

In 2015, the American Society for Gastrointestinal Endoscopy published guidelines on endoscopic procedures for GERD. In its review of the EsophyX and Stretta procedures, the Society noted some positive findings but discrepancies between subjective and objective outcome measures or a lack of objective outcome measures in reported trials, concluding that these techniques represent “potentially new therapeutic indications for GI endoscopy”, but that prospective trials using objective measures of GERD as the primary endpoint could be useful in defining the clinical role of these procedures.

#### **American College of Gastroenterology**

In 2013, updated guidelines from the American College of Gastroenterology indicated the use of current endoscopic therapy or TIF could not be recommended as an alternative to medical or traditional surgical therapy (conditional recommendation, moderate level of evidence). The guidelines also cited limited data on small numbers of subjects and short duration of follow-up.

#### **Society of American Gastrointestinal and Endoscopic Surgeons**

In 2017, the Society of American Gastrointestinal and Endoscopic Surgeons (SAGES) provided a clinical spotlight review on endoluminal treatments for GERD. The SAGES gave a strong recommendation based on moderate-quality evidence that TIF using EsophyX can be performed with an acceptable safety risk in selected patients. The SAGES concluded that EsophyX results in better control of GERD symptoms than proton pump inhibitor (PPI) treatment in the short term (6 months), and leads to similar improvements in objective GERD measures compared with PPIs. TIF appears to lose effectiveness during longer-term follow-up and is associated with moderate patient satisfaction scores. SAGES found no comparative, controlled trials between TIF and surgical fundoplication, but preliminary evidence suggested that surgical fundoplication can be used safely after TIF failure.

The SAGES gave a strong recommendation based on moderate-quality evidence that Stretta is safe for adults and significantly improves health-related quality of life score, heartburn scores, the incidence of esophagitis, and esophageal acid exposure in patients with GERD. Stretta was found to decrease PPI use by about 50%, and be more effective than PPIs, but less effective compared to fundoplication. The effectiveness of the procedure decreases over time.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.





# Louisiana

## Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease

Policy # 00123

Original Effective Date: 06/24/2002

Current Effective Date: 03/08/2021

### **American Society of General Surgeons**

In 2011, the American Society of General Surgeons issued a position statement on transoral fundoplication stating that "ASGS supports the use of transoral fundoplication by trained General Surgeons for the treatment of symptomatic chronic gastroesophageal reflux disease (GERD) in patients who fail to achieve satisfactory response to a standard dose of Proton Pump Inhibitor (PPI) therapy or for those who wish to avoid the need for a lifetime of medication dependence."

### **National Institute for Health and Care Excellence**

In 2013, the National Institute for Health and Care Excellence (NICE) updated its guidance on endoscopic radiofrequency treatment for GERD, concluding: "The evidence on the safety of endoscopic radiofrequency ablation for gastro-esophageal reflux disease is adequate in the short and medium term, but there is uncertainty about longer-term outcomes. With regard to efficacy, there is evidence of symptomatic relief, but objective evidence on reduction of reflux is inconclusive....." The NICE noted "concern on the part of some specialists about the possibility that symptoms may improve as a result of denervation caused by the procedure; if that were the case then failure to recognize and treat reflux might lead to complications in the long term."

In 2011, the NICE issued guidance on endoluminal gastroplication for GERD, concluding that "The evidence on endoluminal gastroplication for gastroesophageal reflux disease raises no major safety concerns. Evidence from a number of RCTs [randomized controlled trials] shows a degree of efficacy in terms of reduced medication requirement in the short term, but changes in other efficacy outcomes are inconsistent, and there is no good evidence of sustained improvement in esophageal pH measurements...."

In 2004, the NICE issued guidance on bulking agents for GERD, concluding that "Current evidence on the safety and efficacy of endoscopic injection of bulking agents for gastro-esophageal reflux disease does not appear adequate for this procedure to be used without special arrangements....." The NICE (2016) removed guidance on endoscopic bulking agents/hydrogel implants from guidelines on treatment for "dyspepsia and gastro-esophageal reflux" because the product had been withdrawn by the manufacturer.

### **U.S. Preventive Services Task Force Recommendations**

Not applicable.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease

Policy # 00123

Original Effective Date: 06/24/2002

Current Effective Date: 03/08/2021

**Medicare National Coverage**

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

**Ongoing and Unpublished Clinical Trials**

Some currently ongoing and unpublished trials that might influence this review are listed in Table 1.

**Table 1. Summary of Key Trials**

<b>NCT No.</b>	<b>Trial Name</b>	<b>Planned Enrollment</b>	<b>Completion Date (status)</b>
<i>Ongoing</i>			
NCT04306380	Transoral Incisionless Fundoplication Database Repository (TIF)	500	Dec 2030
<i>Unpublished</i>			
NCT01110811 <sup>a</sup>	A Randomized Controlled Trial Comparing Transoral Incisionless Fundoplication (TIF) Using EsophyX With Sham Procedure for the Treatment of PPI Dependent GERD: the TIF vs Sham Study	60	Dec 2018 (completed)
NCT01118585 <sup>a</sup>	Prospective Outcome Evaluation of Transoral Incisionless Fundoplication (TIF) for the Treatment of Gastroesophageal Reflux Disease (GERD): The TIF Registry Study	278	Dec 2018 (completed)
NCT02366169 <sup>a</sup>	A Worldwide Post-Market Surveillance Registry to Assess the Medigus Ultrasonic Surgical Endostapler (MUSE™) System for the Treatment of GERD	200	Dec 2019 (unknown)

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease

Policy # 00123

Original Effective Date: 06/24/2002

Current Effective Date: 03/08/2021

NCT: national clinical trial.

<sup>a</sup> Denotes industry-sponsored or cosponsored trial.

### **References**

1. Blue Cross and Blue Shield Association, Medical Policy Reference Manual, “Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease”, 2.01.38, January 2021.
2. Katz PO, Gerson LB, Vela MF. Guidelines for the diagnosis and management of gastroesophageal reflux disease. *Am J Gastroenterol*. Mar 2013; 108(3): 308-28; quiz 329. PMID 23419381
3. van Pinxteren B, Sigterman KE, Bonis P, et al. Short-term treatment with proton pump inhibitors, H2-receptor antagonists and prokinetics for gastro-oesophageal reflux disease-like symptoms and endoscopy negative reflux disease. *Cochrane Database Syst Rev*. Nov 10 2010; (11): CD002095. PMID 21069670
4. Ip S, Bonis P, Tatsoni A, et al. Comparative Effectiveness of Management Strategies for Gastroesophageal Reflux Disease (Evidence Report/Technology Assessment No. 1). Rockville, MD: Agency for Healthcare Research and Quality; 2005.
5. Ip S, Chung M, Moorthy D, et al. Management strategies for gastroesophageal reflux disease: An update (Comparative Effectiveness Review No. 29). Rockville, MD: Agency for Healthcare Research and Quality; 2011.
6. Humphries LA, Hernandez JM, Clark W, et al. Causes of dissatisfaction after laparoscopic fundoplication: the impact of new symptoms, recurrent symptoms, and the patient experience. *Surg Endosc*. May 2013; 27(5): 1537-45. PMID 23508812
7. Hummel K, Richards W. Endoscopic treatment of gastroesophageal reflux disease. *Surg Clin North Am*. Jun 2015; 95(3): 653-67. PMID 25965137
8. Ihde GM. The evolution of TIF: transoral incisionless fundoplication. *Therap Adv Gastroenterol*. 2020; 13: 1756284820924206. PMID 32499834
9. Food and Drug Administration (FDA). 510(k) Summary: EsophyX (K160960). 2016; [https://www.accessdata.fda.gov/cdrh\\_docs/pdf16/K160960.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf16/K160960.pdf).
10. Food and Drug Administration (FDA). EsophyX Summary K171307. 2017; [https://www.accessdata.fda.gov/cdrh\\_docs/pdf17/K171307.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf17/K171307.pdf).
11. Food and Drug Administration (FDA). EsophyX Summary K172811. 2017; [https://www.accessdata.fda.gov/cdrh\\_docs/pdf17/K172811.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf17/K172811.pdf).

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease

Policy # 00123

Original Effective Date: 06/24/2002

Current Effective Date: 03/08/2021

12. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Transesophageal Endoscopic Treatments for Gastroesophageal Reflux Disease. TEC Assessment. 2003;Volume 18:Tab 20. PMID
13. McCarty TR, Itidiare M, Njei B, et al. Efficacy of transoral incisionless fundoplication for refractory gastroesophageal reflux disease: a systematic review and meta-analysis. *Endoscopy*. Jul 2018; 50(7): 708-725. PMID 29625507
14. Richter JE, Kumar A, Lipka S, et al. Efficacy of Laparoscopic Nissen Fundoplication vs Transoral Incisionless Fundoplication or Proton Pump Inhibitors in Patients With Gastroesophageal Reflux Disease: A Systematic Review and Network Meta-analysis. *Gastroenterology*. Apr 2018; 154(5): 1298-1308.e7. PMID 29305934
15. Hunter JG, Kahrilas PJ, Bell RC, et al. Efficacy of transoral fundoplication vs omeprazole for treatment of regurgitation in a randomized controlled trial. *Gastroenterology*. Feb 2015; 148(2): 324-333.e5. PMID 25448925
16. Trad KS, Barnes WE, Simoni G, et al. Transoral incisionless fundoplication effective in eliminating GERD symptoms in partial responders to proton pump inhibitor therapy at 6 months: the TEMPO Randomized Clinical Trial. *Surg Innov*. Feb 2015; 22(1): 26-40. PMID 24756976
17. Trad KS, Fox MA, Simoni G, et al. Transoral fundoplication offers durable symptom control for chronic GERD: 3-year report from the TEMPO randomized trial with a crossover arm. *Surg Endosc*. Jun 2017; 31(6): 2498-2508. PMID 27655380
18. Trad KS, Barnes WE, Prevou ER, et al. The TEMPO Trial at 5 Years: Transoral Fundoplication (TIF 2.0) Is Safe, Durable, and Cost-effective. *Surg Innov*. Apr 2018; 25(2): 149-157. PMID 29405886
19. Hakansson B, Montgomery M, Cadiere GB, et al. Randomised clinical trial: transoral incisionless fundoplication vs. sham intervention to control chronic GERD. *Aliment Pharmacol Ther*. Dec 2015; 42(11-12): 1261-70. PMID 26463242
20. Witteman BP, Conchillo JM, Rinsma NF, et al. Randomized controlled trial of transoral incisionless fundoplication vs. proton pump inhibitors for treatment of gastroesophageal reflux disease. *Am J Gastroenterol*. Apr 2015; 110(4): 531-42. PMID 25823768
21. Toomey P, Teta A, Patel K, et al. Transoral incisionless fundoplication: is it as safe and efficacious as a Nissen or Toupet fundoplication?. *Am Surg*. Sep 2014; 80(9): 860-7. PMID 25197871
22. Frazzoni M, Conigliaro R, Manta R, et al. Reflux parameters as modified by EsophyX or laparoscopic fundoplication in refractory GERD. *Aliment Pharmacol Ther*. Jul 2011; 34(1): 67-75. PMID 21539587

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease

Policy # 00123

Original Effective Date: 06/24/2002

Current Effective Date: 03/08/2021

23. Stefanidis G, Viazis N, Kotsikoros N, et al. Long-term benefit of transoral incisionless fundoplication using the esophyx device for the management of gastroesophageal reflux disease responsive to medical therapy. *Dis Esophagus*. Feb 01 2017; 30(3): 1-8. PMID 27868281
24. Testoni PA, Testoni S, Mazzoleni G, et al. Long-term efficacy of transoral incisionless fundoplication with Esophyx (Tif 2.0) and factors affecting outcomes in GERD patients followed for up to 6 years: a prospective single-center study. *Surg Endosc*. Sep 2015; 29(9): 2770-80. PMID 25480624
25. Testoni PA, Testoni S, Distefano G, et al. Transoral incisionless fundoplication with Esophyx for gastroesophageal reflux disease: clinical efficacy is maintained up to 10 years. *Endosc Int Open*. May 2019; 7(5): E647-E654. PMID 31058207
26. Huang X, Chen S, Zhao H, et al. Efficacy of transoral incisionless fundoplication (TIF) for the treatment of GERD: a systematic review with meta-analysis. *Surg Endosc*. Mar 2017; 31(3): 1032-1044. PMID 27495332
27. Lipka S, Kumar A, Richter JE. No evidence for efficacy of radiofrequency ablation for treatment of gastroesophageal reflux disease: a systematic review and meta-analysis. *Clin Gastroenterol Hepatol*. Jun 2015; 13(6): 1058-67.e1. PMID 25459556
28. Corley DA, Katz P, Wo JM, et al. Improvement of gastroesophageal reflux symptoms after radiofrequency energy: a randomized, sham-controlled trial. *Gastroenterology*. Sep 2003; 125(3): 668-76. PMID 12949712
29. Aziz AM, El-Khayat HR, Sadek A, et al. A prospective randomized trial of sham, single-dose Stretta, and double-dose Stretta for the treatment of gastroesophageal reflux disease. *Surg Endosc*. Apr 2010; 24(4): 818-25. PMID 19730952
30. Arts J, Bisschops R, Blondeau K, et al. A double-blind sham-controlled study of the effect of radiofrequency energy on symptoms and distensibility of the gastro-esophageal junction in GERD. *Am J Gastroenterol*. Feb 2012; 107(2): 222-30. PMID 22108449
31. Fass R, Cahn F, Scotti DJ, et al. Systematic review and meta-analysis of controlled and prospective cohort efficacy studies of endoscopic radiofrequency for treatment of gastroesophageal reflux disease. *Surg Endosc*. Dec 2017; 31(12): 4865-4882. PMID 28233093
32. Coron E, Sebille V, Cadiot G, et al. Clinical trial: Radiofrequency energy delivery in proton pump inhibitor-dependent gastro-oesophageal reflux disease patients. *Aliment Pharmacol Ther*. Nov 01 2008; 28(9): 1147-58. PMID 18616516
33. Kalapala R, Shah H, Nabi Z, et al. Treatment of gastroesophageal reflux disease using radiofrequency ablation (Stretta procedure): An interim analysis of a randomized trial. *Indian J Gastroenterol*. Sep 2017; 36(5): 337-342. PMID 29030802

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease

Policy # 00123

Original Effective Date: 06/24/2002

Current Effective Date: 03/08/2021

34. Zerbib F, Sacher-Huvelin S, Coron E, et al. Randomised clinical trial: oesophageal radiofrequency energy delivery versus sham for PPI-refractory heartburn. *Aliment Pharmacol Ther.* Aug 2020; 52(4): 637-645. PMID 32656869
35. Liang WT, Yan C, Wang ZG, et al. Early and Midterm Outcome After Laparoscopic Fundoplication and a Minimally Invasive Endoscopic Procedure in Patients with Gastroesophageal Reflux Disease: A Prospective Observational Study. *J Laparoendosc Adv Surg Tech A.* Aug 2015; 25(8): 657-61. PMID 26258269
36. Ma L, Li T, Liu G, et al. Stretta radiofrequency treatment vs Toupet fundoplication for gastroesophageal reflux disease: a comparative study. *BMC Gastroenterol.* May 27 2020; 20(1): 162. PMID 32460696
37. Liang WT, Wang ZG, Wang F, et al. Long-term outcomes of patients with refractory gastroesophageal reflux disease following a minimally invasive endoscopic procedure: a prospective observational study. *BMC Gastroenterol.* Oct 10 2014; 14: 178. PMID 25304252
38. Noar M, Squires P, Noar E, et al. Long-term maintenance effect of radiofrequency energy delivery for refractory GERD: a decade later. *Surg Endosc.* Aug 2014; 28(8): 2323-33. PMID 24562599
39. Ganz RA, Fallon E, Wittchow T, et al. A new injectable agent for the treatment of GERD: results of the Durasphere pilot trial. *Gastrointest Endosc.* Feb 2009; 69(2): 318-23. PMID 19185691
40. Fockens P, Cohen L, Edmundowicz SA, et al. Prospective randomized controlled trial of an injectable esophageal prosthesis versus a sham procedure for endoscopic treatment of gastroesophageal reflux disease. *Surg Endosc.* Jun 2010; 24(6): 1387-97. PMID 20198491
41. Feretis C, Benakis P, Dimopoulos C, et al. Endoscopic implantation of Plexiglas (PMMA) microspheres for the treatment of GERD. *Gastrointest Endosc.* Apr 2001; 53(4): 423-6. PMID 11275880
42. Muthusamy VR, Lightdale JR, Acosta RD, et al. The role of endoscopy in the management of GERD. *Gastrointest Endosc.* 2015; 81(6): 1305-10. PMID 25863867
43. Society of American Gastrointestinal and Endoscopic Surgeons. Clinical Spotlight Review: Endoluminal Treatments for Gastroesophageal Reflux Disease (GERD). 2017; <https://www.sages.org/publications/guidelines/endoluminal-treatments-for-gastroesophageal-reflux-disease-gerd/>.
44. American Society of General Surgeons (ASGS). Coverage of Transoral fundoplication. 2011; <https://theasgs.org/position-statements/coverage-of-transoral-fundoplication-2/>.
45. National Institute for Health and Care Excellence (NICE). Endoscopic radiofrequency ablation for gastro-oesophageal reflux disease [IPG461]. 2013;

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease

Policy # 00123

Original Effective Date: 06/24/2002

Current Effective Date: 03/08/2021

<https://www.nice.org.uk/guidance/ipg461>.

46. National Institute for Health and Care Excellence (NICE). Endoluminal gastroplication for gastro-oesophageal reflux disease [IPG404]. 2011; <https://www.nice.org.uk/guidance/ipg404>.

47. National Institute for Health and Care Excellence (NICE). Endoscopic injection of bulking agents for gastro-oesophageal reflux disease [IPG55]. 2004; <https://www.nice.org.uk/guidance/ipg55>.

### **Policy History**

Original Effective Date: 06/24/2002

Current Effective Date: 03/08/2021

06/20/2002 Medical Policy Committee review.

06/24/2002 Managed Care Advisory Council approval.

04/01/2004 Medical Director review

04/20/2004 Medical Policy Committee review. Format revision. No substance change to policy.

04/26/2004 Managed Care Advisory Council approval

04/05/2006 Medical Director review

04/19/2006 Medical Policy Committee review. Format revision, including addition of FDA and or other governmental regulatory approval and rationale/source. Coverage eligibility unchanged.

04/02/2008 Medical Director review

04/16/2008 Medical Policy Committee approval. No change to coverage eligibility.

04/02/2009 Medical Director review

04/15/2009 Medical Policy Committee approval. No change to coverage eligibility. Title changed to match BCBSA.

04/08/2010 Medical Policy Committee approval

04/21/2010 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

04/07/2011 Medical Policy Committee review

04/13/2011 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

04/12/2012 Medical Policy Committee review

04/25/2012 Medical Policy Implementation Committee approval. Added NDO Plicator and EsophyX procedures as examples of transesophageal endoscopic gastroplasty.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease

Policy # 00123

Original Effective Date: 06/24/2002

Current Effective Date: 03/08/2021

	Investigational statements were combined on biocompatible polymer and PMMA beads as bulking agents.
04/04/2013	Medical Policy Committee review
04/24/2013	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
04/03/2014	Medical Policy Committee review
04/23/2014	Medical Policy Implementation Committee approval. Removed the second Investigational statement regarding transesophageal radiofrequency to create thermal lesions of the gastrointestinal junction (i.e., the Stretta procedure) as a treatment of GERD.
04/02/2015	Medical Policy Committee approval
04/20/2015	Medical Policy Implementation Committee approval. Coverage eligibility unchanged. Updated rationale and references.
08/03/2015	Coding update: ICD10 Diagnosis code section added; ICD9 Procedure code section removed.
10/01/2015	Coding update
01/01/2016	Coding update
04/07/2016	Medical Policy Committee approval
04/20/2016	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
01/01/2017	Coding update: Removing ICD-9 Diagnosis Codes
01/05/2017	Medical Policy Committee approval
01/18/2017	Medical Policy Implementation Committee approval. Removed transesophageal endoscopic gastroplasty from the policy and policy statements.
02/01/2018	Medical Policy Committee review
02/21/2018	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
02/07/2019	Medical Policy Committee review
02/20/2019	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
02/06/2020	Medical Policy Committee review
02/12/2020	Medical Policy Implementation Committee approval. Removed biocompatible liquid polymer from the current investigational statement. Added two investigational statements for endoscopic GERD procedures for transoral

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.





# Louisiana

## Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease

Policy # 00123

Original Effective Date: 06/24/2002

Current Effective Date: 03/08/2021

incisionless fundoplication (TIF) using the Esophyx device and transesophageal radiofrequency using the Stretta device.

05/05/2020 Coding update

09/11/2020 Coding update

02/04/2021 Medical Policy Committee review

02/10/2021 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

09/30/2021 Coding update

Next Scheduled Review Date: 02/2022

### **Coding**

*The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)‡, copyright 2020 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.*

*The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.*

CPT is a registered trademark of the American Medical Association.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease

Policy # 00123

Original Effective Date: 06/24/2002

Current Effective Date: 03/08/2021

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	43201, 43210, 43236, 43257, 43499
HCPCS	No codes
ICD-10 Diagnosis	K21.00-K21.01, K21.9, K22.70, K22.710-K22.711, K22.719, K22.8 Added codes eff 10/1/2021: K22.81-K22.89

\*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
  - 1. Consultation with the Blue Cross and Blue Shield Association technology assessment program (TEC) or other nonaffiliated technology evaluation center(s);
  - 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
  - 3. Reference to federal regulations.

‡ Indicated trademarks are the registered trademarks of their respective owners.

**NOTICE:** If the Patient’s health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease

Policy # 00123

Original Effective Date: 06/24/2002

Current Effective Date: 03/08/2021

**NOTICE:** Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.