



# Louisiana

## sapropterin dihydrochloride (Kuvan<sup>®</sup>, generics)

Policy # 00303

Original Effective Date: 08/17/2011

Current Effective Date: 03/08/2021

*Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.*

### When Services May Be Eligible for Coverage

*Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:*

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Based on review of available data, the Company may consider the use of sapropterin dihydrochloride (Kuvan<sup>®</sup>, generics)<sup>‡</sup> for the treatment of hyperphenylalaninemia (HPA) due to tetrahydrobiopterin- (BH4-) responsive phenylketonuria (PKU) to be **eligible for coverage**.<sup>\*\*</sup>

#### Patient Selection Criteria

Coverage eligibility for the use of sapropterin dihydrochloride (Kuvan, generics) for the treatment of HPA will be considered when both of the following criteria are met:

- Patient has a diagnosis of HPA due to BH4- PKU; AND
- Patient is on a phenylalanine (Phe) restricted diet; AND
- If the request is for brand Kuvan, patient has tried and failed (e.g., intolerance or inadequate response) GENERIC sapropterin dihydrochloride unless there is clinical evidence or patient history that suggests GENERIC sapropterin dihydrochloride will be ineffective or cause an adverse reaction to the patient.

*(Note: this specific patient criterion is an additional Company requirement for coverage eligibility and will be denied as not medically necessary<sup>\*\*</sup> if not met).*

### When Services Are Considered Not Medically Necessary

Based on review of available data, the Company considers the use of brand Kuvan when the patient has not tried and failed GENERIC sapropterin dihydrochloride to be **not medically necessary**.<sup>\*\*</sup>

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



sapropterin dihydrochloride (Kuvan<sup>®</sup>, generics)

Policy # 00303

Original Effective Date: 08/17/2011

Current Effective Date: 03/08/2021

## **When Services Are Considered Investigational**

*Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.*

Based on review of available data, the Company considers the use of sapropterin dihydrochloride (Kuvan) when patient selection criteria are not met (except the criterion denoted above as **not medically necessary\*\***) to be **investigational**.\*

## **Background/Overview**

Kuvan is a synthetic preparation of naturally occurring BH<sub>4</sub><sup>-</sup>, which is a cofactor for the enzyme phenylalanine hydroxylase (PAH). PAH hydroxylates Phe through an oxidative reaction to form tyrosine. In patients with PKU, PAH activity is deficient or absent. Treatment with BH<sub>4</sub><sup>-</sup> can activate residual PAH enzyme, improve the normal oxidative metabolism of Phe, and decrease Phe levels in some patients. In patients with PKU who are responsive to treatment, blood Phe levels decrease within 24 hours after administration, although maximal effect on Phe levels may take up to one month. In 8-day and 6-week pivotal trials, response to Kuvan treatment was defined as  $\geq 30\%$  decrease in blood Phe concentration from baseline. However, in some cases (e.g., patients with mild HPA, neonates or children <4 years of age, or pregnant women), a 20% reduction in blood Phe concentration may be sufficient. Kuvan is supplied as brand and generic 100 mg tablets. Both the brand and generic are also available in unit dosed powder packets for oral solution containing either 100 mg or 500 mg of the drug. The recommended starting dose of Kuvan is 10mg/kg/day taken once daily. Doses of Kuvan may be adjusted in the range of 5-20mg/kg taken once daily. Doses of Kuvan above 20mg/kg/day have not been evaluated.

## **Phenylketonuria**

PKU is an autosomal recessive disorder caused by a defect in PAH. Kuvan is a synthetic formulation of BH<sub>4</sub><sup>-</sup>. BH<sub>4</sub><sup>-</sup> has many roles in the body. One of which is to convert amino acids, such as Phe, to precursors of serotonin and dopamine via PAH. The lack of BH<sub>4</sub><sup>-</sup> could result in PKU from HPA. This buildup could lead to severe brain damage, behavioral abnormalities, mental retardation, speech impediments, etc. Kuvan works in the body to supplement the BH<sub>4</sub><sup>-</sup>, which in turn helps to convert Phe via PAH appropriately and therefore lowers the levels of Phe in the blood.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

sapropterin dihydrochloride (Kuvan<sup>®</sup>, generics)

Policy # 00303

Original Effective Date: 08/17/2011

Current Effective Date: 03/08/2021

## **FDA or Other Governmental Regulatory Approval**

### **U.S. Food and Drug Administration (FDA)**

In December 2007, the U.S. FDA granted marketing approval for Kuvan (sapropterin dihydrochloride) tablets, the first specific drug therapy approved to reduce blood Phe levels in patients with HPA due to BH4-responsive PKU.

### **Rationale/Source**

This medical policy was developed through consideration of peer-reviewed medical literature generally recognized by the relevant medical community, U.S. Food and Drug Administration approval status, nationally accepted standards of medical practice and accepted standards of medical practice in this community, Blue Cross and Blue Shield Association technology assessment program (TEC) and other non-affiliated technology evaluation centers, reference to federal regulations, other plan medical policies, and accredited national guidelines.

Kuvan was studied in four clinical trials. Study 1 was a multicenter, open-label, uncontrolled clinical trial of 489 patients with PKU, ages 8 to 48 years (mean 22 years), who had baseline blood Phe levels  $\geq 450$   $\mu\text{mol/L}$  and who were not on Phe-restricted diets. All patients received treatment with Kuvan 10mg/kg/day for 8 days. For the purposes of this study, response to Kuvan treatment was defined as a  $\geq 30\%$  decrease in blood Phe from baseline. At Day 8, 96 patients (20%) were identified as responders.

Study 2 was a multicenter, double-blind, placebo-controlled study of 88 patients with PKU who responded to Kuvan in Study 1. After a washout period from Study 1, patients were randomized equally to either Kuvan 10mg/kg/day (N=41) or placebo (N=47) for 6 weeks. Efficacy was assessed by the mean change in blood Phe level from baseline to Week 6 in the Kuvan-treated group as compared to the mean change in the placebo group. The results showed that at baseline, the mean standard deviation ( $\pm\text{SD}$ ) blood Phe level was 843 ( $\pm 300$ )  $\mu\text{mol/L}$  in the Kuvan-treated group and 888 ( $\pm 323$ )  $\mu\text{mol/L}$  in the placebo group. At Week 6, the Kuvan-treated group had a mean ( $\pm\text{SD}$ ) blood Phe level of 607 ( $\pm 377$ )  $\mu\text{mol/L}$ , and the placebo group had a mean blood Phe level of 891 ( $\pm 348$ )  $\mu\text{mol/L}$ . At Week 6, the Kuvan- and placebo-treated groups had mean changes in blood Phe level of  $-239$  and  $6$   $\mu\text{mol/L}$ , respectively (mean percent changes of  $-29\%$  ( $\pm 32$ ) and  $3\%$  ( $\pm 33$ ), respectively). The difference between the groups was statistically significant ( $p < 0.001$ ).

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

sapropterin dihydrochloride (Kuvan<sup>®</sup>, generics)

Policy # 00303

Original Effective Date: 08/17/2011

Current Effective Date: 03/08/2021

Study 3 was a multicenter, open-label, extension study in which 80 patients who responded to Kuvan treatment in Study 1 and completed Study 2 underwent 6 weeks of forced dose-titration with 3 different doses of Kuvan. Treatments consisted of 3 consecutive 2-week courses of Kuvan at doses of 5, then 20, and then 10 mg/kg/day. Blood Phe level was monitored after 2 weeks of treatment at each dose level. At baseline, mean ( $\pm$ SD) blood Phe was 844 ( $\pm$ 398)  $\mu$ mol/L. At the end of treatment with 5, 10, and 20 mg/kg/day, mean ( $\pm$ SD) blood Phe levels were 744 ( $\pm$ 384)  $\mu$ mol/L, 640 ( $\pm$ 382)  $\mu$ mol/L, and 581 ( $\pm$ 399)  $\mu$ mol/L, respectively.

Study 4 was a multicenter study of 90 children with PKU, ages 4 to 12 years, who were on Phe-restricted diets and who had blood Phe levels  $\leq$ 480 $\mu$ mol/L at screening. All patients were treated with open-label Kuvan 20 mg/kg/day for 8 days. Response to Kuvan was defined as a  $\geq$ 30% decrease in blood Phe from baseline at Day 8. At Day 8, 50 patients (56%) had a  $\geq$ 30% decrease in blood Phe.

## References

1. Kuvan [package insert]. Novato, CA: BioMarin Pharmaceuticals; March 2020.
2. Drugs at FDA. Sapropterin (Kuvan).

## Policy History

Original Effective Date: 08/17/2011

Current Effective Date: 03/08/2021

08/04/2011 Medical Policy Committee review

08/17/2011 Medical Policy Implementation Committee approval. New policy.

08/02/2012 Medical Policy Committee review

08/15/2012 Medical Policy Implementation Committee approval. No change to coverage.

02/19/2013 Coding updated

08/01/2013 Medical Policy Committee review

08/21/2013 Medical Policy Implementation Committee approval. Reworded the patient selection criteria, however there is no coverage change. Added background on phenylketonuria. Changed the rationale/source section to info from the package insert regarding the clinical trials.

08/07/2014 Medical Policy Committee review

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

sapropterin dihydrochloride (Kuvan<sup>®</sup>, generics)

Policy # 00303

Original Effective Date: 08/17/2011

Current Effective Date: 03/08/2021

08/20/2014	Medical Policy Implementation	Committee approval.	Coverage eligibility unchanged.
08/06/2015	Medical Policy Committee review		
08/19/2015	Medical Policy Implementation	Committee approval.	Coverage eligibility unchanged.
08/04/2016	Medical Policy Committee review		
08/17/2016	Medical Policy Implementation	Committee approval.	Coverage eligibility unchanged.
08/03/2017	Medical Policy Committee review		
08/23/2017	Medical Policy Implementation	Committee approval.	Coverage eligibility unchanged.
08/09/2018	Medical Policy Committee review		
08/15/2018	Medical Policy Implementation	Committee approval.	Coverage eligibility unchanged.
08/01/2019	Medical Policy Committee review		
08/14/2019	Medical Policy Implementation	Committee approval.	Coverage eligibility unchanged.
08/06/2020	Medical Policy Committee review		
08/12/2020	Medical Policy Implementation	Committee approval.	Coverage eligibility unchanged.
02/04/2021	Medical Policy Committee review		
02/10/2021	Medical Policy Implementation	Committee approval.	Title changed to reflect availability of generics. Generic added to policy with new criterion requiring trial of generic prior to brand.

Next Scheduled Review Date: 02/2022

\*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

sapropterin dihydrochloride (Kuvan<sup>®</sup>, generics)

Policy # 00303

Original Effective Date: 08/17/2011

Current Effective Date: 03/08/2021

- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
1. Consultation with the Blue Cross and Blue Shield Association technology assessment program (TEC) or other nonaffiliated technology evaluation center(s);
  2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
  3. Reference to federal regulations.

**\*\*Medically Necessary (or “Medical Necessity”)** - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. In accordance with nationally accepted standards of medical practice;
- B. Clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and
- C. Not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.

For these purposes, “nationally accepted standards of medical practice” means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

‡ Indicated trademarks are the registered trademarks of their respective owners.

**NOTICE:** If the Patient’s health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.





sapropterin dihydrochloride (Kuvan<sup>®</sup>, generics)

Policy # 00303

Original Effective Date: 08/17/2011

Current Effective Date: 03/08/2021

**NOTICE:** Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.