



Louisiana

Light Therapy for Vitiligo

Policy # 00699

Original Effective Date: 05/01/2020

Current Effective Date: 03/08/2021

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Note: Dermatologic Applications of Photodynamic Therapy is addressed separately in medical policy 00098.

Note: Light Therapy for Psoriasis is addressed separately in medical policy 00131.

Note: In-office narrow band UVB (NB-UVB) phototherapy is not subject of this medical policy, only PUVA, home NB-UVB light box and targeted phototherapy (e.g. excimer lasers and excimer lamps).

When Services Are Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Based on review of available data, the Company may consider psoralen plus ultraviolet A for the treatment of vitiligo that is not responsive to other forms of conservative therapy (eg, topical corticosteroids, coal/tar preparations, NB-UVB ultraviolet light) to be **eligible for coverage.****

When Services May Be Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Based on review of available data, the Company may consider home ultraviolet light therapy to be **eligible for coverage.****

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Light Therapy for Vitiligo

Policy # 00699

Original Effective Date: 05/01/2020

Current Effective Date: 03/08/2021

Patient Selection Criteria

Coverage eligibility may be considered for home ultraviolet light therapy when ALL of the following criteria are met:

- The device must be approved for home use by the Food and Drug Administration (FDA) and appropriate for the body surface and area treated; AND
- The member has generalized vitiligo affecting > 10% of body surface area, with documented response to in-office phototherapy, for which narrowband ultraviolet B (NB-UVB) is prescribed and indicated; AND
- Long-term maintenance UV light treatment is planned and patient is unable to attend office based treatment; AND
- The member is motivated, able to administer the treatment correctly, willing and able to keep records of treatments and attend regular follow-up visits with prescribing physician.

Note: Tanning beds are not eligible for coverage.

When Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Based on review of available data, the Company considers targeted phototherapy (e.g. excimer lasers or excimer lamps) for the treatment of vitiligo to be **investigational**.*

Policy Guidelines

The following therapies are currently being used to treat vitiligo: topical medications and narrowband ultraviolet B (NB-UVB) light therapy (in office or home). The most appropriate comparison for targeted phototherapy (e.g. excimer laser or excimer lamp) is NB-UVB, which is considered a standard treatment for active and/or widespread vitiligo based on efficacy and safety.

During psoralen plus ultraviolet A therapy, the patient needs to be assessed on a regular basis to determine the effectiveness of the therapy and the development of side effects. These evaluations are essential to ensure that the exposure dose of radiation is kept to the minimum compatible with adequate control of the disease. Therefore, psoralen plus ultraviolet A is generally not recommended for home therapy.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Light Therapy for Vitiligo

Policy # 00699

Original Effective Date: 05/01/2020

Current Effective Date: 03/08/2021

Background/Overview

Vitiligo

Vitiligo is an idiopathic skin disorder that causes depigmentation of sections of skin, most commonly on the extremities. Depigmentation occurs because melanocytes are no longer able to function properly. The cause of vitiligo is unknown; it is sometimes considered an autoimmune disease. The most common form of the disorder is nonsegmental vitiligo in which depigmentation is generalized, bilateral, symmetrical, and increases in size over time. In contrast, segmental vitiligo, also called asymmetric or focal vitiligo, covers a limited area of skin. The typical natural history of vitiligo involves stepwise progression with long periods in which the disease is static and relatively inactive, and relatively shorter periods in which areas of pigment loss increase.

Treatment

There are numerous medical and surgical treatments aimed at decreasing disease progression and/or attaining repigmentation. Topical corticosteroids, alone or in combination with topical vitamin D₃ analogues, are common first-line treatments for vitiligo. Alternative first-line therapies include topical calcineurin inhibitors, systemic steroids, and topical antioxidants. Treatment options for vitiligo recalcitrant to first-line therapy include, among others, light box therapy with narrowband ultraviolet B and psoralen plus ultraviolet A (PUVA).

Targeted phototherapy with handheld lamps or lasers is also being evaluated. Potential advantages of targeted phototherapy include the ability to use higher treatment doses and to limit exposure to surrounding tissue. Original ultraviolet B devices consisted of a Phillips TL-01 fluorescent bulb with a maximum wavelength (λ max) of 311 nm. Subsequently, xenon chloride lasers and lamps were developed as targeted ultraviolet B treatment devices; these devices generate monochromatic or very narrowband radiation with a λ max of 308 nm. Targeted phototherapy devices are directed at specific lesions or affected areas, thus limiting exposure to the surrounding normal tissues. They may, therefore, allow higher dosages compared with a lightbox, which could result in fewer treatments.

PUVA uses a psoralen derivative in conjunction with long-wavelength ultraviolet A (UVA) light (sunlight or artificial) for photochemotherapy of skin conditions. Psoralens are tricyclic furocoumarins that occur in certain plants and can also be synthesized. They are available in oral and topical forms. Oral PUVA is generally given 1.5 hours before exposure to UVA radiation.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Light Therapy for Vitiligo

Policy # 00699

Original Effective Date: 05/01/2020

Current Effective Date: 03/08/2021

Topical PUVA therapy refers to the direct application of psoralen to the skin with subsequent exposure to UVA light. With topical PUVA, UVA exposure is generally administered within 30 minutes of psoralen application.

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

In 2001, XTRAC^{TM‡} (PhotoMedex), a xenon chloride (XeCl) excimer laser, was cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process for the treatment of skin conditions such as vitiligo. The 510(k) clearance has subsequently been obtained for a number of targeted UVB lamps and lasers, including newer versions of the XTRAC system including the XTRAC Ultra^{TM‡}, the VTRAC^{TM‡} lamp (PhotoMedex), the BCclear^{TM‡} lamp (Lumenis), the 308 excimer lamp phototherapy system (Quantel Medical), MultiClear Multiwavelength Targeted Phototherapy System, Psoria-Light^{TM‡}, and the Excilite^{TM‡} and Excilite μ ^{TM‡} XeCl lamps. The intended use of all of these devices includes vitiligo among other dermatologic indications. Some light-emitting devices are handheld. FDA product code: GEX.

The oral psoralen product, Oxsoresalen-Ultra (methoxsalen soft gelatin capsules), has been approved by the FDA and is made by Bausch Health; a generic product is also available from various manufacturers. Topical psoralen products (Oxsoresalen; Valeant Pharmaceuticals) and methoxsalen hard gelatin capsules have been discontinued.

Rationale/Source

Vitiligo is an idiopathic skin disorder that causes depigmentation of sections of skin, most commonly on the extremities. Topical corticosteroids, alone or in combination with topical vitamin D₃ analogues, are common first-line treatments for vitiligo. Alternative first-line therapies include topical calcineurin inhibitors, systemic steroids, and topical antioxidants. Treatment options for vitiligo recalcitrant to first-line therapy include, among others, ultraviolet B, light box therapy, and psoralen plus ultraviolet A (PUVA). Targeted phototherapy is also being evaluated.

For individuals who have vitiligo who receive targeted phototherapy, the evidence includes systematic reviews of randomized controlled trials (RCTs), 2 individual RCTs, and 2 retrospective studies. Relevant outcomes are a change in disease status, quality of life, and treatment-related morbidity. Individual studies tend to have small sample sizes, and few were designed to isolate the

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Light Therapy for Vitiligo

Policy # 00699

Original Effective Date: 05/01/2020

Current Effective Date: 03/08/2021

effect of laser therapy. Two meta-analyses were attempted; however, results from a meta-analysis could not be verified because the selected studies were not available in English, and 1 estimate was imprecise due to the small number of studies and participants. RCTs have shown targeted phototherapy to be associated with statistically significant improvements in Vitiligo Area Scoring index scores and/or repigmentation compared to alternate treatment options. However, 1 of the RCTs only showed marginal differences between groups in these outcomes limiting clinical significance, and the second compared phototherapy to oral vitamin E, which is not an optimal comparator. Overall, there is a lack of clinical trial evidence that compares this technique with more conservative treatments or no treatment/placebo. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have vitiligo who have not responded to conservative therapy who receive PUVA (photochemotherapy), the evidence includes systematic reviews and RCTs. Relevant outcomes are a change in disease status, quality of life, and treatment-related morbidity. There is some evidence from randomized studies, mainly those published before 1985, that PUVA is more effective than a placebo for treating vitiligo. When compared with narrowband ultraviolet B in meta-analyses, results have shown that patients receiving narrowband ultraviolet B experienced higher rates of repigmentation than patients receiving PUVA, though the differences were not statistically significant. Based on the available evidence and clinical guidelines, PUVA may be considered in patients with vitiligo who have not responded adequately to conservative therapy. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

Supplemental Information

Practice Guidelines and Position Statements

British Association of Dermatologists et al.

In 2008, guidelines on the diagnosis and management of vitiligo were published by a collaboration of several U.K. organizations, including the British Association of Dermatologists, the Royal College of Physicians of London, and the Cochrane Skin Group. As of October 2020, an update for this guideline is currently in progress. The guidelines included the following statements (Table 1).

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Light Therapy for Vitiligo

Policy # 00699

Original Effective Date: 05/01/2020

Current Effective Date: 03/08/2021

Table 1. Guidelines on the Diagnosis and Management of Vitiligo

Recommendation	GOE	LOE
PUVA therapy should be considered for the treatment of vitiligo only in adults who cannot be adequately managed with more conservative treatments. PUVA is not recommended in children.	D	4
If phototherapy is to be used for treating nonsegmental vitiligo, NB-UVB should usually be used in preference to oral PUVA.	A	1+
A trial of PUVA therapy should be considered only for adults with widespread vitiligo, or localized vitiligo associated with a significant impact on a patient's quality of life. Ideally, this treatment should be reserved for patients with darker skin types.	D	3
Before starting PUVA treatment, patients should be made aware that there is no evidence that this treatment alters the natural history of vitiligo. They should also be made aware that not all patients respond, and that some sites on the body, such as the hands and feet, respond poorly in all patients. They should also be informed of the limit to the number of treatments due to possible adverse effects.	D	3

GOE: grade of evidence; LOE: level of evidence; NB-UVB: narrowband ultraviolet B; PUVA: psoralens with ultraviolet A.

European Dermatology Forum

In 2013, the European Dermatology Forum published consensus guidelines on the management of vitiligo. The guidelines stated that oral psoralens with ultraviolet A are commonly used in adults with generalized vitiligo as a second-line treatment. The guidelines also stated that targeted phototherapy is indicated for localized vitiligo, particularly small lesions of recent onset and childhood vitiligo, to avoid adverse effects due to total body irradiation and when total body irradiation is contraindicated. The guidelines were based on expert opinion, not a systematic review of the literature.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Light Therapy for Vitiligo

Policy # 00699

Original Effective Date: 05/01/2020

Current Effective Date: 03/08/2021

Vitiligo Working Group

The Vitiligo Working Group is supported by the National Institute of Arthritis and Musculoskeletal and Skin Diseases, part of the National Institutes of Health. In 2017, the group published guidelines on current and emerging treatments for vitiligo. The Working Group indicated that psoralens with ultraviolet A have largely been replaced by narrowband ultraviolet B, but that “PUVA may be considered in patients with darker Fitzpatrick skin phototypes or those with treatment-resistant vitiligo (level I evidence).” The Working Group also stated that “Targeted phototherapy (excimer lasers and excimer lamps) can be considered when <10% of body surface area is affected (level II evidence).”

U.S. Preventive Services Task Force Recommendation

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

Ongoing and Unpublished Clinical Trials

A search of [ClinicalTrials.gov](https://clinicaltrials.gov) in October 2020 did not identify any ongoing or unpublished trials that may influence this review.

References

1. Blue Cross and Blue Shield Association, Medical Policy Reference Manual, “Targeted Phototherapy and Psoralen with Ultraviolet A for Vitiligo”, 2.01.86, January 2021.
2. Blue Cross and Blue Shield of Louisiana, “Light Therapy for Psoriasis”, 00131, December 14, 2020.
3. Lopes C, Trevisani VF, Melnik T. Efficacy and Safety of 308-nm Monochromatic Excimer Lamp Versus Other Phototherapy Devices for Vitiligo: A Systematic Review with Meta-Analysis. *Am J Clin Dermatol*. Feb 2016; 17(1): 23-32. PMID 26520641
4. Whitton ME, Pinart M, Batchelor J, et al. Interventions for vitiligo. *Cochrane Database Syst Rev*. Feb 24 2015; (2): CD003263. PMID 25710794
5. Sun Y, Wu Y, Xiao B, et al. Treatment of 308-nm excimer laser on vitiligo: A systemic review of randomized controlled trials. *J Dermatolog Treat*. 2015; 26(4): 347-53. PMID 25428573

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Light Therapy for Vitiligo

Policy # 00699

Original Effective Date: 05/01/2020

Current Effective Date: 03/08/2021

6. Yang YS, Cho HR, Ryou JH, et al. Clinical study of repigmentation patterns with either narrow-band ultraviolet B (NBUVB) or 308 nm excimer laser treatment in Korean vitiligo patients. *Int J Dermatol*. Mar 2010; 49(3): 317-23. PMID 20465673
7. Poolsuwan P, Churee C, Pattamadilok B. Comparative efficacy between localized 308-nm excimer light and targeted 311-nm narrowband ultraviolet B phototherapy in vitiligo: A randomized, single-blind comparison study. *Photodermatol Photoimmunol Photomed*. Oct 12 2020. PMID 33047405
8. Nistico S, Chiricozzi A, Saraceno R, et al. Vitiligo treatment with monochromatic excimer light and tacrolimus: results of an open randomized controlled study. *Photomed Laser Surg*. Jan 2012; 30(1): 26-30. PMID 22054204
9. Fa Y, Lin Y, Chi XJ, et al. Treatment of vitiligo with 308-nm excimer laser: our experience from a 2-year follow-up of 979 Chinese patients. *J Eur Acad Dermatol Venereol*. Feb 2017; 31(2): 337-340. PMID 27538097
10. Dong DK, Pan ZY, Zhang J, et al. Efficacy and Safety of Targeted High-Intensity Medium-Band (304-312 nm) Ultraviolet B Light in Pediatric Vitiligo. *Pediatr Dermatol*. May 2017; 34(3): 266-270. PMID 28318054
11. Bae JM, Jung HM, Hong BY, et al. Phototherapy for Vitiligo: A Systematic Review and Meta-analysis. *JAMA Dermatol*. Jul 01 2017; 153(7): 666-674. PMID 28355423
12. Njoo MD, Spuls PI, Bos JD, et al. Nonsurgical repigmentation therapies in vitiligo. Meta-analysis of the literature. *Arch Dermatol*. Dec 1998; 134(12): 1532-40. PMID 9875190
13. Yones SS, Palmer RA, Garibaldinos TM, et al. Randomized double-blind trial of treatment of vitiligo: efficacy of psoralen-UV-A therapy vs Narrowband-UV-B therapy. *Arch Dermatol*. May 2007; 143(5): 578-84. PMID 17519217
14. Gawkrödger DJ, Ormerod AD, Shaw L, et al. Guideline for the diagnosis and management of vitiligo. *Br J Dermatol*. Nov 2008; 159(5): 1051-76. PMID 19036036
15. Taieb A, Alomar A, Bohm M, et al. Guidelines for the management of vitiligo: the European Dermatology Forum consensus. *Br J Dermatol*. Jan 2013; 168(1): 5-19. PMID 22860621
16. Rodrigues M, Ezzedine K, Hamzavi I, et al. Current and emerging treatments for vitiligo. *J Am Acad Dermatol*. Jul 2017; 77(1): 17-29. PMID 28619557

Policy History

Original Effective Date: 05/01/2020

Current Effective Date: 03/08/2021

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Light Therapy for Vitiligo

Policy # 00699

Original Effective Date: 05/01/2020

Current Effective Date: 03/08/2021

02/06/2020 Medical Policy Committee review

02/12/2020 Medical Policy Implementation Committee approval. New policy.

02/04/2021 Medical Policy Committee review

02/10/2021 Medical Policy Implementation Committee approval. Removed “box” from home ultraviolet box therapy in the coverage section. Coverage eligibility unchanged.

Next Scheduled Review Date: 02/2022

Coding

The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)‡, copyright 2020 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	96912, 96913, 96920, 96921, 96922, 96999

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Light Therapy for Vitiligo

Policy # 00699

Original Effective Date: 05/01/2020

Current Effective Date: 03/08/2021

HCPCS	E0691, E0692, E0693, E0694
ICD-10 Diagnosis	H02.73, L80

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
 - 1. Consultation with the Blue Cross and Blue Shield Association technology assessment program (TEC) or other nonaffiliated technology evaluation center(s);
 - 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
 - 3. Reference to federal regulations.

**Medically Necessary (or “Medical Necessity”) - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. In accordance with nationally accepted standards of medical practice;
- B. Clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and
- C. Not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Light Therapy for Vitiligo

Policy # 00699

Original Effective Date: 05/01/2020

Current Effective Date: 03/08/2021

For these purposes, “nationally accepted standards of medical practice” means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

‡ Indicated trademarks are the registered trademarks of their respective owners.

NOTICE: If the Patient’s health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

©2021 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.