



Louisiana

Bioimpedance Devices for Detection and Management of Lymphedema

Policy # 00780

Original Effective Date: 06/01/2022

Current Effective Date: 06/01/2022

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Note: Surgical Treatments for Breast Cancer-Related Lymphedema is addressed separately in medical policy 00635.

Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Based on review of available data, the Company considers devices using bioimpedance (bioelectrical impedance spectroscopy) for use in the diagnosis, surveillance, or treatment of patients with lymphedema, including use in subclinical secondary lymphedema, to be **investigational**.*

Background/Overview

Lymphedema

Lymphedema is an accumulation of fluid due to disruption of lymphatic drainage. Lymphedema can be caused by congenital or inherited abnormalities in the lymphatic system (primary lymphedema) but is most often caused by acquired damage to the lymphatic system (secondary lymphedema). Breast cancer treatment is one of the most common causes of secondary lymphedema. Both the surgical removal of lymph nodes and radiotherapy are associated with development lymphedema in patients with breast cancer. In a systematic review of 72 studies (N=29612 women), DiSipio et al (2013) reported that approximately 1 in 5 women who survive breast cancer will develop arm lymphedema. Risk factors for development of lymphedema that had a strong level of evidence were extensive surgery (ie, axillary-lymph-node dissection, greater number of lymph nodes dissected, mastectomy) and being overweight or obese.

Diagnosis and Staging

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Bioimpedance Devices for Detection and Management of Lymphedema

Policy # 00780

Original Effective Date: 06/01/2022

Current Effective Date: 06/01/2022

A diagnosis of secondary lymphedema is based on history (e.g., cancer treatment, trauma) and physical examination (localized, progressive edema and asymmetric limb measurements) when other causes of edema can be excluded. Imaging, such as MRI, computed tomography, ultrasound, or lymphoscintigraphy, may be used to differentiate lymphedema from other causes of edema in diagnostically challenging cases.

Table 1 lists International Society of Lymphology guidance for staging lymphedema based on "softness" or "firmness" of the limb and the changes with an elevation of the limb.

Table 1. Recommendations for Staging Lymphedema

Stage	Description
Stage 0 (subclinical)	Swelling is not evident and most patients are asymptomatic despite impaired lymphatic transport
Stage I (mild)	Accumulation of fluid that subsides (usually within 24 hours) with limb elevation; soft edema that may pit, without evidence of dermal fibrosis
Stage II (moderate)	Does not resolve with limb elevation alone; limb may no longer pit on examination
Stage III (severe)	Lymphostatic elephantiasis; pitting can be absent; skin has trophic changes

Management and Treatment

Lymphedema is treated using elevation, compression, and exercise. Conservative therapy may consist of several features depending on the severity of the lymphedema. Patients are educated on the importance of self-care including hygiene practices to prevent infection, maintaining ideal body weight through diet and exercise, and limb elevation. Compression therapy consists of repeatedly applying padding and bandages or compression garments. Manual lymphatic drainage is a light pressure massage performed by trained physical therapists or by patients designed to move fluid from obstructed areas into functioning lymph vessels and lymph nodes. Complete decongestive therapy is a multiphase treatment program involving all of the previously mentioned conservative treatment components at different intensities. Pneumatic compression pumps may also be considered as an adjunct to conservative therapy or as an alternative to self-manual lymphatic drainage in patients who have difficulty performing self-manual lymphatic drainage. In patients with more

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Bioimpedance Devices for Detection and Management of Lymphedema

Policy # 00780

Original Effective Date: 06/01/2022

Current Effective Date: 06/01/2022

advanced lymphedema after fat deposition and tissue fibrosis has occurred, palliative surgery using reductive techniques such as liposuction may be performed.

Bioimpedance Spectroscopy

Bioimpedance spectroscopy is based on the theory that the level of opposition to the flow of electric current (impedance) through the body is inversely proportional to the volume of fluid in the tissue. In lymphedema, with the accumulation of excess interstitial fluid, tissue impedance decreases.

Bioimpedance has been proposed as a diagnostic test for this condition. In usual care, lymphedema is recognized clinically or via limb measurements. However, management via bioelectrical impedance spectroscopy has been proposed as a way to implement early treatment of subclinical lymphedema to potentially reduce its severity.

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

Devices that have been cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process to aid in the assessment of lymphedema are summarized in Table 2.

Table 2. FDA Cleared Bioimpedance Spectroscopy Devices for Lymphedema

Year	Device	Manufacturer	510(k) Number	Indication
2018	SOZO	ImpediMed (Carlsbad, CA)	K180126	For adults at risk of lymphedema. Supports the measurement of extracellular fluid volume differences between the limbs and is presented to the clinician on an L-Dex scale as an aid to their clinical assessment of lymphedema.

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Bioimpedance Devices for Detection and Management of Lymphedema

Policy # 00780

Original Effective Date: 06/01/2022

Current Effective Date: 06/01/2022

				The device is only indicated for patients who will have or who have had lymph nodes, from the axillary and/or pelvic regions, either removed, damaged or irradiated.
2015	MoistureMeterD	Delfin Technologies (Stamford, CT)	K143310	Supports local assessment of tissue water differences between affected and contralateral non-affected arm tissues to aid in forming a clinical judgment of unilateral lymphedema in women. The device is not intended to make diagnosis or predict arm lymphedema.
2007	ImpediMed L-Dex™‡ U400	ImpediMed (Carlsbad, CA)	K050415	Supports the measurement of extracellular fluid volume differences between the arms to aid in the clinical assessment of unilateral lymphedema of the arm in women. This device is not intended to diagnose or predict

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Bioimpedance Devices for Detection and Management of Lymphedema

Policy # 00780

Original Effective Date: 06/01/2022

Current Effective Date: 06/01/2022

				lymphedema of an extremity.
--	--	--	--	-----------------------------

FDA product code: OBH.

Rationale/Source

This medical policy was developed through consideration of peer-reviewed medical literature generally recognized by the relevant medical community, U.S. Food and Drug Administration approval status, nationally accepted standards of medical practice and accepted standards of medical practice in this community, technology evaluation centers, reference to federal regulations, other plan medical policies, and accredited national guidelines.

Secondary lymphedema may develop following treatment for breast cancer. Bioimpedance, which uses resistance to electrical current to compare the composition of fluid compartments, could be used as a tool to diagnose lymphedema.

For individuals who have known or suspected lymphedema who receive bioimpedance spectroscopy, the evidence includes a systematic review, 1 RCT, 1 prospective comparative observational study, and multiple uncontrolled observational studies. Relevant outcomes are test validity, symptoms, and quality of life. Diagnostic accuracy studies have found a poor correlation between bioimpedance analysis and the reference standard (volume displacement or circumferential measurement). Interim results from an ongoing RCT comparing bioimpedance with standard tape measure following treatment for breast cancer have been published. Overall, 109 of 508 (21.5%) patients received early treatment due to reaching a pre-determined threshold to trigger an intervention. A total of 12 triggering patients progressed to clinical lymphedema (2 in the bioimpedance group [4.9%] and 10 in the tape measure group [14.7%]; P=0.130). The RCT was limited by its open-label design and lack of reporting of important health outcomes. The single prospective comparative study found a significantly lower rate of clinical lymphedema in patients managed with bioimpedance devices but had several limitations, including nonrandomized design, lack of blinding, lack of complete data on a substantial proportion of enrolled patients, and lack of a systematic method for diagnosing lymphedema in the control group. Retrospective studies suggested that postoperative bioimpedance monitoring is feasible but provide limited information about its efficacy. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Bioimpedance Devices for Detection and Management of Lymphedema

Policy # 00780

Original Effective Date: 06/01/2022

Current Effective Date: 06/01/2022

Supplemental Information

Clinical Input From Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

In response to requests, input was received from 2 specialty societies and 2 academic medical centers while this policy was under review in 2011. Three of 4 reviewers agreed that bioimpedance devices are considered investigational for diagnosis, surveillance, and treatment of patients with lymphedema. The fourth reviewer, from an academic medical center, considered the use of the technology a reasonable alternative, especially in situations in which minor lymphedema can have a large impact on a patient. One specialty society supported further research into the effectiveness of this technology and recommended reimbursement in the context of relevant clinical trials.

Practice Guidelines and Position Statements

National Comprehensive Cancer Network Clinical Practice Guidelines on Survivorship (v.2.2020) and Breast Cancer (v.6.2020) recommend education, monitoring, and referral for lymphedema management as needed. Neither guideline mentions bioimpedance.

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 3.

Table 3. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
---------	------------	--------------------	-----------------

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Bioimpedance Devices for Detection and Management of Lymphedema

Policy # 00780

Original Effective Date: 06/01/2022

Current Effective Date: 06/01/2022

<i>Ongoing</i>			
NCT03978754	Assessment of Breast Cancer-Related Arm Lymphedema—Comparison of Traditional Measurement Methods and Indocyanine Green (ICG) Lymphography	300	Jan 2022 (recruiting as of Jun 2019)
NCT02743858	A Prospective Surveillance Program for Assessment and Treatment of Breast Cancer-Related Lymphedema After Axillary Lymph Node Dissection	850	Apr 2021 (recruiting as of Feb 2020)
NCT01544335	Evaluation of the Validity of BIS as a Tool for Quantification of Lymphedema Through Comparison With Perometry and Self-Report	270	Dec 2020
<i>Unpublished</i>			

BIS: bioimpedance spectroscopy; NCT: national clinical trial.

References

1. DiSipio T, Rye S, Newman B, et al. Incidence of unilateral arm lymphoedema after breast cancer: a systematic review and meta-analysis. *Lancet Oncol.* May 2013; 14(6): 500-15. PMID 23540561
2. International Society of Lymphology Executive Committee. The Diagnosis and Treatment of Peripheral Lymphedema: 2016 Consensus Document of the International Society of Lymphology. 2016; <https://journals.uair.arizona.edu/index.php/lymph/article/view/20106>.
3. Pusic AL, Cemal Y, Albornoz C, et al. Quality of life among breast cancer patients with lymphedema: a systematic review of patient-reported outcome instruments and outcomes. *J Cancer Surviv.* Mar 2013; 7(1): 83-92. PMID 23212603
4. Oremus M, Walker K, Dayes I, et al. Technology Assessment: Diagnosis and treatment of secondary lymphedema. Rockville, MD: Agency for Healthcare Research and Quality; 2010.
5. Cornish BH, Chapman M, Hirst C, et al. Early diagnosis of lymphedema using multiple frequency bioimpedance. *Lymphology.* Mar 2001; 34(1): 2-11. PMID 11307661

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Bioimpedance Devices for Detection and Management of Lymphedema

Policy # 00780

Original Effective Date: 06/01/2022

Current Effective Date: 06/01/2022

6. Hayes S, Janda M, Cornish B, et al. Lymphedema secondary to breast cancer: how choice of measure influences diagnosis, prevalence, and identifiable risk factors. *Lymphology*. Mar 2008; 41(1): 18-28. PMID 18581955
7. Barrio AV, Eaton A, Frazier TG. A Prospective Validation Study of Bioimpedance with Volume Displacement in Early-Stage Breast Cancer Patients at Risk for Lymphedema. *Ann Surg Oncol*. Dec 2015; 22 Suppl 3: S370-5. PMID 26085222
8. Blaney JM, McCollum G, Lorimer J, et al. Prospective surveillance of breast cancer-related lymphoedema in the first-year post-surgery: feasibility and comparison of screening measures. *Support Care Cancer*. Jun 2015; 23(6): 1549-59. PMID 25398360
9. Ridner SH, Dietrich MS, Cowher MS, et al. A Randomized Trial Evaluating Bioimpedance Spectroscopy Versus Tape Measurement for the Prevention of Lymphedema Following Treatment for Breast Cancer: Interim Analysis. *Ann Surg Oncol*. Oct 2019; 26(10): 3250-3259. PMID 31054038
10. Soran A, Ozmen T, McGuire KP, et al. The importance of detection of subclinical lymphedema for the prevention of breast cancer-related clinical lymphedema after axillary lymph node dissection; a prospective observational study. *Lymphat Res Biol*. Dec 2014; 12(4): 289-94. PMID 25495384
11. Laidley A, Anglin B. The Impact of L-Dex((R)) Measurements in Assessing Breast Cancer-Related Lymphedema as Part of Routine Clinical Practice. *Front Oncol*. 2016; 6: 192. PMID 27656420
12. Koelmeyer LA, Borotkanics RJ, Alcorso J, et al. Early surveillance is associated with less incidence and severity of breast cancer-related lymphedema compared with a traditional referral model of care. *Cancer*. Mar 15 2019; 125(6): 854-862. PMID 30521080
13. Kilgore LJ, Korentager SS, Hangge AN, et al. Reducing Breast Cancer-Related Lymphedema (BCRL) Through Prospective Surveillance Monitoring Using Bioimpedance Spectroscopy (BIS) and Patient Directed Self-Interventions. *Ann Surg Oncol*. Oct 2018; 25(10): 2948-2952. PMID 29987599
14. Whitworth PW, Cooper A. Reducing chronic breast cancer-related lymphedema utilizing a program of prospective surveillance with bioimpedance spectroscopy. *Breast J*. Jan 2018; 24(1): 62-65. PMID 29063664
15. Erdogan Iyigun Z, Selamoglu D, Alco G, et al. Bioelectrical impedance for detecting and monitoring lymphedema in patients with breast cancer. Preliminary results of the florence nightingale breast study group. *Lymphat Res Biol*. Mar 2015; 13(1): 40-5. PMID 25526543

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Bioimpedance Devices for Detection and Management of Lymphedema

Policy # 00780

Original Effective Date: 06/01/2022

Current Effective Date: 06/01/2022

16. Shah C, Vicini F, Beitsch P, et al. The use of bioimpedance spectroscopy to monitor therapeutic intervention in patients treated for breast cancer related lymphedema. *Lymphology*. Dec 2013; 46(4): 184-92. PMID 25141461
17. Lim SM, Han Y, Kim SI, et al. Utilization of bioelectrical impedance analysis for detection of lymphedema in breast Cancer survivors: a prospective cross sectional study. *BMC Cancer*. Jul 08 2019; 19(1): 669. PMID 31286884
18. Kaufman DI, Shah C, Vicini FA, et al. Utilization of bioimpedance spectroscopy in the prevention of chronic breast cancer-related lymphedema. *Breast Cancer Res Treat*. Dec 2017; 166(3): 809-815. PMID 28831632
19. Whitworth PW, Shah C, Vicini F, et al. Preventing Breast Cancer-Related Lymphedema in High-Risk Patients: The Impact of a Structured Surveillance Protocol Using Bioimpedance Spectroscopy. *Front Oncol*. 2018; 8: 197. PMID 29946531
20. National Comprehensive Cancer Network. Clinical Practice Guidelines in Oncology: Survivorship. Version 2.2020. https://www.nccn.org/professionals/physician_gls/pdf/survivorship.pdf.
21. National Comprehensive Cancer Network. Clinical Practice Guidelines in Oncology: Breast Cancer. Version 6.2020. https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf.

Policy History

Original Effective Date: 06/01/2022

Current Effective Date: 06/01/2022

03/03/2022 Medical Policy Committee review

03/09/2022 Medical Policy Implementation Committee approval. New policy.

Next Scheduled Review Date: 03/2023

Coding

The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)‡, copyright 2021 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Bioimpedance Devices for Detection and Management of Lymphedema

Policy # 00780

Original Effective Date: 06/01/2022

Current Effective Date: 06/01/2022

The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	93702
HCPCS	No codes
ICD-10 Diagnosis	I97.2, Z90.10-Z90.13

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Bioimpedance Devices for Detection and Management of Lymphedema

Policy # 00780

Original Effective Date: 06/01/2022

Current Effective Date: 06/01/2022

diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:

1. Consultation with technology evaluation center(s);
2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
3. Reference to federal regulations.

‡ Indicated trademarks are the registered trademarks of their respective owners.

NOTICE: If the Patient's health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

©2022 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.