



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Note: Microarray-based Gene Expression Analysis for Prostate Cancer Management is addressed separately in medical policy 00403.

Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Based on review of available data, the Company considers the following genetic and protein biomarkers for the diagnosis of prostate cancer to be **investigational***:

- Kallikrein markers (eg, 4Kscore^{™†} Test)
- Metabolomic profiles (eg, Prostarix^{™†})
- PCA3 testing
- TMPRSS fusion genes
- Candidate gene panels
- Mitochondrial DNA variant testing (eg, Prostate Core Mitomics Test^{™†})
- Gene hypermethylation testing (eg, ConfirmMDx^{®†})
- Prostate Health Index (phi).

Based on review of available data, the Company considers single nucleotide variant testing for cancer risk assessment of prostate cancer to be **investigational.***

Policy Guidelines

GENETICS NOMENCLATURE UPDATE

The Human Genome Variation Society nomenclature is used to report information on variants found in DNA and serves as an international standard in DNA diagnostics. It is being implemented for genetic testing medical evidence review updates starting in 2017 (see Table PG1). The Society's nomenclature is recommended by the Human Variome Project, the HUMAN Genome Organization, and by the Human Genome Variation Society itself.

The American College of Medical Genetics and Genomics and the Association for Molecular Pathology standards and guidelines for interpretation of sequence variants represent expert opinion from both organizations, in addition to the College of American Pathologists. These recommendations primarily apply to genetic tests used in clinical laboratories, including genotyping, single genes, panels, exomes, and genomes. Table PG2 shows the recommended standard terminology—"pathogenic," "likely pathogenic,"

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

“uncertain significance,” “likely benign,” and “benign”—to describe variants identified that cause Mendelian disorders.

Table PG1. Nomenclature to Report on Variants Found in DNA

Previous	Updated	Definition
Mutation	Disease-associated variant	Disease-associated change in the DNA sequence
	Variant	Change in the DNA sequence
	Familial variant	Disease-associated variant identified in a proband for use in subsequent targeted genetic testing in first-degree relatives

Table PG2. ACMG-AMP Standards and Guidelines for Variant Classification

Variant Classification	Definition
Pathogenic	Disease-causing change in the DNA sequence
Likely pathogenic	Likely disease-causing change in the DNA sequence
Variant of uncertain significance	Change in DNA sequence with uncertain effects on disease
Likely benign	Likely benign change in the DNA sequence
Benign	Benign change in the DNA sequence

American College of Medical Genetics and Genomics; AMP: Association for Molecular Pathology.

GENETIC COUNSELING

Experts recommend formal genetic counseling for patients who are at risk for inherited disorders and who wish to undergo genetic testing. Interpreting the results of genetic tests and understanding risk factors can be difficult for some patients; genetic counseling helps individuals understand the impact of genetic testing, including the possible effects the test results could have on the individual or their family members. It should be noted that genetic counseling may alter the utilization of genetic testing substantially and may reduce inappropriate testing; further, genetic counseling should be performed by an individual with experience and expertise in genetic medicine and genetic testing methods.

Background/Overview

PROSTATE CANCER

Prostate cancer is the second most common cancer in men, with a predicted 161,360 incidence cases and 26,730 deaths expected in the United States in 2017.

Prostate cancer is a complex, heterogeneous disease, ranging from microscopic tumors unlikely to be life-threatening to aggressive tumors that can metastasize, leading to morbidity or death. Early localized disease can usually be treated with surgery and radiotherapy, although active surveillance may be adopted in men whose cancer is unlikely to cause major health problems during their lifespan or for whom the treatment might be dangerous. In patients with inoperable or metastatic disease, treatment consists of hormonal therapy and possibly chemotherapy. The lifetime risk of being diagnosed with prostate cancer for men in the United States is approximately 16%, while the risk of dying of prostate cancer is 3%. African American men have the highest prostate cancer risk in the United States; the incidence of prostate cancer

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

is about 60% higher and the mortality rate is more than 2 to 3 times greater than that of white men. Autopsy results have suggested that about 30% of men age 55 and 60% of men age 80 who die of other causes have incidental prostate cancer, indicating that many cases of cancer are unlikely to pose a threat during a man's life expectancy.

Grading

The most widely used grading scheme for prostate cancer is the Gleason system. It is an architectural grading system ranging from 1 (well differentiated) to 5 (poorly differentiated); the score is the sum of the primary and secondary patterns. A Gleason score of 6 is low-grade prostate cancer that usually grows slowly; 7 is an intermediate grade; 8 to 10 is high-grade cancer that grows more quickly. Ten-year survival rates stratified by Gleason score have been estimated from the Surveillance, Epidemiology, and End Results registry to be about 98% for scores 2 through 6, 92% for a score of 7 with primary pattern 3 and secondary pattern 4 (3+4), 77% for a score of 7 (4+3), and 70% for scores between 8 and 10.

Numerous genetic alterations associated with development or progression of prostate cancer have been described, with the potential for the use of these molecular markers to improve the selection process of men who should undergo prostate biopsy or rebiopsy after an initial negative biopsy.

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratory-developed tests must meet the general regulatory standards of the Clinical Laboratory Improvement Amendments. Laboratories that offer laboratory-developed tests must be licensed under the Clinical Laboratory Improvement Amendments for high-complexity testing. The following laboratories are certified under the Clinical Laboratory Improvement Amendments: BioReference Laboratories and GenPath Diagnostics (subsidiaries of OPKO Health; 4Kscore[®])[‡], ARUP Laboratories, Mayo Medical Laboratories, LabCorp, BioVantra, others (PCA3 assay), Clinical Research Laboratory (Prostate Core Mitomic Test[™])[‡], MDx Health (ConfirMDx), and Innovative Diagnostics (phi[™])[‡]. To date, the U.S. FDA has chosen not to require any regulatory review of this test.

In February 2012, the Progenesa[®] PCA3 Assay (Gen-Probe; now Hologic, Marlborough, MA) was approved by FDA through the premarket approval process. According to the company's press release, this assay is "indicated for use in conjunction with other patient information to aid in the decision for repeat biopsy in men 50 years of age or older who have had 1 or more previous negative prostate biopsies and for whom a repeat biopsy would be recommended by a urologist based on the current standard of care, before consideration of Progenesa PCA3 assay results." FDA product code: OYM.

In June 2012, proPSA, a blood test used to calculate the Prostate Health Index (phi; Beckman Coulter, Brea, CA) was approved by FDA through the premarket approval process. The phi test is indicated as an aid to distinguish prostate cancer from a benign prostatic condition in men ages 50 and older with prostate-

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

specific antigen levels of 4 to 10 ng/mL and with digital rectal exam findings that are not suspicious. According to the manufacturer, the test reduces the number of prostate biopsies. FDA product code: OYA.

Centers for Medicare and Medicaid Services (CMS)

There is no national coverage determination (NCD). In the absence of an NCD, coverage decisions are left to the discretion of local Medicare carriers.

Rationale/Source

Genetic and protein biomarker tests are best evaluated within the framework of a diagnostic or prognostic test because such frameworks provide diagnostic and prognostic information that assists in clinical management decisions. Assessment of a diagnostic or prognostic tool typically focuses on 3 categories of evidence: (1) analytic validity (ability of the test to accurately and reliably measure the marker of interest); (2) clinical validity (ie, statistically significant association between the test result and health outcomes); and (3) clinical utility (ie, demonstration that use of the diagnostic or prognostic information clinically can improve the net health outcome compared with patient management without use of the tool). Because these tests are used as an adjunct to the usual diagnostic workup, it is important to evaluate whether the tests provide incremental information above the standard workup to determine whether the tests have utility in clinical practice.

This review evaluates evidence for genetic and protein biomarkers for the purpose of guiding decision making about biopsy or rebiopsy (see Appendix Table 1 for genetic testing categories).

BIOMARKER TESTING FOR SELECTION OF MEN FOR INITIAL PROSTATE BIOPSY

Clinical Context and Test Purpose

The purpose of genetic and protein biomarker testing for prostate cancer is to inform the selection of men who should undergo initial biopsy. Conventional decision-making tools for identifying men for prostate biopsy include digital rectal exam (DRE), serum prostate-specific antigen (PSA), and patient risk factors such as age, race, and family history of prostate cancer.

DRE has relatively low interrater agreement among urologists, with estimated sensitivity, specificity, and positive predictive value (PPV) for diagnosis of prostate cancer of 59%, 94%, and 28%, respectively. DRE might have a higher PPV in the setting of elevated PSA.

The risk of prostate cancer increases with increasing PSA levels; an estimated 15% of men with a PSA level of 4 ng/mL or less and normal DRE, 30% to 35% of men with a PSA level between 4 ng/mL and 10 ng/mL, and more than 67% of men with a PSA level greater than 10 ng/mL will have biopsy-detectable prostate cancer. Use of PSA levels in screening has improved detection of prostate cancer. The European Randomized Study of Screening for Prostate Cancer (ERSPC) trial and Göteborg Randomised Prostate Cancer Screening Trial trials demonstrated that biennial PSA screening reduces the risk of being diagnosed with metastatic prostate cancer.

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

However, elevated PSA levels are not specific to prostate cancer; levels can be elevated due to infection, inflammation, trauma, or ejaculation. In addition, there are no clear cutoffs for cancer positivity with PSA. Using a common PSA level cutoff of 4.0 ng/mL, the American Cancer Society (2010) systematically reviewed the literature and calculated pooled estimates of elevated PSA sensitivity of 21% for detecting any prostate cancer and 5% for detecting high-grade cancers with an estimated specificity of 91%.

PSA screening in the general population is controversial. The U.S. Preventive Services Task Force recommended against PSA-based screening (D recommendation) in 2012; the recommendations are currently being updated, and the draft released in 2017 advises individualized decision making about screening for prostate cancer after discussion with a clinician for men ages 55 to 69 (C recommendation) and recommends against PSA-based screening in men 70 and older (D recommendation). Guidelines published by the American Cancer Society and the American Urological Association (AUA) have endorsed consideration of PSA screening based on age, other risk factors, and estimated life expectancy.

The utility of PSA screening depends on whether screening can lead to management changes that improve the net health outcome. Results from the National Health Service–supported Prostate Testing for Cancer and Treatment trial (2016) demonstrated no difference in 10-year prostate cancer mortality rates between the treatment strategies of active monitoring, radical prostatectomy, and external-beam radiotherapy in clinically localized prostate cancer detected by PSA testing.

Existing screening tools have led to unnecessary prostate biopsies. More than 1 million prostate biopsies are performed annually in the United States, with a resulting cancer diagnosis in 20% to 30% of men. About one-third of men who undergo prostate biopsy experience transient pain, fever, bleeding, and urinary difficulties. Serious biopsy risks (eg, bleeding or infection requiring hospitalization) are rare, with estimated rates ranging from less than 1% to 3%.

Given the risk, discomfort, burden of biopsy, and low diagnostic yield, there is a need for noninvasive tests that distinguish potentially aggressive tumors that should be referred for biopsy from clinically insignificant localized tumors or other prostatic conditions that do not need biopsy with the goal of avoiding low-yield biopsy.

The following PICOTS were used to select literature to inform this review.

Patients

The relevant populations are men for whom an initial prostate biopsy is being considered because of clinical symptoms (eg, difficulty with urination, elevated PSA).

The population for which these tests could be most informative is men in the indeterminate or “gray zone” range of PSA on repeat testing with unsuspecting DRE findings. Repeat PSA testing is important because results initially reported to be between 4 and 10 ng/mL frequently revert to normal. The gray zone for PSA

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

levels is usually between 3 or 4 ng/mL and 10 ng/mL, but PSA levels vary with age. Age-adjusted normal PSA ranges have been proposed but not standardized or validated.

Screening of men with a life expectancy of less than 10 years is unlikely to be useful because most prostate cancer progresses slowly. However, the age range for which screening is most useful is controversial. The ERSPC and Rotterdam trials observed benefits of screening only in men up to about 70 years old.

Interventions

For assessing future prostate cancer risk, numerous studies have demonstrated the association between many genetic and protein biomarker tests and prostate cancer. Commercially available tests for selection of men for initial prostate biopsy include those described in Table 1.

Table 1. Commercially Available Tests to Determine Candidate for Prostate Initial Biopsy

Test	Manufacturer	Description
4Kscore	OPKO lab	Blood test that measures 4 prostate-specific kallikreins, which are combined into an algorithm to produce a score
Prostate Health Index (phi)	Beckman Coulter	Blood test that combines several components of PSA with an algorithm to produce a score

PSA: prostate-specific antigen.

In addition to commercially available tests, single-nucleotide variant (SNV) testing as part of genome-scanning tests for prostate cancer risk assessment are offered by a variety of laboratories, such as Navigenics (now Life Technologies), LabCorp (23andme), and ARUP Laboratories (deCODE), as laboratory-developed tests.

Comparators

Standard clinical examination for determining who goes to biopsy might include DRE, review of history of PSA values, along with consideration of risk factors such as age, race, and family history. The ratio of free (or unbound) PSA to total PSA (percent free PSA) is lower in men who have prostate cancer than in those who do not. A percent free PSA cutoff of 25% has been shown to have a sensitivity and specificity of 95% and 20%, respectively, for men with total PSA levels between 4.0 ng/mL and 10.0 ng/mL.

The best way to combine all risk information to determine who should go to biopsy is not standardized. Risk algorithms have been developed that incorporate clinical risk factors into a risk score or probability. Two examples are the Prostate Cancer Prevention Trial (PCPT) predictive model and the Rotterdam Prostate Cancer risk calculator (also known as the ERSPC-Risk Calculator 4 [ERSPC-RC]). In 2016, the AUA and the Society of Abdominal Radiology recommended that high-quality prostate magnetic resonance imaging (MRI), if available, should be strongly considered in any patient with a prior negative biopsy who has persistent clinical suspicion for prostate cancer and who is under evaluation for a possible repeat biopsy.

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

Outcomes

The beneficial outcome of the test is to avoid a negative biopsy for prostate cancer. A harmful outcome is failure to undergo a biopsy that would be positive for prostate cancer, especially when disease is advanced or aggressive. Thus the relevant measures of clinical validity are the sensitivity and negative predictive value (NPV). The appropriate reference standard is biopsy, though prostate biopsy is an imperfect diagnostic tool. Biopsies can miss cancers and repeat biopsies are sometimes needed to confirm diagnosis. Detection rates vary by biopsy method and patient characteristics, with published estimates between 14% and 22% for the initial biopsy.

Timing

The timeframe of interest for calculating performance characteristics is time to biopsy result. Men who forgo biopsy based on test results could miss or delay diagnosis of cancer. Longer follow-up would be necessary to determine effects on overall survival.

Setting

Initial screening using PSA levels and DRE may be performed in the primary care setting with referral to specialty (urologist) care for suspicious findings and biopsy. Clinical practice on screening methods and frequency vary widely.

Kallikreins Biomarkers and 4Kscore Test

The 4Kscore test (OPKO Lab) uses results from a blood test to generate a risk score estimating the probability of finding high-grade prostate cancer (defined as a Gleason score ≥ 7) if a prostate biopsy were performed. The intended use of the test is to aid in a decision whether to proceed with a prostate biopsy. A kallikrein is a subgroup of enzymes that cleaves peptide bonds in proteins. The intact prostate-specific antigen and human kallikrein 2 (hK2) tests are immunoassays that employ distinct mouse monoclonal antibodies. The score combines the measurement of 4 prostate-specific kallikreins (total PSA, free PSA, intact PSA, hK2), with an algorithm including patient age, DRE (nodules or no nodules), and a prior negative prostate biopsy.

The manufacturer's website states that the ideal patient for the 4Kscore is one whose other test results are equivocal. The test is not intended for patients with a previous diagnosis of prostate cancer, who have had a DRE in the previous 4 days, who have received 5 α reductase inhibitor therapy in the previous 6 months, or who have undergone treatment for symptomatic benign prostatic hypertrophy in the previous 6 months.

Analytic Validity

Measures of analytic validity include sensitivity (detection rate), specificity (1 – false-positive rate); reliability (repeatability of test results), and assay robustness (resistance to small changes in preanalytic or analytic variables). As described above, the 4Kscore combines 4 blood biomarkers. Total and free PSA are measured using FDA-approved kits from Roche Diagnostics. Intact PSA and hK2 are proprietary assays validated by OPKO. Only one published study was found describing any components of analytic validity for a test of kallikrein biomarkers. In 2006, Vaisanen et al reported on results of a new method to reduce false

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272
 Original Effective Date: 10/20/2010
 Current Effective Date: 01/17/2018

high results by eliminating assay interference in measurement of intact, free PSA and free hK2. Using 1092 female heparin plasma samples as controls and 957 male samples, they optimized the protocol for immunoassays by replacing monoclonal capture or tracer antibodies with F(ab')₂ or recombinant Fab fragments. They tested the new method on another set of 444 samples and found that the optimized assay eliminated 70% to 85% of the falsely elevated results. Other measures of analytic validity were not found in the literature or the 4Kscore website. The laboratories that perform the analyses for 4Kscore are certified under the Clinical Laboratory Improvement Amendments.

Clinical Validity

At least 13 retrospective studies and a prospective study have estimated the performance characteristics of a risk score derived from 4 kallikrein (KLK) biomarkers. Many studies appear to be developmental work for the currently marketed version of the test. In general, the comparators used in these studies were other risk calculators or models that included terms for age, total PSA, and occasionally other risk factors. The reference standard was usually biopsy. Some studies performed in Sweden had long-term follow-up from a national registry of prostate cancer. The eligibility criteria included a lower limit of PSA (2 or 3 ng/mL) in most studies with no upper limit, and men with and without positive DRE. The studies included a mix of men who had or did not have previous PSA testing or biopsies. Mathematical methods used to calculate the KLK risk score varied across studies with respect to whether KLK values derived from plasma or serum measurements. Additionally, there was variance across studies in the additional risk factors included in the model (age, DRE, biopsies, other risk factors), and how KLK marker values were entered into the model (linearly, with splines or cubic splines). The area under the receiver operating characteristic (AUROC) curve, or a similar metric, was calculated in all studies. The estimated area under the curve (AUC) for the KLK model ranged from 0.72 to 0.90 and was numerically higher than the comparator in all studies except Carlsson et al (2013), who compared the KLK model with a clinical model including the length of benign tissue. However, the confidence intervals (CIs) for AUC of the KLK model frequently overlapped with those of the comparator. A few studies have provided results for the KLK model calculated with and without each of the 4 KLK. In many cases, the addition of intact PSA and hK2 did not significantly improve the model. In Bryant et al (2015), the CIs of the AUC for 4 KLK model overlapped considerably with a model that included age, total and free PSA for any grade, and high-grade cancer. Nordstrom et al (2015) included a comparison with another biomarker test (phi) and found both tests had very similar AUCs.

Our review of the clinical validity of the 4Kscore only includes studies that stated use of the marketed 4Kscore version of the KLK model. The marketed version of the test appears to have been used in 3 studies. Cutoffs for categorizing risk into low, medium, or high levels were only given in Konety et al (2015) and therefore sensitivity and NPV have not been calculated. Results of the studies are summarized in Table 2. Two studies were conducted in the United States and are discussed in the following paragraphs.

Table 2. Clinical Validity Studies of the 4Kscore for Diagnosing High-Grade Prostate Cancer

Study	Population	Reference Standard	Blinded Comparison With Reference Standard	Performance Characteristics (95% CI)
-------	------------	--------------------	--	--------------------------------------

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

Study	Population	Reference Standard	Blinded Comparison With Reference Standard	Performance Characteristics (95% CI)		
				4Kscore	Comparators	
Borque-Fernando et al (2016) (Spain)	51 men scheduled for biopsy for suspicion of prostate cancer	Clinical consensus of 4 urologists after review of biopsy of ≥10 cores	NR	AUC=0.79 (0.66 to 0.89)	ERSPC-RC risk calculator AUC=0.81 (0.67 to 0.90)	PCPT risk calculator AUC=0.75 (0.61 to 0.86)
Konety et al (2015) (U.S.)	171 men of high-volume 4Kscore users who had biopsy results	Biopsy	NR	Low vs intermediate/high risk ^a : • Sen=0.97 (0.87 to 0.99) • Spec=0.12 (0.07 to 0.19) • PPV=0.37 (0.29 to 0.45) • NPV=0.87 (0.58 to 0.98)	NA	
Parekh et al (2015) (U.S.)	1012 men scheduled for biopsy regardless of PSA or clinical findings	Biopsy with ≥10 cores	Yes	AUC=0.82 (0.79 to 0.85)	Risk model without intact PSA and hK2 AUC=0.75 (0.71 to 0.79)	PCPT modified risk calculator ^b AUC=0.74 (NR)

AUC: area under the curve; CI: confidence interval; ERSPC-RC: European Randomized Study of Screening for Prostate Cancer Risk Calculator; NA: not available; NPV: negative predictive value; NR: not reported; PCPT: Prostate Cancer Prevention Trial; PPV: positive predictive value; PSA: prostate-specific antigen; Sen: Sensitivity; Spec: Specificity.

^a Calculated from value provided in article considering low risk to be a negative result and intermediate/high risk to be a positive result.

^b Excluding the term for family history because it was not known in this cohort.

Performance of the 4Kscore test was validated in 1012 patients enrolled in a blinded, prospective study at 26 urology centers in the United States. Enrollment was open to all men scheduled for a prostate biopsy, regardless of age, PSA level, DRE, or prior prostate biopsy. Each patient underwent a transrectal ultrasound-guided prostate biopsy of at least 10 cores. A blinded blood sample collected before biopsy was sent to OPKO Lab for the 4 KLK markers. Results of the KLK markers, prostate biopsy histopathology, patient age, DRE, and prior biopsy status were unblinded and analyzed.

Most participants (86%) were white; 85 (8%) African American men were included. At baseline, 247 (24%) men had an abnormal DRE, 348 (34%) had a PSA level less than 4 ng/mL, and 104 (10%) had PSA level greater than 10 ng/mL. Approximately 25% of the men were older than 70 years. Biopsies were negative in 54% (n=542) of cases, and showed low-grade (all Gleason grade 6) prostatic cancer in 24% (n=239) and high-grade cancer in 23% (n=231). Statistical analysis of 4Kscore test clinical data had AUROC of 0.82 (95% CI, 0.79 to 0.85) for the detection of high-grade prostate cancer; the AUROC for the PCPT risk calculator model was 0.74, but a precision estimate was not given.

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

Subsection Summary: Clinical Validity

The intended use population is not well defined. In addition, there is uncertainty regarding clinical performance characteristics such as sensitivity, specificity, and predictive value due to the following factors: a lack of standardization of cutoffs to recommend biopsy, study populations including men with low (<4 ng/mL) and high (>10 ng/mL) baseline PSA levels, positive DRE results likely outside the intended use population, and lack of comparison with models using information from standard clinical examination. African Americans have a high burden of morbidity and mortality but were not well represented in the study populations. The evidence needed to conclude clinical validity is insufficient. Longer term data on the incidence of prostate cancer in men who did not have a biopsy following testing with the marketed version of 4Kscore are not available. However, the Stattin et al (2015) case-control study, which was a nested cohort study of more than 17,000 Swedish men, estimated that, for men ages 60 with PSA levels of 3 or higher and a KLK risk score less than 10%, the risk of metastasis at 20 years was 1.95% (95% CI, 0.64% to 4.66%).

Clinical Utility

No studies reporting direct evidence of utility for clinical outcomes were found. Various cutoffs for the KLK probability score were used in decision curve analyses to estimate the number of biopsies vs cancers missed. Parekh et al (2015) estimated that 307 biopsies could have been avoided and 24 cancer diagnoses would have been delayed with a 9% 4Kscore cutoff for biopsy and 591 biopsies would have been avoided with 48 diagnoses delayed with a 15% cutoff. However, inferences on clinical utility cannot be made due to deficiencies in estimating the clinical validity that are described in the previous section.

Konety et al (2015) reported on results of a survey of 35 U.S. urologists identified through the 4Kscore database at OPKO Lab as belonging to practices that were large users of the test. All 611 patients of participating urologists who were referred for abnormal PSA or DRE and had a 4Kscore test were included. Six percent of the men had an abnormal DRE; the distribution of PSA levels was not reported. Urologists, who received the 4Kscore as a continuous risk percentage, were retrospectively asked about their plans for biopsy before and after receiving the test results and whether the 4Kscore test results influenced their decisions. Scores were grouped into 3 risk categories: less than 7.5%, low risk; 7.5% to 19.9%, intermediate risk; and 20% or more, high risk. The physicians reported that the 4Kscore results influenced decisions in 89% of men and that the test led to a 64.6% reduction in prostate biopsies. The 4Kscore risk categories correlated highly ($p < 0.001$) with biopsy outcomes in 171 men with biopsy results. Calculated performance characteristics are shown in Table 2. No other risk calculators were included as comparators.

Absent direct evidence of clinical utility, a chain of evidence might be constructed. The 4Kscore test is associated with a diagnosis of aggressive prostate cancer. The incremental value of the 4Kscore concerning clinical examination and risk calculators in the intended use population is unknown due to deficiencies in estimating clinical validity described in the previous section. There is no prospective evidence that use of 4Kscore changes management decisions. The chain of evidence is incomplete.

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

Section Summary: Kallikreins Biomarkers and 4Kscore Test

Published data on most components of the analytic validity of the 4Kscore test are lacking. At least 13 studies have reported on clinical validity of the KLK biomarkers, but only three clearly used the marketed version of the 4Kscore test. The eligibility criteria for these studies had a lower limit for screening PSA but no upper limit. Given that the test manufacturer's website states that the test is for men with inconclusive results, the inclusion of men with PSA levels greater than 10 ng/mL and positive DRE in the validation studies is likely not reflective of the intended use population. Studies that provide data on the incremental value of the components of the test show only small improvements with the intact PSA and hKA components (components specific to the 4Kscore). The 2 studies performed in U.S. men did not provide estimates (with confidence intervals) of validity compared with a standard clinical examination with a ratio of free or unbound PSA to total PSA (percent free PSA). Very few data are available on longer term clinical outcomes of men who were not biopsied based on 4Kscore results. No direct evidence supports the clinical utility of the test, and the chain of evidence is incomplete due to the limitations in estimates of clinical validity and utility.

proPSA and Prostate Health Index

The Prostate Health Index (phi; Beckman Coulter) is an assay combining results of 3 blood serum immunoassays (total PSA, free PSA, [-2]proPSA [p2PSA]) numerically to produce a "phi score." This score is calculated in a routine laboratory using Beckman Coulter equipment and software with the phi algorithm incorporated in the software using the following formula: $([-2]proPSA/free\ PSA) \times \sqrt{total\ PSA}$. It has been suggested that the PSA isoform p2PSA might better distinguish between prostate cancer and benign prostatic conditions.

The phi score has been approved by FDA for distinguishing prostate cancer from benign prostatic conditions in men 50 years and older with above-normal total PSA readings between 4.0 ng/mL and 10 ng/mL who have had a negative DRE. The manufacturer's website states that the test gives men "accurate information on what an elevated PSA level might mean and the probability of finding cancer on biopsy" and when "combined with family and patient history, the phi results can be used to determine the best individualized patient management decisions."

Analytic Validity

The FDA Summary of Safety and Effectiveness Data (SSED) provides data on the analytic validity of the assay. The analytic validity was also reviewed by the National Institute for Health and Care Excellence (NICE) in 2015. The limit of blank of p2PSA was 0.5 pg/mL, the limit of detection was 0.7 pg/mL, and the limit of quantification was 3.23 pg/mL. Accuracy was calculated by the percentage recovery of measured p2PSA pg/mL in 6 male serum samples containing different known amounts of purified p2PSA.

Clinical Validity

Systematic Reviews

Several systematic reviews and meta-analyses have described the clinical validity of p2PSA and phi. The characteristics of the reviews are shown in Table 3. The reviews cover studies reported between 1990 and

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272
 Original Effective Date: 10/20/2010
 Current Effective Date: 01/17/2018

2014. All primary studies were observational and most were retrospective. All reviews included studies of men with a positive, negative, or inconclusive DRE; only 2 of the 5 reviews restricted eligibility to studies including PSA levels between 2 ng/mL and 10 ng/mL. The Wang et al (2014) review included only studies that had sufficient information to distinguish aggressive from indolent prostate cancer. The 2 most recent reviews (Nicholson et al [2015], Pecoraro et al [2016],) included most of the studies covered in the older reviews; we review them in more detail below.

Nicholson et al performed a systematic review and health technology assessment commissioned to support the development of NICE guidance (2015) for diagnosing prostate cancer with the prostate cancer antigen 3 (PCA3) and phi assays. The search included the databases MEDLINE, EMBASE, the Cochrane Library, Web of Science, Medion, Aggressive Research Intelligence Facility database, ClinicalTrials.gov, International Standard Randomised Controlled Trial Number Register, and World Health Organization International Clinical Trials Registry Platform. The review included studies of men whose initial prostate biopsies were negative or equivocal and studies using clinical examination or clinical examination plus MRI as comparators. Pooled estimates were not reported due to heterogeneity.

Pecoraro et al performed a search of MEDLINE, EMBASE, Web of Science, Scopus, and the Cochrane Register of Diagnostic Test Accuracy Studies for studies including men with a PSA level between 2 ng/mL and 10 ng/mL. The quality of each study was assessed using the Quality Assessment of Diagnostic Accuracy Studies (QUADAS-2) checklist, and the evidence was evaluated using the GRADE approach. Random-effects bivariate models were used to calculate pooled estimates.

Table 3. Characteristics of Systematic Reviews on the Clinical Validity of phi for Diagnosing Prostate Cancer

Study	Dates	Key Inclusion Criteria ^a	Design
Nicholson et al (2015)	2000-2014	Initial prostate biopsy was negative or equivocal, 6+ cores in initial biopsy	Prospective and mixed (prospective/retrospective) OBS
Pecoraro et al (2016)	2003-2014	PSA 2-10 ng/mL	Prospective, retrospective, and mixed (prospective/retrospective) OBS
Bruzzese et al (2014)	2009-2013	TRUS biopsy (6+ cores) for diagnosis; PSA 2-10 ng/mL; first biopsy	Retrospective and prospective OBS
Wang et al (2014)	2000-2014	Biopsy reference standard	Prospective, retrospective, and mixed (prospective/retrospective) OBS
Filella et al (2013)	1990-2011	Biopsy reference standard	Retrospective and prospective OBS

OBS: observational; PSA: prostate-specific antigen; TRUS: transrectal ultrasound.

^a Results from all studies were with or without digital rectal exam.

Results of the systematic reviews and meta-analyses are shown in Table 4. Pecoraro et al included 17 studies with 6912 men. They rated most of the primary studies as low quality due to the design (most were retrospective), lack of blinding of outcome assessors to reference standard results, lack of clear cutoffs for diagnosis, and lack of explicit diagnostic question. Pooled estimates had high heterogeneity across studies but with generally low specificity of phi at 90% sensitivity.

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

Nicholson et al included 4 studies with 767 men that included estimates of clinical assessment alone vs clinical assessment plus phi. They concluded that the implication of adding phi to clinical assessment was unclear. Due to heterogeneity in cutoffs used in the primary studies, it was not possible to identify thresholds to use in a clinical setting, and the clinical relevance of many reported outcomes was unclear.

Table 4. Results of Systematic Reviews on the Clinical Validity of phi for Diagnosing Prostate Cancer

Study	Studies	N (Range)	Outcomes	Results (95% CI)
Pecoraro et al (2016)	17	6912 (63-1091)	<ul style="list-style-type: none"> Diagnostic performance Any prostate cancer 	Pooled specificity at 90% sensitivity: <ul style="list-style-type: none"> phi=31% (29% to 33%) total PSA=25% (23% to 27%)
Nicholson et al (2015)	4	767 (95-280)	<ul style="list-style-type: none"> Diagnostic performance Any prostate cancer 	<ul style="list-style-type: none"> No pooled estimates AUC range: <ul style="list-style-type: none"> phi plus clinical assessment, 0.65-0.81 Clinical assessment alone, 0.62-0.75 Derived sensitivity (at given specificity) <ul style="list-style-type: none"> phi plus clinical assessment, 42% (at 80%), 25% (at 90%), 19% (at 95%) Clinical assessment alone, 48% (at 80%), 23% (at 90%), 17% (at 95%)
Bruzzese et al (2014)	8	3173 (64-896)	<ul style="list-style-type: none"> Diagnostic performance Any prostate cancer 	<ul style="list-style-type: none"> Significant heterogeneity Sensitivity range: <ul style="list-style-type: none"> phi, 31%-90% %fPSA, 12%-90% Specificity range: <ul style="list-style-type: none"> phi, 30%-90% %fPSA, 11%-90% AUC: <ul style="list-style-type: none"> phi=0.74 (0.70 to 0.77) %fPSA=0.63 (0.58 to 0.67)
Wang et al (2014)	12		<ul style="list-style-type: none"> Diagnostic performance, high-grade (Gleason score ≥ 7) prostate cancer 	<ul style="list-style-type: none"> Pooled sensitivity, 90% (87% to 92%) Pooled specificity, 17% (14% to 19%) Diagnostic OR=3.06 (1.61 to 5.84) Pooled AUC=0.67 (0.57 to 0.77)
Filella et al (2013)	13	3928 (63-1091)	<ul style="list-style-type: none"> Diagnostic performance Any prostate cancer 	<ul style="list-style-type: none"> Significant heterogeneity Pooled specificity at 90% sensitivity, 32% (29% to 34%)

AUC: area under the curve; CI: confidence interval; OR: odds ratio; PSA: prostate-specific antigen; %fPSA: ratio of free (unbound) PSA to total PSA.

Clinical Studies

The pivotal study described in the FDA SSED included men 50 years and older with nonsuspicious DRE and PSA levels between 4 ng/mL and 10 ng/mL who had a histologically confirmed diagnosis. The null hypothesis was that the phi specificity at 95% sensitivity would be no greater than the specificity of percent

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

free PSA. Seven sites in the United States enrolled 658 men between 2008 and 2009 (97% enrolled prospectively, 3% enrolled retrospectively). Eighty-one percent of participants were white, 5% were African American, and 1% was Asian. At 95% sensitivity, using a phi cutoff of 22.1, the specificity was 14.1% (precision not reported) for phi compared with 9.9% for percent free PSA. AUC was 0.71 (95% CI, 0.67 to 0.75) for phi compared with 0.65 (95% CI, 0.61 to 0.69) for percent free PSA.

Additional studies have been published since the systematic reviews. In 2015 Fossati et al conducted a case-control study with 1036 European men younger than 60 years of age. They reported that phi had a higher AUC (0.70; 95% CI, 0.64 to 0.76) than total PSA (0.55; 95% CI, 0.48 to 0.61) in men younger than 60 years for detecting any prostate cancer. At 91% sensitivity, phi and total PSA had similar specificity (11.1% [95% CI, 6.8% to 16.8%] vs 10.5% [95% CI, 6.4% to 16.1%]) and an NPV (76.0% [95% CI, 59.3% to 92.7%] vs 75.0% [95% CI, 57.7% to 92.3%]), respectively. At the best combination of sensitivity and specificity (phi cutoff ≥ 41.2 , total PSA cutoff ≥ 5.72), phi had a sensitivity of 64.2% (95% CI, 51.5% to 75.5%), a specificity of 63.2% (95% CI, 55.5% to 70.4%), and an NPV of 81.8% (95% CI, 75.2% to 88.4%) while total PSA had a sensitivity of 52.2% (95% CI, 39.7% to 64.6%), a specificity of 52.0% (95% CI, 44.3% to 59.7%), and an NPV of 73.6% (95% CI, 65.7% to 81.4%). A decision curve analysis found that using a model with age, prostate volume, total PSA, free PSA, percent free PSA, and phi with a probability cutoff of 10% would avoid 13% of biopsies while missing 0% of cancers; a cutoff of 20% would avoid 51% of biopsies while missing 18% of cancers; and a cutoff of 50% would avoid 94% of biopsies while missing 66% of cancers.

In 2016 Boegemann et al reported on results of a study of 769 European men ages 65 years and younger scheduled for initial or repeat prostate biopsy who were prospectively and retrospectively enrolled. The investigators compared phi with other PSA measures for detecting clinically significant vs insignificant cancer (PRIAS-criteria: T-stage T1c/T2; Gleason score ≤ 6 ; number of positive cores per biopsies ≤ 2 ; total PSA ≤ 10 ng/mL; PSA density < 0.2 ng/mL). The AUC for phi (0.72; 95% CI, 0.68 to 0.76) was higher than that for total PSA (0.62; 95% CI, 0.58 to 0.66) or percent free PSA (0.64; 95% CI, 0.60 to 0.68).

Morote et al (2016) reported numerically higher but not statistically significantly higher AUC for phi compared with total PSA or total free PSA for detecting aggressive prostate cancer in 357 men with PSA levels between 3 ng/mL and 10 ng/mL scheduled for first biopsy in a retrospective study in Spain. Similarly, Yu et al (2016) reported numerically but not statistically significantly higher AUC for phi vs total PSA in 114 men in China with PSA levels between 2 ng/mL and 10 ng/mL and negative DRE.

Porpiglia et al (2016) reported on the results of an observational retrospective study of 120 men with prostate cancer who received radical prostatectomy but were eligible for active surveillance. Multiparametric MRI (mpMRI), phi, and PCA3 were performed on the single cohort. The base model for predicting pathologically confirmed significant prostate cancer (PCSPCa) had an AUC of 0.71. Relative to the base model for predicting PCSPCa, phi increased the AUC by 4% (0.75; $p < 0.01$), PCA3 increased the AUC by 1% (0.72; $p < 0.01$), and mpMRI increased the AUC by 7% (0.78; $p < 0.01$). Results of mpMRI provided the

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

greatest net benefit in predicting the presence of PCSPCa while phi provided a small incremental benefit in prediction.

Subsection Summary: Clinical Validity

Many studies and systematic reviews of these studies have reported on the clinical validity of phi. In general, the comparator was a component of PSA (total PSA, free PSA) but have not included other risk factors from a standard clinical exam. Most of the primary studies included men with positive, negative, and inconclusive DRE and men with PSA levels outside of the 4- to 10-ng/mL range. African Americans have a high burden of morbidity and mortality but were not well represented in the study populations. There is no standardization of cutoffs used in a clinical setting for diagnosis and data on the diagnostic accuracy of phi for distinguishing clinically significant from insignificant cancer are lacking. A direct comparison between phi and mpMRI demonstrated a higher net benefit for mpMRI in predicting PCSPCa.

Clinical Utility

No studies directly measuring the effect of phi on clinical outcomes were found. A chain of evidence might be used to demonstrate clinical utility if each link in the chain is intact. The phi test is associated with a diagnosis of prostate cancer, although data on association with a diagnosis of aggressive prostate cancer are lacking. The phi test provided better diagnostic information than other measures of PSA alone, but decisions made with phi result plus other risk factors from clinical examination were not provided in most studies. Optimal cutoffs for classifying men into risk groups have not been standardized. No studies were found describing differences in management based on phi risk assessment. The chain of evidence is incomplete.

Section Summary: proPSA and Prostate Health Index

The analytic validity of phi has been established. At least 4 systematic reviews including a NICE assessment have been reported and included many primary studies. In general, selected studies included some men outside of the intended use population (PSA levels outside of the 4- to 10-ng/mL range and abnormal DRE). Comparisons with diagnosis with clinical examination were lacking. The cutoffs for categorizing men into risk groups in clinical practice have not been standardized, and therefore there is heterogeneity in reporting of performance characteristics and decision curve analyses.

BIOMARKER TESTING FOR SELECTION OF MEN FOR REPEAT PROSTATE BIOPSY

Clinical Context and Test Purpose

The purpose of genetic and protein biomarker testing for prostate cancer is to inform the selection of men who should undergo repeat biopsy. The conventional decision-making tools for identifying men for prostate biopsy include DRE, serum PSA, and patient risk factors such as age, race, and family history of prostate cancer are previously described in the section for selection of men for initial prostate biopsy.

Given the risk, discomfort, burden of biopsy, and the low diagnostic yield, there is a need for noninvasive tests that distinguish potentially aggressive tumors that should be referred for rebiopsy from clinically

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272
Original Effective Date: 10/20/2010
Current Effective Date: 01/17/2018

insignificant localized tumors or other prostatic conditions that do not need rebiopsy with the goal of avoiding low-yield biopsy.

The following PICOTS were used to select literature that provides evidence relevant to this review.

Patients

The relevant populations are men for whom a rebiopsy is being considered because the results of an initial prostate biopsy were negative or equivocal and other clinical symptoms remain suspicious.

Interventions

For assessing future prostate cancer risk, numerous studies have demonstrated the association between many genetic and protein biomarker tests and prostate cancer. Commercially available tests for selection of men for repeat prostate biopsy include those described in Table 5.

Table 5. Commercially Available Tests to Determine Candidate for Prostate Repeat Biopsy

Test	Manufacturer	Description
4Kscore	OPKO lab	Blood test that measures 4 prostate-specific kallikreins, which are combined into an algorithm to produce a score
Progensia	<ul style="list-style-type: none"> Hologic Gen-Probe Many labs offer <i>PCA3</i> tests (eg, ARUP Laboratories, Mayo Medical Laboratories, LabCorp) 	Urine test that measures <i>PCA3</i> mRNA
ConfirmMDx	MDxHealth	Measures hypermethylation of 3 genes in tissue sample
Prostate Health Index (phi)	Beckman Coulter	Blood test that combines several components of PSA with an algorithm to produce a score
Prostate Core Mitomics Test (PCMT)	Mitomics (formerly Genesis Genomics)	Measures deletions in mitochondrial DNA by polymerase chain reaction in tissue sample

PSA: prostate-specific antigen.

In addition to commercially available tests, SNV testing as part of genome-scanning tests for prostate cancer risk assessment is offered by a variety of laboratories, such as Navigenics (now Life Technologies), LabCorp (23andme), and ARUP Laboratories (deCODE), as laboratory-developed tests.

Comparators

Standard clinical examination for determining who goes to biopsy might include DRE, review of history of PSA values, along with consideration of risk factors such as age, race, and family history. The ratio of free (unbound) PSA to total PSA is lower in men who have prostate cancer than in those who do not. A percent

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

free PSA cutoff of 25% has been shown to have a sensitivity and specificity of 95% and 20%, respectively, for a group of men with total PSA levels between 4.0 ng/mL and 10.0 ng/mL.

The best way to combine all of the risk information to determine who should go to biopsy is not standardized. Risk algorithms have been developed that incorporate clinical risk factors into a risk score or probability. Two examples are the PCPT predictive model and the ERSPC-RC). The AUA and the Society of Abdominal Radiology recently recommended that high-quality prostate MRI, if available, should be strongly considered in any patient with a prior negative biopsy who has persistent clinical suspicion for prostate cancer and who is under evaluation for a possible repeat biopsy.

Outcomes

The beneficial outcome of the test is to avoid a negative biopsy for prostate cancer. A harmful outcome is failure to undergo a biopsy that would be positive for prostate cancer, especially when disease is advanced or aggressive. Thus the relevant measures of clinical validity are the sensitivity and negative predictive value. The appropriate reference standard is biopsy, though prostate biopsy is an imperfect diagnostic tool. Biopsies can miss cancers and repeat biopsies are sometimes needed to confirm the diagnosis. Detection rates vary by biopsy method and patient characteristics, with published estimates between 10% and 28% for a second biopsy and 5% and 10% for a third biopsy.

Timing

The timeframe of interest for calculating performance characteristics is time to biopsy result. Men who forgo biopsy based on test results could miss or delay diagnosis of cancer. Longer follow-up would be necessary to determine effects on overall survival.

Setting

Screening using PSA levels and DRE may be performed in the primary care setting with referral to specialty (urologist) care for suspicious findings and biopsy. Clinical practice on screening methods and frequency vary widely.

Progenesa PCA3 Assay

PCA3 is overexpressed in prostate cancer, and *PCA3* messenger RNA can be detected in urine samples collected after prostate massage. When normalized using PSA to account for prostate cells released into the urine (*PCA3* score), the test has significantly improved specificity compared with serum PSA and may better discriminate patients with benign findings on (first or second) biopsy from those with malignant biopsy results.

The Progenesa *PCA3* assay (Hologic Gen-Probe) has been approved by FDA to aid in the decision for repeat biopsy in men 50 years or older who have had 1 or more negative prostate biopsies and for whom a repeat biopsy would be recommended based on current standard of care. The Progenesa *PCA3* assay should not be used for men with atypical small acinar proliferation on their most recent biopsy. The manufacturer's website states that the test is intended to identify men who have negative first biopsy results

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

to determine who needs a follow-up biopsy and that a PCA3 score less than 25 is associated with a decreased likelihood of a positive biopsy.

Analytic Validity

The analytic validity of the ProgenSA PCA3 has been reviewed by FDA and NICE. Limit of blank was reported as 0.50 pg/mL, limit of detection was 0.69 pg/mL, and limit of quantitation was 3.23 pg/mL. No assay interference was recorded in the SSED report. The SSED report included carryover studies with a 0% false-positive rate for negative samples interspersed with high-titer samples. Accuracy was calculated by percent recovery of PCA3 compared with ultraviolet-determined copies per milliliters of an 8-member panel of female urine spiked with in vitro transcript; the minimum was 90% and the maximum was 118%. Precision as measured by percent coefficient of variation for within- and between-laboratory variation ranged from 12.3 to 25 for PCA3 score in 3 control samples. Linearity was assessed with 11 samples with in vitro transcripts in processed female urine, and the deviation from linearity for PCA3 was less than 9%.

Clinical Validity

Systematic Reviews

Several systematic reviews and meta-analyses have described the clinical validity of ProgenSA. The characteristics of the reviews are in Table 6. The reviews cover studies reported up to 2014. All primary studies selected were observational, although 1 study used the placebo arm from a randomized controlled trials and a validation trial not included in the reviews are described below. Reviewers selected studies of men with positive, negative, or inconclusive DRE without restrictions on PSA levels. The 2 most recent reviews (Cui et al [2016], Nicholson et al [2015]) are detailed below.

In 2016 Cui et al reported on results of a systematic review that searched PubMed and EMBASE for case-control or cohort studies. Quality was assessed using the QUADAS tool. Pooled estimates were calculated using random-effects models and summarized ROCs when evidence of threshold effect was detected. Nicholson et al was described in the section on phi. In brief, the NICE assessment included studies with men for whom initial prostate biopsy results were negative or equivocal.

Table 6. Characteristics of Systematic Reviews on the Clinical Validity of ProgenSA for Diagnosing Prostate Cancer

Study	Dates	Key Inclusion Criteria	Design
Cui et al (2016)	Up to 2014	Biopsy as reference standard	Prospective, retrospective (case-control or cohort) OBS
Nicholson et al (2015) (NICE)	2000-2014	Initial prostate biopsy negative or equivocal, 6+ cores in initial biopsy, with/without DRE	Prospective and mixed (prospective/retrospective) OBS (1 included a cohort from an RCT)
Bradley et al (2013) (AHRQ)	1990-2012	Comparative data	Cohort studies
Ruiz-Aragon et al (2010)	2000-2009	Men undergoing prostate cancer screening, biopsy reference standard	Prospective and retrospective OBS

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

AHRQ: Agency for Healthcare Research and Quality; DRE: digital rectal exam; NICE: National Institute for Health and Care Excellence; OBS: observational; RCT: randomized controlled trial.

Results from the systematic reviews are shown in Table 7. Cui et al included 46 studies with over 12,000 men. The quality of the selected studies was rated as moderate to high. The most common PCA3 cutoff for categorizing low and high risk was 35; 25 studies had a PCA3 cutoff of 35. Most were performed in the United States and Europe; five were conducted in Asia. The estimates of AUC were lower for studies including men having repeated (0.68; 95% CI, 0.67 to 0.70) vs initial (0.80; 95% CI, 0.78 to 0.82) biopsies. AUC values were 0.74 (95% CI, 0.73 to 0.76) for studies with a cutoff value of 35 and 0.77 (95% CI, 0.75 to 0.79) for studies with a cutoff value equal to 35, although the group with varying cutoff (\neq 35) had a greater range and more variable performance estimates.

Nicholson et al included 13 studies describing 11 cohorts including one from the placebo arm of a randomized controlled trial. Reviewers found that referral criteria for repeat biopsy were often unclear and varied across studies. The criteria also differed based on whether normal or abnormal DREs were included, and the mean or median PSA, when reported, ranged from 4.9 to 11.0 ng/mL. The prevalence of cancer on repeat biopsy varied from 11.4% to 68.3%. Meta-analyses were not performed due to heterogeneity. Some studies used PCA3 scores as a continuous variable and others created risk categories. The addition of PCA3 to clinical assessment, as a continuous or categorical variable, generally led to improvement in AUC. The comparisons were mixed concerning the diagnostic odds ratio (OR), although most studies found increased diagnostic accuracy for PCA3 plus clinical assessment compared with clinical assessment alone. Studies that fixed sensitivity and derived specificity and those that reported decision curve analysis had mixed results. Reviewers concluded that the clinical benefit of PCA3 in combination with clinical assessment was not confirmed.

Table 7. Results of Systematic Reviews on the Clinical Validity for PCA3 or Progenza for Diagnosing Prostate Cancer

Study	Studies	N (Range)	Outcomes	Results (95% CI)
Cui et al (2016)	46	12,295 (NR)	<ul style="list-style-type: none"> Diagnostic performance Any prostate cancer 	<ul style="list-style-type: none"> Sensitivity range, 47%-95% Significant heterogeneity and threshold effect Pooled estimate, 0.65 (0.63 to 0.66) Specificity range, 22%-100% Significant heterogeneity and threshold effect Pooled estimate: 73% (72% to 74%) Negative LR: Pooled estimate, 48% (44% to 52%) AUC=0.75 (0.74 to 0.77)
Nicholson et al (2015) (NICE)	11	3336 (41-1072)	<ul style="list-style-type: none"> Diagnostic performance Any prostate cancer 	<ul style="list-style-type: none"> AUC range: Clinical assessment alone, 0.55-0.75 Clinical assessment plus PCA3, 0.61-0.76 Derived sensitivity (at given specificity) range: Clinical assessment alone, 44%-48% (at 80%) Clinical assessment plus PCA3, 39%-46% (at 80%)
Bradley et al (2013)	43	9719 biopsies or prostatectomies	<ul style="list-style-type: none"> Diagnostic performance 	<ul style="list-style-type: none"> Derived sensitivity (at given specificity) tPSA=91% (at 20%)

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

(AHRQ)	(32-1246)		
Ruiz-Aragon et al (2010)	14 3467 (30-563)	<ul style="list-style-type: none"> Any prostate cancer and aggressive prostate cancer Diagnostic performance Any prostate cancer 	<ul style="list-style-type: none"> PCA3=96% (at 20%) Unable to compare performance for aggressive prostate cancer Sensitivity range, 47%-82% Pooled sensitivity, 85% (84% to 87%) Specificity range, 56%-89% Pooled specificity, 96% (96% to 97%) Pooled negative LR=0.15 (0.13 to 0.18)

AHRQ: Agency for Healthcare Research and Quality; AUC: area under the curve; CI: confidence interval; LR: likelihood ratio; NICE: National Institute for Health and Care Excellence; NR: not reported; tPSA: total prostate-specific antigen.

Randomized Controlled Trials

In 2014, the National Cancer Institute conducted a prospective trial to validate the diagnostic use of the PCA3 assay to complement PSA-based detection of prostate cancer. The target population included men who had been screened for prostate cancer, primarily with a PSA test, some of whom had undergone a previous prostate biopsy. The study included 859 men from 11 centers in the United States. The primary study end point was the diagnosis of prostate cancer on biopsy, and the secondary study end point was the diagnosis of high-grade prostate cancer, defined as a Gleason score greater than 6. The primary analyses, including PCA3 thresholds, were determined a priori, and statistical power was based on independent analyses of prevalidation data from similar cohorts. Of the men in the study, 562 were presenting for their initial prostate biopsy. PPV was 80% (95% CI, 72% to 86%), and using a PCA3 score of more than 60, diagnostic sensitivity and specificity of PCA3 score were 42% (95% CI, 36% to 48%) and 91% (95% CI, 87% to 94%), respectively. For patients who underwent a repeat biopsy, the NPV was 88% (95% CI, 81% to 93%), and, by using a PCA3 score of less than 20, sensitivity and specificity were 76% (95% CI, 64% to 86%) and 52% (95% CI, 45% to 58%), respectively. For the detection of high-grade cancer, performance of PCPT risk calculator was improved by adding PCA3 results to the risk calculator factors, with an AUC improvement of 0.74 to 0.78 for initial biopsy and 0.74 to 0.79 on repeat biopsy (p<0.003).

Clinical Studies

The pivotal study describing in the FDA SSED for Progenesa included 495 men from 15 clinical sites who had at least 1 negative prostate biopsy and whose urologist recommended repeat biopsy. Prostate biopsy was performed using each site's local standard procedure. A total of 433 (87.5%) were white, 45 (9.1%) were African American or black, and 11 (2.2%) were Asian. A valid PCA3 score and biopsy result were available for 466 men. Using a PCA3 cutoff score of 25, the performance characteristics for positive biopsy were as follows: sensitivity, 77.5% (95% CI, 68.4% to 84.5%); specificity, 57.1% (95% CI, 52.0% to 62.1%); PPV, 33.6% (95% CI, 30.0% to 37.2%); and NPV, 90.0% (95% CI, 86.5% to 93.1%). In other words, 208 men in the study might have been spared an unnecessary repeat biopsy if a cutoff of 25 was used to recommend repeat biopsy. On the other hand, 23 of the men who had a biopsy positive for prostate cancer might have had their diagnosis delayed due to negative PCA3 result.

Clinical studies have compared PCA3 results with clinical examination and risk calculators; those focused on distinguishing between aggressive and indolent cancer are particularly relevant. Ankerst et al (2008) reported that incorporating the PCA3 score into the PCPT risk calculator improved the diagnostic accuracy

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

of the calculator (from an AUC of 0.653 to 0.696). Chun et al (2009), using a multivariate nomogram, demonstrated a 5% gain in predictive accuracy when *PCA3* was incorporated with other predictive variables such as age, DRE results, PSA levels, prostate volume, and biopsy history. In a 2011 study of 218 patients with PSA levels of 10 ng/mL or less, Perdonà et al performed a head-to-head comparison of these 2 risk assessment tools and suggested both might have value in clinical decision making.

Several studies have evaluated the *PCA3* score as a tool for distinguishing between patients with indolent cancers who may need only active surveillance and those with aggressive cancers who warrant aggressive therapy. Three studies from 2008—Haese et al, Nakanishi et al, and Whitman et al—demonstrated an association between *PCA3* scores and evidence of tumor aggressiveness. However, these findings were not confirmed in a 2006 study by Bostwick et al or a 2008 study by van Gils et al. Au-prich et al (2011) reported that *PCA3* scores appeared to enhance identification of indolent disease but not pathologically advanced or aggressive cancer.

Tosoian et al (2010) reported on a short-term prospective cohort study evaluating the *PCA3* score in relation to outcomes in an active surveillance program involving 294 patients. The *PCA3* score did not distinguish between patients who had stable disease and those with more aggressive features.

Clinical Utility

Clinical utility studies using assay results for decision making for initial biopsy, repeat biopsy, or treatment have not been reported, nor have studies of the effects of using assay results on clinical outcomes. Several studies using decision analysis to estimate the cost-benefit tradeoff between reduction in unnecessary biopsies and missed prostate cancers have been published. One group reported potential reductions in unnecessary biopsies of 48% to 52%, with attendant increases in missed prostate cancers of 6% to 15% using either a *PCA3*-based nomogram or *PCA3* level corrected for prostate volume (*PCA3* density). Although both studies were prospective, neither assessed utility of the test for clinical decision making because all patients underwent biopsy. Merdan et al (2015) used decision analysis to simulate long-term outcomes associated with use of the *PCA3* score to trigger repeat biopsy compared with the PCPT risk calculator in men with at least 1 previous negative biopsy and elevated PSA levels. They estimated that incorporating the *PCA3* score of 25 (biopsy threshold) into the decision to recommend repeat biopsy could avoid 55.4% of repeat biopsies, with a 0.93% reduction in the 10-year survival rate.

Given the lack of direct evidence of utility, a chain of evidence would be needed to demonstrate clinical utility. The *PCA3* test is associated with a diagnosis of prostate cancer, although data on its association with a diagnosis of aggressive prostate cancer are lacking. The *PCA3* test provided better diagnostic information than other measures of PSA, but comparison with decisions made using risk factors from clinical examination was not provided in most studies. No prospective studies were found describing differences in management based on *PCA3* risk assessment. The chain of evidence is incomplete.

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

Section Summary: Progenesa PCA3 Assay

The analytic validity of Progenesa has been established. At least 4 systematic reviews, including a health technology assessment, have been reported and included many primary studies. Studies of the PCA3 score as a diagnostic test for prostate cancer have reported sensitivities and specificities in the moderate range. In general, these studies are preliminary and report on clinical performance characteristics in different populations and with various assay cutoff values, reflecting the lack of standardization in performance and interpretation of PCA3 test results. Cutoffs for recommending repeat biopsy with the Progenesa test have been suggested by the manufacturer and were used in a validation study for FDA approval. The clinical utility of the PCA3 test is uncertain because there is no evidence that its use can change management in ways that improve outcomes.

Gene Hypermethylation and ConfirmMDx

Epigenetic changes—chromatin protein modifications that do not involve changes to the underlying DNA sequence but can change gene expression—have been identified in specific genes. An extensive literature has reported significant associations between epigenetic DNA modifications and prostate cancer. Several investigators have evaluated detection of hypermethylation products in biologic fluids for early detection of prostate cancer. Promoters of 3 genes (adenomatous polyposis coli [*APC*], *GSTP1*, *RARβ2*) were identified early as potentially involved in prostate carcinogenesis. *GSTP1* is the most widely studied methylation marker for prostate cancer, usually as a diagnostic application. Studies in the late 2000s of *GSTP1* hypermethylation using tissue samples reported conflicting results. Sunami et al (2009) assayed blood from 40 healthy individuals and 83 men with prostate cancer using a 3-gene cohort (*GSTP1*, *RASSF1*, *RARβ2*) and demonstrated sensitivity of 28% for cancer patients. Trock et al (2012) conducted a small (86-patient) diagnostic exploratory cohort study and showed that hypermethylation of *APC* was associated with high sensitivity and specificity for cancer on repeat biopsy.

In a 2012 meta-analysis by Van Neste et al, 30 peer-reviewed studies of hypermethylation of *GSTP1* and other genes in prostate tissue were evaluated. The pooled estimates of sensitivity for *GSTP1* to distinguish prostate cancer from normal in biopsies (328 cases, 263 controls) was 82%, with 95% specificity, 95% NPV, and 85% PPV. The combination of *GSTP1*, *APC*, and *RARβ* had a sensitivity of 95%, specificity of 95%, NPV of 99%, and PPV of 95%. Reviewers suggested that a valuable first step in diagnostic use might be to test for methylated genes to select patients undergoing prostate biopsy who might not require repeat biopsy.

Following the 2012 meta-analysis, several studies reported on associations between DNA hypermethylation at various gene loci (*RASSF1A*, *APC*, *GSTP1*, *PTGS2*, *RARβ*, *TIG1*, *AOX1*, *C1orf114*, *GAS6*, *HAPLN3*, *KLF8*, *MOB3B*) and prostate cancer. In contrast, Kachakova et al (2013) found that *HIST1H4K* hypermethylation was more likely due to aging than to prostate carcinogenesis.

ConfirmMDx (MDxHealth) is a commercially available test for gene methylation intended to distinguish true- from false-negative prostate biopsies to avoid the need for repeat biopsy in cases of a true negative and to

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

identify men who may need a repeat biopsy. The test measures methylation of the genes *GSTP1*, *APC*, and *RASSF1*.

Analytic Validity

Goessl et al (2002) confirmed in 26 patients that neoplastic transformation could be identified in washings of prostate biopsies by *GSTP1* promoter hypermethylation using methylation-specific polymerase chain reaction (PCR). Chu et al (2002) described a protocol for real-time, quantitative, methylation-sensitive PCR for detecting the methylation change in the 5' regulatory sequence flanking the *GSTP1* gene that was more sensitive than conventional nested PCR (DNA test limitations were 0.048 ng and 0.64 ng, respectively). Mehrotra et al (2008) confirmed that a field effect was detectable for *APC*, *RARβ2*, and *RASSF1A* up to 3 mm from the malignant core. Van Neste et al (2012) described a study evaluating a multiplex assay consisting of 3 genes: *GSTP1*, *APC*, and *RASSF1*. Thirty cancer-positive tissue samples and 12 cancer-free controls were analyzed with 4 singleplex vs 1 multiplex assay. A control gene (*ACTB*) was used to estimate DNA quantity and quality in 2 replicates. The ratio of *ACTB* copies ranged from 0.73 to 1.17 (outlier removed) for the multiplex assay, with a median ratio of 1.0. *ACTB* copy numbers were higher for the multiplex assay than for a singleplex assay (median, 1.5-fold copy increase). A linear regression model yielded amplification factors of 1.57, 1.19, 4.13, and 1.25 for the *ACTB*, *GSTP1*, *APC*, and *RASSF1* assays, respectively, with consistently high R^2 values (>0.90). Biopsies consisting of 10, 20, and 40 μm from formalin-fixed, paraffin-embedded tissue blocks from the minimization cohort were tested and compared (outliers were removed). The effect of the original sample volume on the relative DNA yield was minor, indicating that samples as small as 20 μm can be used to detect methylation. Older samples showed lower relative DNA yields ($p < 0.001$), indicating that the age of formalin-fixed, paraffin-embedded samples does have a negative impact on DNA quality and quantity. Other measures of analytic validity were not found in the literature or the ConfirmMDx website. The laboratory that performs the analyses for ConfirmMDx is certified under the Clinical Laboratory Improvement Amendments.

Clinical Validity

Two blinded multicenter validation studies of the ConfirmMDx test have been performed. Partin et al (2014) reported on results of the DOCUMENT study; it evaluated archived, cancer-negative prostate biopsy core tissue samples from 350 men from 5 U.S. urology centers. All patients underwent repeat biopsy within 24 months. Men with 2 consecutive negative biopsies were classified as controls and men with a negative biopsy followed by a positive biopsy were classified as cases. Thirty (9%) men were excluded from analysis because of noneligibility (n=2), insufficient DNA (n=1), insufficient biopsy cores (n=23), or detection of adenocarcinoma in the first biopsy based on central pathology review (n=4); 320 men were included in analysis (92 cases, 228 controls). Median age was 62 years (range, not given). Median PSA level was 5.3 ng/mL; 23% of men had PSA levels less than 4 ng/mL and 10% had a PSA level of 10 ng/mL or higher. Sixty percent of men had a normal DRE. Forty-two (13%) of the men were black, 232 (73%) were white, and 13 (4%) were Asian. The ConfirmMDx test, performed on the first biopsy, resulted in a NPV of 88% (95% CI, 85% to 91%), sensitivity of 62% (95% CI, 51% to 72%), and specificity of 64% (95% CI, 57% to 70%). The study was not powered to determine accurately the performance characteristics in a subgroup of black patients, but the estimated sensitivity was 77% (95% CI, 46% to 95%), specificity was 66% (95% CI,

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

46% to 82%), and NPV was 93% (85% CI, 82% to 97%). Multivariate analysis of potential predictors of cancer on repeat biopsy, corrected for age, PSA, DRE, first biopsy histopathology characteristics, and race, showed that the ConfirmMDx test was the most significant independent predictor of patient outcome (OR=2.69; 95% CI, 1.60 to 4.51).

The MATLOC study, reported by Stewart et al (2013), tested archived cancer-negative prostate biopsy needle core tissue samples from 498 men from the U.K. and Belgium. Patients underwent repeat biopsy within 30 months; cases had a positive second biopsy while controls had a negative second biopsy. A total of 483 men were included in the analysis (87 cases, 396 controls). The median PSA level was 5.9 ng/mL; 21% of men had PSA levels less than 4 ng/mL and 18% had PSA levels of 10 ng/mL or higher. Seventy-three percent of men had benign DRE. The ConfirmMDx test, performed on the first biopsy, resulted in a NPV of 90% (95% CI, 87% to 93%), sensitivity of 68% (95% CI, 57% to 77%), and specificity of 64% (95% CI, 59% to 69%). Multivariate analysis of potential predictors of cancer on repeat biopsy, corrected for patient age, PSA, DRE, and first biopsy histopathology characteristics, showed that the ConfirmMDx test was the most significant independent predictor of patient outcome (OR=3.17; 95% CI, 1.81 to 5.53).

In 2016, Van Neste et al reported on results of combined data from the DOCUMENT and MATLOC studies to investigate whether DNA methylation intensities were associated with high-grade (Gleason score, ≥ 7) prostate cancer. DNA methylation was the most significant and important predictor of high-grade cancer, resulting in an NPV of 96% (precision not reported).

Clinical Utility

In 2014, Wojno et al reported on a field observation study in which practicing urologists at 5 centers used the ConfirmMDx test to evaluate at least 40 men with previous cancer-negative biopsies who were considered at risk for prostate cancer. Centers reported whether patients who had a negative test assay result had undergone a repeat biopsy at the time of the analysis. Median patient follow-up after the assay results were received was 9 months. A total of 138 patients were included in the analysis. The median PSA level was 4.7 ng/mL. Repeat biopsies had been performed in 6 (4.3%) of the 138 men with a negative ConfirmMDx test, in which no cancer was identified.

In 2013, Aubry et al analyzed the expected reduction in biopsies associated with ConfirmMDx use. Using the MATLOC estimates of performance characteristics for ConfirmMDx, the authors estimated that 1106 biopsies per 1 million people would be avoided. The study did not include a decision analysis comparing the tradeoff in reduction in biopsies and missed cancers.

MDxHealth completed enrollment into the PASCUAL trial in April 2015. The PASCUAL trial is described as an observational trial of ConfirmMDx to evaluate the impact of the test (clinical utility) on physician decisions for repeat biopsy. Results have not yet been published.

No studies were found that directly show the effects of using ConfirmMDx results on clinical outcomes. Given the lack of direct evidence of utility, a chain of evidence would be needed to demonstrate clinical

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

utility. The ConfirmMDx test is associated with a diagnosis of prostate cancer and aggressive prostate cancer. The validity studies of the ConfirmMDx test included men in the intended use population but did not compare performance characteristics with clinical examination plus percent free PSA. One survey of urologists who had previously used the ConfirmMDx test found that most ConfirmMDx negative patients did not have a biopsy. Prospective data on utility should be available after completion of PASCUAL. No data are available on the longer term clinical outcomes of the men who did not have biopsy based on ConfirmMDx results. The chain of evidence is incomplete.

Section Summary: Gene Hypermethylation and ConfirmMDx

Two clinical validation studies have reported on the clinical validity of the ConfirmMDx score in the intended use population. The studies did not provide estimates of validity compared with a standard clinical examination with percent free PSA. No data are available on the long-term clinical outcomes or clinical utility of the test. The indirect chain of evidence is incomplete due to the limitations of evidence on the comparative clinical validity and utility.

TMPRSS Fusion Genes and Mi-Prostate

TMPRSS2 is an androgen-regulated transmembrane serine protease that is preferentially expressed in normal prostate tissue. In prostate cancer, it may be fused to an E26 transformation-specific (ETS) family transcription factor (*ERG*, *ETV1*, *ETV4*, *ETV5*), which modulates transcription of target genes involved in cell growth, transformation, and apoptosis. The result of gene fusion with an ETS transcription gene is that the androgen-responsive promoter of *TMPRSS2* upregulates expression of the ETS gene, suggesting a mechanism for neoplastic transformation. Fusion genes may be detected in tissue, serum, or urine.

TMPRSS2-ERG gene rearrangements have been reported in 50% or more of primary prostate cancer samples. Although *ERG* appears to be the most common ETS family transcription factor involved in the development of fusion genes, not all are associated with *TMPRSS2*. About 6% of observed rearrangements are seen with *SLC45A3*, and about 5% appear to involve other types or rearrangement.

In 2014, Yao et al published a systematic review with meta-analysis of *TMPRSS2-ERG* for the detection of prostate cancer. Literature was searched through July 2013, and 32 articles were included. Pooled sensitivity, specificity, positive likelihood ratio, and negative likelihood ratio were 47% (95% CI, 46% to 49%), 93% (95% CI, 92% to 94%), 8.9 (95% CI, 5.7 to 14.1), and 0.49 (95% CI, 0.43 to 0.55), respectively. Statistical heterogeneity was high ($I^2 > 85\%$). It was unclear whether studies in screening populations were pooled with enriched patient samples (eg, elevated PSA and/or negative biopsy). There also was variability in the type of tissue samples analyzed (urine, prostatic secretions, biopsy, surgical specimens); the type of *TMPRSS2-ERG* assays used (fluorescence in situ hybridization, immunohistochemistry, real-time reverse transcriptase PCR, transcription-mediated amplification); and in *TMPRSS2-ERG* threshold cutoff values.

The Mi-Prostate (MiPS) is a test using the *TMPRSS2-ERG* gene to produce a risk probability for detection of prostate cancer and aggressive prostate cancer by standard biopsy. The probability score is calculated with logistic regression models that incorporate serum PSA, or the PCPT version 1.0, and urine *T2-ERG*

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

and PCA3 scores. The test was developed by, and is only available from, the University of Michigan MLabs, and may be used to decide about monitoring PSA levels or pursuing a prostate biopsy.

Analytic Validity

The MiPS test uses results from the Progensa PCA3 test that demonstrated analytic validity in the FDA submission. The amounts of urine *TMPRSS2-ERG* are determined using transcription-mediated amplification assays. No peer-reviewed, full-length publications describing the analytic validity of the *TMPRSS2-ERG* assays were identified.

Clinical Validity

Tomlins et al (2011) developed a transcription-mediated amplification assay to measure *TMPRSS2-ERG* fusion transcripts in parallel with PCA3. Combining results from the *TMPRSS2-ERG* and PCA tests and incorporating them into the multivariate PCPT risk calculator appeared to improve identification of patients with clinically significant cancer using Epstein criteria and high-grade cancer on biopsy. Although the study was large (1312 men at multiple centers), it was confounded by assay modifications during the study and by use of cross-validation rather than independent validation, using independent training and testing sets.

In 2013, this same group evaluated 45 men using a multivariable algorithm that included serum PSA plus urine *TMPSS2-ERG* and PCA3 from a post-DRE sample. Samples were collected before prostate biopsy at 2 centers. For cancer prediction, sensitivity and specificity were 80% and 90%, respectively. AUC was 0.88.

In 2016, Tomlins et al published results of a validation study of the MiPS score in 1244 prospectively collected, post-DRE urine samples from 7 U.S. clinics. A total of 1225 of the specimens had sufficient materials for both *TMPSS2-ERG* and PCA3 analysis and were included. Eighty percent of patients were presenting for initial biopsy. Seventy-three percent were white; the percentage of African Americans was not given. Approximately 25% of the men were older than 70. Twenty-three percent had an abnormal DRE, and the median PSA level was 4.7 ng/mL. The AUCs for predicting any cancer using PSA alone, PCPT risk calculator alone, and the MiPS score alone were 0.59, 0.64, and 0.76, respectively (CIs not given, $p < 0.001$ for MiPS vs PCPT). The AUCs for predicting high-grade cancer were 0.65, 0.71, and 0.78, respectively ($p < 0.001$ for MiPS vs PCPT). A MiPS score threshold for recommending biopsy has not been provided, and so sensitivity and NPV were not calculated.

Clinical Utility

Tomlins et al (2016) also used decision-curve analysis to estimate the number of biopsies that would have been performed and cancers that would have been missed using a MiPS risk cutoff for biopsy in their cohort. Compared with a biopsy-all strategy, using a MiPS cutoff for aggressive cancer of 15% would have avoided 36% of biopsies while missing 7.0% of any prostate cancer and 1.6% of high-grade prostate cancer diagnoses. Using the PCPT risk calculator cutoff of 15% for aggressive cancer would have avoided 68% of biopsies while missing 25% of any cancer and 8% of high-grade cancer.

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

No studies were found that directly show the effects of using MiPS results on clinical outcomes. Given the lack of direct evidence of utility, a chain of evidence would be needed to demonstrate clinical utility. The MiPS test is associated with a diagnosis of prostate cancer and aggressive prostate cancer. The clinical validity study of the MiPS test included men with relevant PSA levels but also included men with positive DRE who would not likely forgo biopsy. The clinical validation study included comparison of performance characteristics with standard risk calculators; comparison with percent free PSA was not provided. Confirmation of performance characteristics is needed. No prospective data are available on using the MiPS score for decision making. No data are available on the longer term clinical outcomes of the men who did not have biopsy based on MiPS results. The chain is incomplete.

Section Summary: *TMPRSS Fusion Genes and Mi-Prostate*

Concomitant detection of *TMPRSS2-ERG* and *PCA3* may more accurately identify men with prostate cancer. However, current evidence is insufficient to support its use. Estimated accuracy varies across available studies. The MiPS test has preliminary data suggesting improved clinical validity compared with the PCPT risk calculator in a validation study, but independent confirmation of clinical validity and comparison with percent free PSA is needed. Data on analytic validity and clinical utility are lacking.

Prostate Core Mitomics Test

The Prostate Core Mitomics Test (PCMT; Mitomics; formerly Genesis Genomics) is a proprietary test intended to determine whether a patient has prostate cancer, despite a negative prostate biopsy, by analyzing deletions in mitochondrial DNA by PCR to detect “tumor field effect.” The test is performed on the initial negative prostate biopsy tissue. According to the company website, a negative PCMT result confirms the result of the negative biopsy (ie, the patient does not have prostate cancer) and that the patient can avoid a second biopsy, but a positive PCMT means that the patient is at high risk of undiagnosed prostate cancer. The website also states that physicians should consider using PCMT for patients who have a negative initial biopsy but continue to have elevated PSA, rising PSA, irregular DRE, atypical small acinar proliferation, high-grade prostatic intraepithelial neoplasia, or inconclusive biopsy.

Analytic Validity

No peer-reviewed, full-length publications on the analytic validity of the commercially available PCMT test were identified.

Clinical Validity

A 2006 study retrospectively analyzed mitochondrial DNA variants from 3 tissue types from 24 prostatectomy specimens: prostate cancer, adjacent benign tissue, and benign tissue distant to the tumor (defined as tissue from a nondiseased lobe or at least 10-cell diameters from the tumor if in the same lobe). Prostate needle biopsy tissue (from 12 individuals referred for biopsy) that were histologically benign were used as controls. Results from the prostatectomy tissue analysis showed that 16 (66.7%) of 24 had variants in all 3 tissue types, 22 (91.7%) of 24 had variants in malignant samples, 19 (79.2%) of 24 in adjacent benign samples, and 22 of 24 in distant benign glands. Overall, 273 somatic variants were observed in this sample set. In the control group, 7 (58.3%) patients had between 1 and 5 genetic alterations, mainly in

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

noncoding regions. The authors concluded that the variants found in the malignant group vs the control group differed significantly and that mitochondrial DNA variants are an indicator of malignant transformation in prostate tissue.

In 2008, Maki et al reported on the discovery and characterization of a 3.4-kilobase mitochondrial genome deletion and its association with prostate cancer. A pilot study analyzed 38 benign biopsy specimens from 22 patients, 41 malignant biopsy specimens from 24 patients, and 29 proximal to malignant (PTM) biopsy specimens from 22 patients. All patients with malignant biopsies had a subsequent prostatectomy, and the diagnosis of cancer was confirmed. The PTM biopsy samples were negative for cancer and were from the cohort that underwent prostatectomy. A confirmation study used 98 benign biopsy specimens from 22 patients, 75 malignant biopsy specimens from 65 patients, and 123 PTM biopsy specimens from 96 patients. In the confirmation study, patients had to have at least 2 successive negative biopsies; the first negative biopsy was used for analyses. For both the pilot and confirmation studies, samples for analysis were selected based on a review of pathology reports. The levels of the variation were measured by quantitative PCR and, based on PCR cycle threshold, data were used to calculate a score for each biopsy sample. In the pilot study, the scores were statistically significant between benign and malignant ($p < 0.000$) and benign and proximal ($p < 0.003$) samples. The PTM samples closely resembled the malignant sample, with no statistically significant resolution between the scores ($p < 0.833$), to which the authors attributed to a field cancerization phenomenon. Results from the larger confirmation study were similar. Compared with histopathologic examination of the benign and malignant samples, the sensitivity and specificity were 80% and 71%, respectively, and the AUROC was 0.83 for the deletion. A blinded, external validation study showed a sensitivity and specificity of 83% and 79% and the AUROC of 0.87.

In 2010, Robinson et al assessed the clinical value of the 3.4-kilobase deletion described in the Maki study in predicting rebiopsy outcomes. Levels of the deletion were measured by quantitative PCR in prostate biopsies negative for cancer from 101 patients who underwent repeat biopsy within 1 year and had known outcomes. Of the 101 first biopsies, the diagnosis was normal in 8, atypical and/or had prostatic intraepithelial neoplasia in 50, and hyperplasia or inflammation in 43. The clinical performance of the deletion was calculated with the use of an empirically established cycle threshold cutoff, the lowest cycle threshold as diagnostic of prostate cancer, and the histopathologic diagnosis on second biopsy. Final data were based on 94 patients, who on second biopsy had 20 malignant and 74 benign diagnoses. The cycle cutoff gave a sensitivity and specificity of 84% and 54%, respectively, with an AUROC of 0.75. NPV was 91%.

Clinical Utility

No peer-reviewed, full-length publications on the clinical utility of the commercially available PCMT test was identified.

Section Summary: Prostate Core Mitomics Test

The PCMT test has preliminary data on performance characteristics in a small validation study, but independent confirmation of clinical validity is needed. The studies did not provide estimates of validity

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

compared with a standard clinical examination. No data are available on the long-term clinical outcomes. Data on analytic validity and clinical utility are lacking.

Candidate Gene Panels and SNV Testing

Because no single gene marker that is both highly sensitive and highly specific for diagnosing prostate cancer has been found, particularly in men already known to have elevated PSA levels, some investigators are combining several markers into a single diagnostic panel. Although promising in concept, only single studies of various panels have been published, and none apparently is offered as a clinical service.

SNVs occur when a single nucleotide is replaced with another, and they are the most common type of genetic variation in humans. They occur normally throughout the genome and can act as biologic markers for disease association. Genome-wide association studies have identified correlations between prostate cancer risk and specific SNVs. However, it is widely accepted that individually, SNV-associated disease risk is low and of no value in screening for disease, although multiple SNVs in combination may account for a higher proportion of prostate cancer. Investigators have begun to explore the use of algorithms incorporating information from multiple SNVs to increase the clinical value of testing.

Ma et al (2014) examined various algorithms for cancer diagnosis and prognosis using urine and plasma levels of multiple genes, including *PCA3*, *PSA*, *TMPRSS2*, and *ERG*. One algorithm distinguished prostate cancer from benign prostatic hypertrophy with an AUC of 0.78. Another algorithm distinguished men with a Gleason score 7 or higher for men with a Gleason score less than 7 (AUC=0.88). Combination of these 2 algorithms into a scoring system predicted the presence of a Gleason score 7 or higher in 75% of men. Qu et al (2013) reported on preliminary results of a 3-gene panel (androgen receptor [*AR*], *PTEN*, *TMPRSS2-ERG*) analyzed by fluorescence in situ hybridization. Thirty-one percent of 110 archived primary tumor samples and 97 metastatic tumor samples from a separate cohort of patients were analyzable. Chromosomal abnormalities were detected in 53% of primary prostate cancers and in 87% of metastatic tumors ($p < 0.001$).

In 2015, Leyten et al reported on the development of a gene panel using specimens from 133 patients that included 3 urinary biomarkers (*HOXC6*, *TDRD1*, *DLX1*). When the gene panel was used with PSA, the combined AUC for predicting high-grade prostate cancer was 0.81 (95% CI, 0.75 to 0.86), which was higher than the concurrently measured Progenesa AUC of 0.68 (95% CI, 0.62 to 0.75). Xiao et al (2016) reported the development of an 8-gene panel (*PMP22*, *HPN*, *LMTK2*, *FN1*, *EZH2*, *GOLM1*, *PCA3*, *GSTP1*) that was able to distinguish high-grade prostate cancer from indolent prostate cancer with a sensitivity of 93% (95% CI, 88% to 97%), a specificity of 70% (95% CI, 36% to 104%), a PPV of 98% (95% CI, 95% to 100%), and an NPV of 61% (95% CI, 25% to 97%) in a specimen cohort of 158 men.

A 2012 Agency for Healthcare Research and Quality report on multigene panels in prostate cancer risk assessment reviewed the literature on SNV panel tests for assessing risk of prostate cancer. All studies included in the review had poor discriminative ability for predicting the risk of prostate cancer, had a moderate risk of bias, and none of the panels had been evaluated in routine clinical settings. Reviewers

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

concluded that the evidence on the available SNV panels did not permit a meaningful assessment of analytic validity, that the limited evidence on clinical validity was insufficient to conclude that SNV panels would perform adequately as a screening test, and that there was no evidence on the clinical utility of current panels.

Kader et al (2012) evaluated a panel of 33 prostate cancer-associated SNVs identified from genome-wide association studies in 1654 men. Genetic score was a significant ($p < 0.001$) independent predictor of prostate cancer (OR=1.72; 95% CI, 1.44 to 2.09) after adjusting for clinical variables and family history. Addition of genetic markers to the classification of prostate cancer risk resulted in 33% of men by reclassified into a different risk quartile. Approximately half of these ($n=267$) were downgraded to a lower risk quartile, and the other half ($n=265$) were upgraded into a higher risk quartile. The net reclassification benefit was 10% ($p=0.002$). The authors concluded that, with the additional information of genetic score, the same number of cancers could be detected but with 15% fewer biopsies.

In a 2010 review by Ioannidis et al, 27 gene variants across a large range of chromosomal locations were identified that increased risk for prostate cancer, although, in all cases, the observed incremental risk was modest (OR \leq 1.36).

Lindstrom et al (2011), in a study of 10,501 cases of prostate cancer and 10,831 controls, identified 36 SNVs showing association with prostate cancer risk, including two (rs2735893, rs266849) that showed differential association with Gleason score. Per allele odds ranged from 1.07 to 1.44.

Ishak and Giri (2011) reviewed 11 replication studies involving 30 SNVs (19 in men of African descent, 10 in men with familial prostate cancer). Odds ratios were positively associated with prostate cancer, although the magnitude of association was small (range, 1.11-2.63).

Section Summary: Candidate Gene Panels and SNV Testing

Numerous studies have demonstrated the association of many gene panels and SNVs with prostate cancer. These studies, in early stages of development, have shown a modest degree of association with future risk for prostate cancer. The clinical utility of these tests is uncertain; there is no evidence that information obtained from gene panels or SNV testing can be used to change clinical management in ways that will improve outcomes.

SUMMARY OF EVIDENCE

For individuals who are being considered for an initial prostate biopsy or a repeat biopsy who receive testing for genetic and protein biomarkers of prostate cancer, the evidence includes systematic reviews and meta-analyses and primarily observational studies. Relevant outcomes are overall survival, disease-specific survival, test accuracy and validity, other test performance measures, resource utilization, and quality of life. The evidence supporting clinical utility varies by test but has not been directly shown for any biomarker test. In general, the performance of biomarker testing for predicting biopsy referrals compared with clinical examination, including the ratio of free or unbound PSA to total PSA, is lacking. Procedures for referrals for biopsy based on clinical examination vary, making it difficult to quantify performance characteristics for this

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

comparator. There is also considerable variability in biopsy referral practices based on clinical examination alone, and many biomarker tests do not have standardized cutoffs to recommend a biopsy. Therefore, to determine whether the tests improve the net health outcome, prospective, comparative data are needed on how test results are expected to be used vs how they are being used in practice, because of information about the associated effects on outcomes. Many test validation populations have included men with a positive digital rectal exam, PSA level outside of the gray zone (between 3 or 4 ng/mL and 10 ng/mL), or older men for whom the information from PSA test results are less likely to be informative. African American men have a high burden of morbidity and mortality, but have not been well represented in these study populations. It is unclear how to monitor men with low biomarker risk scores who continue to have symptoms or high or rising PSA levels. Comparative studies of the many biomarkers are lacking, and it is unclear how to use the tests in practice, particularly when test results are contradictory. The evidence is insufficient to determine the effects of the technology on health outcomes.

References

1. Blue Cross and Blue Shield Association, Medical Policy Reference Manual, "Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer", 2.04.33, 11:2017.
2. Siegel RL, Miller KD, Jemal A. Cancer Statistics, 2017. *CA Cancer J Clin*. Jan 2017;67(1):7-30. PMID 28055103
3. Howlader N, Noone AM, Krapcho M, et al. SEER Cancer Statistics Review, 1975-2014. Bethesda, MD: National Cancer Institute; 2017.
4. Ries LAG, Melbert D, Krapcho M, et al. SEER Cancer Statistics Review, 1975-2004. 2007; http://seer.cancer.gov/csr/1975_2004/. Accessed September 6, 2016.
5. Odedina FT, Akinremi TO, Chinegwundoh F, et al. Prostate cancer disparities in Black men of African descent: a comparative literature review of prostate cancer burden among Black men in the United States, Caribbean, United Kingdom, and West Africa. *Infect Agent Cancer*. 2009;4 Suppl 1:S2. PMID 19208207
6. Bell KJ, Del Mar C, Wright G, et al. Prevalence of incidental prostate cancer: A systematic review of autopsy studies. *Int J Cancer*. Oct 1 2015;137(7):1749-1757. PMID 25821151
7. Gleason DF. Classification of prostatic carcinomas. *Cancer Chemother Rep*. Mar 1966;50(3):125-128. PMID 5948714
8. Wright JL, Salinas CA, Lin DW, et al. Prostate cancer specific mortality and Gleason 7 disease differences in prostate cancer outcomes between cases with Gleason 4 + 3 and Gleason 3 + 4 tumors in a population based cohort. *J Urol*. Dec 2009;182(6):2702-2707. PMID 19836772
9. Hoogendam A, Buntinx F, de Vet HC. The diagnostic value of digital rectal examination in primary care screening for prostate cancer: a meta-analysis. *Fam Pract*. Dec 1999;16(6):621-626. PMID 10625141
10. Gosselaar C, Roobol MJ, Roemeling S, et al. The role of the digital rectal examination in subsequent screening visits in the European Randomized Study of Screening for Prostate Cancer (ERSPC), Rotterdam. *Eur Urol*. Sep 2008;54(3):581-588. PMID 18423977
11. Thompson IM, Pauler DK, Goodman PJ, et al. Prevalence of prostate cancer among men with a prostate-specific antigen level < or =4.0 ng per milliliter. *N Engl J Med*. May 27 2004;350(22):2239-2246. PMID 15163773
12. Catalona WJ, Smith DS, Ratliff TL, et al. Measurement of prostate-specific antigen in serum as a screening test for prostate cancer. *N Engl J Med*. Apr 25 1991;324(17):1156-1161. PMID 1707140
13. Aus G, Bergdahl S, Lodding P, et al. Prostate cancer screening decreases the absolute risk of being diagnosed with advanced prostate cancer--results from a prospective, population-based randomized controlled trial. *Eur Urol*. Mar 2007;51(3):659-664. PMID 16934392
14. Buzzoni C, Auvinen A, Roobol MJ, et al. Metastatic prostate cancer incidence and prostate-specific antigen testing: new insights from the European randomized study of screening for prostate cancer. *Eur Urol*. Nov 2015;68(5):885-890. PMID 25791513
15. Arnsrud Godtman R, Holmberg E, Lilja H, et al. Opportunistic testing versus organized prostate-specific antigen screening: outcome after 18 years in the Goteborg randomized population-based prostate cancer screening trial. *Eur Urol*. Sep 2015;68(3):354-360. PMID 25556937

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

16. Hugosson J, Carlsson S, Aus G, et al. Mortality results from the Goteborg randomised population-based prostate-cancer screening trial. *Lancet Oncol.* Aug 2010;11(8):725-732. PMID 20598634
17. Schroder FH, Hugosson J, Roobol MJ, et al. Screening and prostate-cancer mortality in a randomized European study. *N Engl J Med.* Mar 26 2009;360(13):1320-1328. PMID 19297566
18. Wolf AM, Wender RC, Etzioni RB, et al. American Cancer Society guideline for the early detection of prostate cancer: update 2010. *CA Cancer J Clin.* Mar-Apr 2010;60(2):70-98. PMID 20200110
19. Moyer VA, U. S. Preventive Services Task Force. Screening for prostate cancer: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med.* Jul 17 2012;157(2):120-134. PMID 22801674
20. Greene KL, Albertsen PC, Babaian RJ, et al. Prostate specific antigen best practice statement: 2009 update. *J Urol.* Nov 2009;182(5):2232-2241. PMID 19781717
21. Hamdy FC, Donovan JL, Lane JA, et al. 10-year outcomes after monitoring, surgery, or radiotherapy for localized prostate cancer. *N Engl J Med.* Sep 14 2016. PMID 27626136
22. Rosario DJ, Lane JA, Metcalfe C, et al. Short term outcomes of prostate biopsy in men tested for cancer by prostate specific antigen: prospective evaluation within ProtecT study. *BMJ.* 2012;344:d7894. PMID 22232535
23. Gonzalez CM, Averch T, Boyd LA, et al. AUA/SUNA white paper on the incidence, prevention and treatment of complications related to prostate needle biopsy. 2012; https://www.auanet.org/common/pdf/practices-resources/quality/patient_safety/Prostate-Needle-Biopsy-White-Paper.pdf. Accessed September 6, 2016.
24. Lavallee LT, Binette A, Witiuk K, et al. Reducing the harm of prostate cancer screening: repeated prostate-specific antigen testing. *Mayo Clin Proc.* Jan 2016;91(1):17-22. PMID 26688045
25. Partin AW, Brawer MK, Subong EN, et al. Prospective evaluation of percent free-PSA and complexed-PSA for early detection of prostate cancer. *Prostate Cancer Prostatic Dis.* Jun 1998;1(4):197-203. PMID 12496895
26. Thompson IM, Ankerst DP, Chi C, et al. Assessing prostate cancer risk: results from the Prostate Cancer Prevention Trial. *J Natl Cancer Inst.* Apr 19 2006;98(8):529-534. PMID 16622122
27. van Vugt HA, Roobol MJ, Kranse R, et al. Prediction of prostate cancer in unscreened men: external validation of a risk calculator. *Eur J Cancer.* Apr 2011;47(6):903-909. PMID 21163642
28. Rosenkrantz AB, Verma S, Choyke P, et al. Prostate Magnetic Resonance Imaging and Magnetic Resonance Imaging Targeted Biopsy in Patients with a Prior Negative Biopsy: A Consensus Statement of the American Urological Association and the Society of Abdominal Radiology's Prostate Cancer Disease Focused Panel. *J Urol.* Jun 16 2016. PMID 27320841
29. Djavan B, Waldert M, Zlotta A, et al. Safety and morbidity of first and repeat transrectal ultrasound guided prostate needle biopsies: results of a prospective European prostate cancer detection study. *J Urol.* Sep 2001;166(3):856-860. PMID 11490233
30. Lujan M, Paez A, Santonja C, et al. Prostate cancer detection and tumor characteristics in men with multiple biopsy sessions. *Prostate Cancer Prostatic Dis.* 2004;7(3):238-242. PMID 15289810
31. Blue Cross and Blue Shield Technology Evaluation Center (TEC). Special report: recent developments in prostate cancer genetics and genetic testing. *TEC Assessments.* 2008;Volume 23:Tab 7. PMID
32. 4Kscore(R). <http://4kscore.com/>. Accessed September 7, 2016.
33. Vaisanen V, Peltola MT, Lilja H, et al. Intact free prostate-specific antigen and free and total human glandular kallikrein 2. Elimination of assay interference by enzymatic digestion of antibodies to F(ab)'2 fragments. *Anal Chem.* Nov 15 2006;78(22):7809-7815. PMID 17105175
34. Borque-Fernando A, Esteban-Escano LM, Rubio-Briones J, et al. A preliminary study of the ability of the 4Kscore test, the Prostate Cancer Prevention Trial-Risk Calculator and the European Research Screening Prostate-Risk Calculator for predicting high-grade prostate cancer. *Actas Urol Esp.* Apr 2016;40(3):155-163. PMID 26598800
35. Konety B, Zappala SM, Parekh DJ, et al. The 4Kscore(R) test reduces prostate biopsy rates in community and academic urology practices. *Rev Urol.* 2015;17(4):231-240. PMID 26839521
36. Nordstrom T, Vickers A, Assel M, et al. Comparison between the four-kallikrein panel and Prostate Health Index for predicting prostate cancer. *Eur Urol.* Jul 2015;68(1):139-146. PMID 25151013
37. Stattin P, Vickers AJ, Sjoberg DD, et al. Improving the specificity of screening for lethal prostate cancer using prostate-specific antigen and a panel of kallikrein markers: a nested case-control study. *Eur Urol.* Aug 2015;68(2):207-213. PMID 25682340
38. Carlsson S, Maschino A, Schroder F, et al. Predictive value of four kallikrein markers for pathologically insignificant compared with aggressive prostate cancer in radical prostatectomy specimens: results from the European Randomized Study of Screening for Prostate Cancer section Rotterdam. *Eur Urol.* Nov 2013;64(5):693-699. PMID 23683475

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

39. Vickers A, Cronin A, Roobol M, et al. Reducing unnecessary biopsy during prostate cancer screening using a four-kallikrein panel: an independent replication. *J Clin Oncol.* May 20 2010;28(15):2493-2498. PMID 20421547
40. Gupta A, Roobol MJ, Savage CJ, et al. A four-kallikrein panel for the prediction of repeat prostate biopsy: data from the European Randomized Study of Prostate Cancer screening in Rotterdam, Netherlands. *Br J Cancer.* Aug 24 2010;103(5):708-714. PMID 20664589
41. Benchikh A, Savage C, Cronin A, et al. A panel of kallikrein markers can predict outcome of prostate biopsy following clinical work-up: an independent validation study from the European Randomized Study of Prostate Cancer screening, France. *BMC Cancer.* 2010;10:635. PMID 21092177
42. Vickers AJ, Cronin AM, Roobol MJ, et al. A four-kallikrein panel predicts prostate cancer in men with recent screening: data from the European Randomized Study of Screening for Prostate Cancer, Rotterdam. *Clin Cancer Res.* Jun 15 2010;16(12):3232-3239. PMID 20400522
43. Vickers AJ, Gupta A, Savage CJ, et al. A panel of kallikrein marker predicts prostate cancer in a large, population-based cohort followed for 15 years without screening. *Cancer Epidemiol Biomarkers Prev.* Feb 2011;20(2):255-261. PMID 21148123
44. Vickers AJ, Cronin AM, Aus G, et al. A panel of kallikrein markers can reduce unnecessary biopsy for prostate cancer: data from the European Randomized Study of Prostate Cancer Screening in Goteborg, Sweden. *BMC Med.* 2008;6:19. PMID 18611265
45. Vickers AJ, Cronin AM, Aus G, et al. Impact of recent screening on predicting the outcome of prostate cancer biopsy in men with elevated prostate-specific antigen: data from the European Randomized Study of Prostate Cancer Screening in Gothenburg, Sweden. *Cancer.* Jun 1 2010;116(11):2612-2620. PMID 20336781
46. Bryant RJ, Sjoberg DD, Vickers AJ, et al. Predicting high-grade cancer at ten-core prostate biopsy using four kallikrein markers measured in blood in the ProtecT study. *J Natl Cancer Inst.* Jul 2015;107(7). PMID 25863334
47. Parekh DJ, Punnen S, Sjoberg DD, et al. A multi-institutional prospective trial in the USA confirms that the 4Kscore accurately identifies men with high-grade prostate cancer. *Eur Urol.* Sep 2015;68(3):464-470. PMID 25454615
48. phi: Prostate Health Index. <http://prostatehealthindex.us/>. Accessed September 9, 2016.
49. Food and Drug Administration. Summary of Safety and Effectiveness Data. PMA P090026. Quantitative test for determination of [-2]proPSA levels. Silver Spring, MD: Food and Drug Administration; 2012.
50. Nicholson A, Mahon J, Boland A, et al. The clinical effectiveness and cost-effectiveness of the PROGENSA(R) prostate cancer antigen 3 assay and the Prostate Health Index in the diagnosis of prostate cancer: a systematic review and economic evaluation. *Health Technol Assess.* Oct 2015;19(87):1-192. PMID 26507078
51. Pecoraro V, Roli L, Plebani M, et al. Clinical utility of the (-2)proPSA and evaluation of the evidence: a systematic review. *Clin Chem Lab Med.* Jul 1 2016;54(7):1123-1132. PMID 26609863
52. Bruzzese D, Mazzarella C, Ferro M, et al. Prostate health index vs percent free prostate-specific antigen for prostate cancer detection in men with "gray" prostate-specific antigen levels at first biopsy: systematic review and meta-analysis. *Transl Res.* Dec 2014;164(6):444-451. PMID 25035153
53. Wang W, Wang M, Wang L, et al. Diagnostic ability of %p2PSA and prostate health index for aggressive prostate cancer: a meta-analysis. *Sci Rep.* 2014;4:5012. PMID 24852453
54. National Institute for Health and Care Excellence (NICE). Diagnosing prostate cancer: PROGENSA PCA3 assay and Prostate Health Index [DG17]. 2015; <https://www.nice.org.uk/guidance/dg17>.
55. Filella X, Gimenez N. Evaluation of [-2] proPSA and Prostate Health Index (phi) for the detection of prostate cancer: a systematic review and meta-analysis. *Clin Chem Lab Med.* Apr 2013;51(4):729-739. PMID 23154423
56. Fossati N, Lazzeri M, Haese A, et al. Clinical performance of serum isoform [-2]proPSA (p2PSA), and its derivatives %p2PSA and the Prostate Health Index, in men aged <60 years: results from a multicentric European study. *BJU Int.* Jun 2015;115(6):913-920. PMID 24589357
57. Boegemann M, Stephan C, Cammann H, et al. The percentage of prostate-specific antigen (PSA) isoform [-2]proPSA and the Prostate Health Index improve the diagnostic accuracy for clinically relevant prostate cancer at initial and repeat biopsy compared with total PSA and percentage free PSA in men aged <=65 years. *BJU Int.* Jan 2016;117(1):72-79. PMID 25818705
58. Morote J, Celma A, Planas J, et al. Diagnostic accuracy of prostate health index to identify aggressive prostate cancer. An Institutional validation study. *Actas Urol Esp.* Jul-Aug 2016;40(6):378-385. PMID 26923032
59. Yu GP, Na R, Ye DW, et al. Performance of the Prostate Health Index in predicting prostate biopsy outcomes among men with a negative digital rectal examination and transrectal ultrasonography. *Asian J Androl.* Jul-Aug 2016;18(4):633-638. PMID 26975483
60. Sreekumar A, Poisson LM, Rajendiran TM, et al. Metabolomic profiles delineate potential role for sarcosine in prostate cancer progression. *Nature.* Feb 12 2009;457(7231):910-914. PMID 19212411

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

61. Khan AP, Rajendiran TM, Ateeq B, et al. The role of sarcosine metabolism in prostate cancer progression. *Neoplasia*. May 2013;15(5):491-501. PMID 23633921
62. ProgenSA PCA3 assay. <http://www.hologic.com/products/clinical-diagnostics-blood-screening/assays-and-tests/progenSA-pca3-assay>. Accessed September 12, 2016.
63. Food and Drug Administration. Summary of Safety and Effectiveness Data. PMA P100033: PROGENSA PCA3 Assay. Silver Spring, MD: Food and Drug Administration; 2012.
64. Cui Y, Cao W, Li Q, et al. Evaluation of prostate cancer antigen 3 for detecting prostate cancer: a systematic review and meta-analysis. *Sci Rep*. 2016;6:25776. PMID 27161545
65. Rubio-Briones J, Borque A, Esteban LM, et al. Optimizing the clinical utility of PCA3 to diagnose prostate cancer in initial prostate biopsy. *BMC Cancer*. 2015;15:633. PMID 26362197
66. Bradley LA, Palomaki G, Gutman S, et al. PCA3 Testing for the Diagnosis and Management of Prostate Cancer (Comparative Effectiveness Reviews No. 98). Rockville, MD: Agency for Healthcare Research and Quality; 2013.
67. Ruiz-Aragon J, Marquez-Pelaez S. [Assessment of the PCA3 test for prostate cancer diagnosis: a systematic review and meta-analysis]. *Actas Urol Esp*. Apr 2010;34(4):346-355. PMID 20470697
68. Wei JT, Feng Z, Partin AW, et al. Can urinary PCA3 supplement PSA in the early detection of prostate cancer? *J Clin Oncol*. Dec 20 2014;32(36):4066-4072. PMID 25385735
69. Ankerst DP, Groskopf J, Day JR, et al. Predicting prostate cancer risk through incorporation of prostate cancer gene 3. *J Urol*. Oct 2008;180(4):1303-1308; discussion 1308. PMID 18707724
70. Chun FK, de la Taille A, van Poppel H, et al. Prostate cancer gene 3 (PCA3): development and internal validation of a novel biopsy nomogram. *Eur Urol*. Oct 2009;56(4):659-667. PMID 19304372
71. Perdon S, Cavadas V, Di Lorenzo G, et al. Prostate cancer detection in the "grey area" of prostate-specific antigen below 10 ng/ml: head-to-head comparison of the updated PCPT calculator and Chun's nomogram, two risk estimators incorporating prostate cancer antigen 3. *Eur Urol*. Jan 2011;59(1):81-87. PMID 20947244
72. Haese A, de la Taille A, van Poppel H, et al. Clinical utility of the PCA3 urine assay in European men scheduled for repeat biopsy. *Eur Urol*. Nov 2008;54(5):1081-1088. PMID 18602209
73. Nakanishi H, Groskopf J, Fritsche HA, et al. PCA3 molecular urine assay correlates with prostate cancer tumor volume: implication in selecting candidates for active surveillance. *J Urol*. May 2008;179(5):1804-1809; discussion 1809-1810. PMID 18353398
74. Whitman EJ, Groskopf J, Ali A, et al. PCA3 score before radical prostatectomy predicts extracapsular extension and tumor volume. *J Urol*. Nov 2008;180(5):1975-1978; discussion 1978-1979. PMID 18801539
75. Bostwick DG, Gould VE, Qian J, et al. Prostate cancer detected by uPM3: radical prostatectomy findings. *Mod Pathol*. May 2006;19(5):630-633. PMID 16528369
76. van Gils MP, Hessels D, Hulsbergen-van de Kaa CA, et al. Detailed analysis of histopathological parameters in radical prostatectomy specimens and PCA3 urine test results. *Prostate*. Aug 1 2008;68(11):1215-1222. PMID 18500693
77. Auprich M, Chun FK, Ward JF, et al. Critical assessment of preoperative urinary prostate cancer antigen 3 on the accuracy of prostate cancer staging. *Eur Urol*. Jan 2011;59(1):96-105. PMID 20980098
78. Tosoian JJ, Loeb S, Kettermann A, et al. Accuracy of PCA3 measurement in predicting short-term biopsy progression in an active surveillance program. *J Urol*. Feb 2010;183(2):534-538. PMID 20006883
79. Ruffion A, Devonec M, Champetier D, et al. PCA3 and PCA3-based nomograms improve diagnostic accuracy in patients undergoing first prostate biopsy. *Int J Mol Sci*. 2013;14(9):17767-17780. PMID 23994838
80. Ruffion A, Perrin P, Devonec M, et al. Additional value of PCA3 density to predict initial prostate biopsy outcome. *World J Urol*. Aug 2014;32(4):917-923. PMID 24500192
81. Merdan S, Tomlins SA, Barnett CL, et al. Assessment of long-term outcomes associated with urinary prostate cancer antigen 3 and TMPRSS2:ERG gene fusion at repeat biopsy. *Cancer*. Nov 15 2015;121(22):4071-4079. PMID 26280815
82. Suh CI, Shanafelt T, May DJ, et al. Comparison of telomerase activity and GSTP1 promoter methylation in ejaculate as potential screening tests for prostate cancer. *Mol Cell Probes*. Aug 2000;14(4):211-217. PMID 10970725
83. Goessl C, Muller M, Heicappell R, et al. DNA-based detection of prostate cancer in blood, urine, and ejaculates. *Ann N Y Acad Sci*. Sep 2001;945:51-58. PMID 11708494
84. Henrique R, Jeronimo C, Teixeira MR, et al. Epigenetic heterogeneity of high-grade prostatic intraepithelial neoplasia: clues for clonal progression in prostate carcinogenesis. *Mol Cancer Res*. Jan 2006;4(1):1-8. PMID 16446401

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

85. Eilers T, Machtens S, Tezval H, et al. Prospective diagnostic efficiency of biopsy washing DNA GSTP1 island hypermethylation for detection of adenocarcinoma of the prostate. *Prostate*. May 15 2007;67(7):757-763. PMID 17373715
86. Ellinger J, Albers P, Perabo FG, et al. CpG island hypermethylation of cell-free circulating serum DNA in patients with testicular cancer. *J Urol*. Jul 2009;182(1):324-329. PMID 19447423
87. Henrique R, Ribeiro FR, Fonseca D, et al. High promoter methylation levels of APC predict poor prognosis in sextant biopsies from prostate cancer patients. *Clin Cancer Res*. Oct 15 2007;13(20):6122-6129. PMID 17947477
88. Woodson K, O'Reilly KJ, Ward DE, et al. CD44 and PTGS2 methylation are independent prognostic markers for biochemical recurrence among prostate cancer patients with clinically localized disease. *Epigenetics*. Oct-Dec 2006;1(4):183-186. PMID 17998819
89. Ellinger J, Bastian PJ, Jurgan T, et al. CpG island hypermethylation at multiple gene sites in diagnosis and prognosis of prostate cancer. *Urology*. Jan 2008;71(1):161-167. PMID 18242387
90. Sunami E, Shinozaki M, Higano CS, et al. Multimarker circulating DNA assay for assessing blood of prostate cancer patients. *Clin Chem*. Mar 2009;55(3):559-567. PMID 19131636
91. Trock BJ, Brozman MJ, Mangold LA, et al. Evaluation of GSTP1 and APC methylation as indicators for repeat biopsy in a high-risk cohort of men with negative initial prostate biopsies. *BJU Int*. Jul 2012;110(1):56-62. PMID 22077694
92. Van Neste L, Herman JG, Otto G, et al. The epigenetic promise for prostate cancer diagnosis. *Prostate*. Aug 1 2012;72(11):1248-1261. PMID 22161815
93. Ge YZ, Xu LW, Jia RP, et al. The association between RASSF1A promoter methylation and prostate cancer: evidence from 19 published studies. *Tumour Biol*. Apr 2014;35(4):3881-3890. PMID 24353088
94. Moritz R, Ellinger J, Nuhn P, et al. DNA hypermethylation as a predictor of PSA recurrence in patients with low- and intermediate-grade prostate cancer. *Anticancer Res*. Dec 2013;33(12):5249-5254. PMID 24324057
95. Haldrup C, Mundbjerg K, Vestergaard EM, et al. DNA methylation signatures for prediction of biochemical recurrence after radical prostatectomy of clinically localized prostate cancer. *J Clin Oncol*. Sep 10 2013;31(26):3250-3258. PMID 23918943
96. Kachakova D, Mitkova A, Popov E, et al. Evaluation of the clinical value of the newly identified urine biomarker HIST1H4K for diagnosis and prognosis of prostate cancer in Bulgarian patients. *J BUON*. Jul-Sep 2013;18(3):660-668. PMID 24065480
97. Goessl C, Muller M, Heicappell R, et al. Methylation-specific PCR for detection of neoplastic DNA in biopsy washings. *J Pathol*. Mar 2002;196(3):331-334. PMID 11857497
98. Chu DC, Chuang CK, Fu JB, et al. The use of real-time quantitative polymerase chain reaction to detect hypermethylation of the CpG islands in the promoter region flanking the GSTP1 gene to diagnose prostate carcinoma. *J Urol*. Apr 2002;167(4):1854-1858. PMID 11912447
99. Mehrotra J, Varde S, Wang H, et al. Quantitative, spatial resolution of the epigenetic field effect in prostate cancer. *Prostate*. Feb 1 2008;68(2):152-160. PMID 18058812
100. Van Neste L, Bigley J, Toll A, et al. A tissue biopsy-based epigenetic multiplex PCR assay for prostate cancer detection. *BMC Urol*. 2012;12:16. PMID 22672250
101. Stewart GD, Van Neste L, Delvenne P, et al. Clinical utility of an epigenetic assay to detect occult prostate cancer in histopathologically negative biopsies: results of the MATLOC study. *J Urol*. Mar 2013;189(3):1110-1116. PMID 22999998
102. Partin AW, Van Neste L, Klein EA, et al. Clinical validation of an epigenetic assay to predict negative histopathological results in repeat prostate biopsies. *J Urol*. Oct 2014;192(4):1081-1087. PMID 24747657
103. Van Neste L, Partin AW, Stewart GD, et al. Risk score predicts high-grade prostate cancer in DNA-methylation positive, histopathologically negative biopsies. *Prostate*. Sep 2016;76(12):1078-1087. PMID 27121847
104. Wojno KJ, Costa FJ, Cornell RJ, et al. Reduced rate of repeated prostate biopsies observed in ConfirmMDx clinical utility field study. *Am Health Drug Benefits*. May 2014;7(3):129-134. PMID 24991397
105. Aubry W, Lieberthal R, Willis A, et al. Budget impact model: epigenetic assay can help avoid unnecessary repeated prostate biopsies and reduce healthcare spending. *Am Health Drug Benefits*. Jan 2013;6(1):15-24. PMID 24991343
106. Mackinnon AC, Yan BC, Joseph LJ, et al. Molecular biology underlying the clinical heterogeneity of prostate cancer: an update. *Arch Pathol Lab Med*. Jul 2009;133(7):1033-1040. PMID 19642730
107. Yao Y, Wang H, Li B, et al. Evaluation of the TMPRSS2:ERG fusion for the detection of prostate cancer: a systematic review and meta-analysis. *Tumour Biol*. Mar 2014;35(3):2157-2166. PMID 24142545
108. Tomlins SA, Aubin SM, Siddiqui J, et al. Urine TMPRSS2:ERG fusion transcript stratifies prostate cancer risk in men with elevated serum PSA. *Sci Transl Med*. Aug 3 2011;3(94):94ra72. PMID 21813756

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

109. Salami SS, Schmidt F, Laxman B, et al. Combining urinary detection of TMPRSS2:ERG and PCA3 with serum PSA to predict diagnosis of prostate cancer. *Urol Oncol.* Jul 2013;31(5):566-571. PMID 21600800
110. Tomlins SA, Day JR, Lonigro RJ, et al. Urine TMPRSS2:ERG Plus PCA3 for Individualized Prostate Cancer Risk Assessment. *Eur Urol.* Jul 2016;70(1):45-53. PMID 25985884
111. Prostate Core Mitomic Test: Clinical insight. <http://mdnalifesciences.com/prostate-core-mitomic-test/index.php>. Accessed September 21, 2016.
112. Parr RL, Dakubo GD, Crandall KA, et al. Somatic mitochondrial DNA mutations in prostate cancer and normal appearing adjacent glands in comparison to age-matched prostate samples without malignant histology. *J Mol Diagn.* Jul 2006;8(3):312-319. PMID 16825503
113. Maki J, Robinson K, Reguly B, et al. Mitochondrial genome deletion aids in the identification of false- and true-negative prostate needle core biopsy specimens. *Am J Clin Pathol.* Jan 2008;129(1):57-66. PMID 18089489
114. Robinson K, Creed J, Reguly B, et al. Accurate prediction of repeat prostate biopsy outcomes by a mitochondrial DNA deletion assay. *Prostate Cancer Prostatic Dis.* Jun 2010;13(2):126-131. PMID 20084081
115. Ma W, Diep K, Fritsche HA, et al. Diagnostic and prognostic scoring system for prostate cancer using urine and plasma biomarkers. *Genet Test Mol Biomarkers.* Mar 2014;18(3):156-163. PMID 24512523
116. Qu X, Randhawa G, Friedman C, et al. A three-marker FISH panel detects more genetic aberrations of AR, PTEN and TMPRSS2/ERG in castration-resistant or metastatic prostate cancers than in primary prostate tumors. *PLoS One.* 2013;8(9):e74671. PMID 24098661
117. Leyten GH, Hessels D, Smit FP, et al. Identification of a candidate gene panel for the early diagnosis of prostate cancer. *Clin Cancer Res.* Jul 1 2015;21(13):3061-3070. PMID 25788493
118. Xiao K, Guo J, Zhang X, et al. Use of two gene panels for prostate cancer diagnosis and patient risk stratification. *Tumour Biol.* Aug 2016;37(8):10115-10122. PMID 26820133
119. Little J, Wilson B, Carter R, et al. Multigene panels in prostate cancer risk assessment. *Evid Rep Technol Assess (Full Rep).* Jul 2012(209):1-166. PMID 24423032
120. Kader AK, Sun J, Reck BH, et al. Potential impact of adding genetic markers to clinical parameters in predicting prostate biopsy outcomes in men following an initial negative biopsy: findings from the REDUCE trial. *Eur Urol.* Dec 2012;62(6):953-961. PMID 22652152
121. Ioannidis JP, Castaldi P, Evangelou E. A compendium of genome-wide associations for cancer: critical synopsis and reappraisal. *J Natl Cancer Inst.* Jun 16 2010;102(12):846-858. PMID 20505153
122. Lindstrom S, Schumacher F, Siddiq A, et al. Characterizing associations and SNP-environment interactions for GWAS-identified prostate cancer risk markers-results from BPC3. *PLoS One.* 2011;6(2):e17142. PMID 21390317
123. Ishak MB, Giri VN. A systematic review of replication studies of prostate cancer susceptibility genetic variants in high-risk men originally identified from genome-wide association studies. *Cancer Epidemiol Biomarkers Prev.* Aug 2011;20(8):1599-1610. PMID 21715604
124. American Urological Association. Detection of prostate cancer, April 2013. <http://www.auanet.org/education/guidelines/prostate-cancer-detection.cfm>. Accessed March 2015.
125. Evaluation of Genomic Applications in Practice Prevention Working Group. Recommendations from the EGAPP Working Group: does PCA3 testing for the diagnosis and management of prostate cancer improve patient health outcomes? *Genet Med.* Apr 2014;16(4):338-346. PMID 24071797
126. Liss M, Ehdaie B, Loeb S, et al. The Prevention and Treatment of the More Common Complications Related to Prostate Biopsy Update. 2012; updated 2016; <https://www.auanet.org/guidelines/prostate-needle-biopsy-complications>. Accessed October 23, 2017.
127. Rosenkrantz AB, Verma S, Choyke P, et al. Prostate magnetic resonance imaging and magnetic resonance imaging targeted biopsy in patients with a prior negative biopsy: a consensus statement by AUA and SAR. *J Urol.* Dec 2016;196(6):1613-1618. PMID 27320841

Policy History

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

10/14/2010 Medical Policy Committee review

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

10/20/2010	Medical Policy Implementation Committee approval. New policy.
10/06/2011	Medical Policy Committee review
10/19/2011	Medical Policy Implementation Committee approval. Minor change to coverage statement ("prognosis" added to the investigational statement on PCA3).
10/11/2012	Medical Policy Committee review
10/31/2012	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
02/19/2013	Coding updated
10/03/2013	Medical Policy Committee review
10/16/2013	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
12/04/2014	Medical Policy Committee review
12/17/2014	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
08/06/2015	Medical Policy Committee review
08/19/2015	Medical Policy Implementation Committee approval. Added Kallikrein markers (4Kscore test), metabolomics profiles (Prostarix), candidate gene panels, mitochondrial DNA mutation testing (Prostate Core Mitomics test), and gene hypermethylation testing (ConfirmMDx) to INV statement. Title change.
10/06/2016	Medical Policy Committee review
10/19/2016	Medical Policy Implementation Committee approval. No change to coverage.
01/01/2017	Coding update: Removing ICD-9 Diagnosis Codes and CPT coding update
01/05/2017	Medical Policy Committee review
01/18/2017	Medical Policy Implementation Committee approval. Added Prostate Health Index (phi) to investigational statement and rationale. Updated rationale and references.
01/04/2018	Medical Policy Committee review
01/17/2018	Medical Policy Implementation Committee approval. Policy revised to separate initial biopsy and repeat biopsy populations, policy statement otherwise unchanged.
10/29/2018	Coding update
Next Scheduled Review Date:	01/2019

Coding

The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)†, copyright 2017 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Policy # 00272

Original Effective Date: 10/20/2010

Current Effective Date: 01/17/2018

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	81313, 81479, 81539, 81599 Codes added eff 1/1/2018: 81551
HCPCS	No codes
ICD-10 Diagnosis	C61, Z12.5

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. FDA and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
 - 1. Consultation with the Blue Cross and Blue Shield Association technology assessment program (TEC) or other nonaffiliated technology evaluation center(s);
 - 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
 - 3. Reference to federal regulations.

‡ Indicated trademarks are the registered trademarks of their respective owners.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

©2018 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.