



Louisiana

GLP-1 Agonists for Diabetes

Policy # 00324

Original Effective Date: 11/16/2011

Current Effective Date: 10/12/2020

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

When Services May Be Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Based on review of the available data, the Company may consider glucagon like peptide 1 (GLP-1) agonists that are FDA approved for the treatment of diabetes, including but not limited to Byetta[®] (exenatide), Bydureon[®]/Bydureon BCise (exenatide ER), Victoza[®] (liraglutide), Trulicity[™] (dulaglutide), Adlyxin[®] (lixisenatide), Ozempic[®] (semaglutide), and Rybelsus[®] (semaglutide), to be **eligible for coverage**** when the patient selection criteria below are met for the requested drug:

Patient Selection Criteria

Coverage eligibility will be considered when the patient selection criteria are met for the requested drug:

- For Victoza (liraglutide), Trulicity (dulaglutide), Ozempic (semaglutide), or Rybelsus (semaglutide) requests:
 - Patient has type 2 diabetes mellitus; OR
- For Byetta (exenatide), Bydureon/Bydureon BCise (exenatide ER), or Adlyxin (lixisenatide) requests:
 - Patient has type 2 diabetes mellitus; AND
 - Patient has tried and failed (e.g., intolerance or inadequate response) at least TWO of the following products: Victoza (liraglutide), Trulicity (dulaglutide), Ozempic (semaglutide), or Rybelsus (semaglutide).

*(Note: This specific patient criterion is an additional Company requirement for coverage eligibility and will be denied as not medically necessary** if not met)*

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

GLP-1 Agonists for Diabetes

Policy # 00324

Original Effective Date: 11/16/2011

Current Effective Date: 10/12/2020

When Services Are Considered Not Medically Necessary

Based on review of available data, the Company considers the use of Byetta (exenatide), Bydureon/Bydureon BCise (exenatide ER), or Adlyxin (lixisenatide) WITHOUT having tried and failed at least TWO of the following products: Victoza (liraglutide), Trulicity (dulaglutide), Ozempic (semaglutide), or Rybelsus (semaglutide) to be **not medically necessary**.**

When Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Based on review of available data, the Company considers the use of GLP-1 agonists that are FDA approved for the treatment of diabetes for any non-FDA approved indication for that specific drug to be **investigational**.*

Schematic

| Preferred | Non-Preferred |
|---------------------------------------------|----------------------------------------------|
| Trulicity Victoza Ozempic Rybelsus | Byetta Bydureon/Bydureon BCise Adlyxin |

Background/Overview

Byetta, Bydureon/Bydureon BCise, Victoza, Trulicity, Adlyxin, and Ozempic are antihyperglycemic agents for subcutaneous injection. Rybelsus is the only GLP-1 agonist product that is currently in an oral dosage form. These products are incretin mimetic agents that bind and activate the human GLP-1 receptor. Activation of this receptor increases glucose-dependent insulin secretion by pancreatic beta-cells and suppresses glucagon secretion and slows gastric emptying. Byetta, Bydureon/Bydureon BCise, Victoza, Trulicity, Adlyxin, Ozempic, and Rybelsus are FDA approved in adjunct to diet and exercise to improve glycemic control in patients with type 2 diabetes. Byetta is administered subcutaneously twice daily. Bydureon/Bydureon BCise, Trulicity, and Ozempic are administered subcutaneously once weekly, and Victoza and Adlyxin are administered subcutaneously once daily. Rybelsus is taken orally once daily.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

GLP-1 Agonists for Diabetes

Policy # 00324

Original Effective Date: 11/16/2011

Current Effective Date: 10/12/2020

The active ingredients in the non-preferred products have not demonstrated superiority in head to head studies comparing preferred and non-preferred products.

This policy is also intended to ensure that the GLP-1 agonist products approved for the treatment of type 2 diabetes are used for the indication of type 2 diabetes only.

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

All of the products mentioned in this policy are FDA approved for the treatment of type 2 diabetes mellitus.

Rationale/Source

As monotherapy, Byetta 5 mcg or 10 mcg twice daily in adjunct to diet and exercise reduced glycosylated hemoglobin (HbA1c) 0.5 to 0.7% (placebo corrected); a placebo corrected weight loss of 2.7 kg to 2.9 kg was noted at the end of the 24 week trial. In general, Byetta appears to lower HbA1c by 0.5% to 1%. In addition to HbA1c reduction, Byetta reduces food intake, and on average produces a 2 kg to 3 kg weight loss over a 6-month period in diabetic patients. Byetta 10 mcg twice daily as an adjunct to metformin, a sulfonylurea, or both in patients with type 2 diabetes decreased body weight by 1.6 kg to 2.8 kg after 30 weeks. In an interim analysis involving a 52 week open-label uncontrolled extension study, which followed the 30 week double-blind period, the average weight loss in type 2 diabetics (n = 314) after a total of 82 weeks of Byetta therapy was 4.4 kg. A similar analysis of an interim report noted weight loss in type 2 diabetics (n = 92) after 82 weeks of Byetta treatment was 5.3 kg. In a multicenter, open-label, randomized, controlled trial in patients with type 2 diabetes (n = 551), at 26 weeks, treatment with Byetta led to a 2.3 kg reduction in body weight compared with a 1.8 kg increase for patients treated with insulin glargine (Lantus[®])[†]. Addition of Byetta to a thiazolidinedione ([TZD] with or without metformin) resulted in a 1.51 kg mean reduction in bodyweight after 16 weeks. Reductions in bodyweight in type 2 diabetic patients treated with Byetta have been sustained for up to two years.

As monotherapy in a 52 week trial, Victoza 1.2 mg and 1.8 mg in adjunct to diet and exercise resulted in mean HbA1c reduction of 0.8% to 1.1% and a 2.1 kg to 2.5 kg weight reduction. Victoza was studied in combination with one or two other oral anti-diabetic agents in four 26 week studies. When added to metformin, Victoza 1.8 mg and 1.2 mg resulted in a mean placebo corrected HbA1c and

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

GLP-1 Agonists for Diabetes

Policy # 00324

Original Effective Date: 11/16/2011

Current Effective Date: 10/12/2020

weight reduction of 1.1% and 1.1 kg to 1.3 kg, respectively. As add-on to sulfonylurea (glimeperide), Victoza 1.2 mg and 1.8 mg treatment resulted in a placebo corrected mean HbA1c reduction of 1.3% to 1.4%. As part of a triple therapy combination with metformin and glimeperide, Victoza 1.8 mg reduced HbA1c (placebo corrected mean) by 1.1% and resulted in a mean weight reduction of 1.4 kg (placebo corrected). When added to metformin and rosiglitazone mean placebo corrected reduction in HbA1c and weight with Victoza (1.8 mg and 1.2 mg) were 0.9% (both doses) and 2.6 kg and 1.6 kg, respectively. In a head-to-head trial with Byetta, weight was significantly reduced in both Byetta (10 mcg twice daily) and Victoza (1.8 mg daily) and was non-significant between groups (-2.87 kg vs. -3.24 kg, respectively).

A randomized, open-label 24-week comparative trial was conducted with Bydureon and Byetta for safety and efficacy in 252 patients with type 2 diabetes. These patients had inadequate glycemic control with diet and exercise alone or with oral antidiabetic therapy, including metformin, a sulfonylurea, a TZD, or combination of two of those therapies. Patients were treated with diet and exercise alone (19%), a single oral antidiabetic agent (47%), or combination therapy of oral antidiabetic agents (35%). The mean baseline HbA1c was 8.4%. Patients were randomly assigned to receive Bydureon 2 mg once every seven days (weekly) or Byetta (10 mcg twice-daily), in addition to existing oral antidiabetic agents. Patients assigned to Byetta initiated treatment with 5 mcg twice-daily then increased the dose to 10 mcg twice-daily after 4 weeks. The primary endpoint was change in HbA1c from baseline to week 24 (or the last value at time of early discontinuation). Change in body weight was a secondary endpoint. Treatment with Bydureon was superior to Byetta for mean HbA1c reduction over 24 weeks.

Trulicity was studied in various trials as monotherapy as well as in addition to oral therapies and in addition to insulin. As monotherapy, Trulicity lowered the HbA1c from 0.7-0.8% vs. metformin's lowering of 0.6%. Trulicity as monotherapy also lowered the fasting plasma glucose by 26 to 29 mg/dL vs. a lowering of 24 mg/dL with metformin. In the combo therapy trials, the HbA1c lowering ranged from 0.8 to 1.6% depending on the treatment that Trulicity was combined with.

Adlyxin was studied in various trials as monotherapy as well as in addition to oral therapies and in addition to insulin. As monotherapy, Adlyxin lowered the HbA1c by 0.83% from baseline (0.65% difference from placebo). Adlyxin also lowered fasting plasma glucose by 15.84 mg/dL in the monotherapy trial. The range of lowering of HbA1c in the combination studies ranges from 0.7-0.91% depending on what drug Adlyxin was combined with.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

GLP-1 Agonists for Diabetes

Policy # 00324

Original Effective Date: 11/16/2011

Current Effective Date: 10/12/2020

Ozempic has been studied as monotherapy and in combination with metformin, metformin and sulfonylureas, metformin and/or thiazolidinedione, and basal insulin in patients with type 2 diabetes mellitus. The efficacy of Ozempic was compared with placebo, sitagliptin, Bydureon, and insulin glargine. Most trials evaluated the use of Ozempic 0.5 mg, and 1 mg, with the exception of the trial comparing Ozempic and Bydureon where only the 1 mg dose was studied. In patients with type 2 diabetes mellitus, Ozempic produced clinically relevant reduction from baseline in HbA1c compared with placebo. The various HbA1c lowering ranged from 1.1-1.6% depending on the clinical comparison.

Rybelsus' study program included 10 randomized, placebo-controlled or active-comparator trials. Studies included comparisons between Rybelsus and placebo, other oral anti-diabetic agents (Jardiance, Januvia), other GLP-1 agonist products (Victoza, Ozempic), along with additions to metformin and sulfonylurea. The various HbA1c lowering ranged from 0.6-1.9% depending on the clinical comparison.

Based on a review of the available data and in the absence of any of the caveats mentioned, there is no advantage of using the non-preferred agents mentioned in this policy over the preferred agents mentioned in this policy.

References

1. Express Scripts Prior Authorization policy on Glucagon-Like Peptide (GLP-1) Agonists (Byetta [exenatide injection Amylin] and Victoza [liraglutide injection- NovoNordisk]). May 2014.
2. Byetta injection [package insert]. San Diego, CA: Amylin Pharmaceuticals, Inc.; October 2011.
3. Victoza [package insert]. Princeton, NJ: Novo Nordisk, Inc.; May 2013.
4. Bydureon for injectable suspension [package insert]. San Diego, CA: Amylin Pharmaceuticals; May 2013.
5. Moretto TJ, Milton DR, Ridge TD, et al. Efficacy and tolerability of exenatide monotherapy over 24 weeks in antidiabetic drug-naive patients with type 2 diabetes: a randomized, double-blind, placebo-controlled, parallel-group study. *Clin Ther.* 2008;30(8):1448-60.
6. Nathan DM, Buse JB, Davidson MB, Ferrannini E et al. Management of hyperglycemia in type 2 diabetes: a consensus algorithm for the initiation and adjustment of therapy. A consensus statement from the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetes Care.* 2009;32(1):193-203.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

GLP-1 Agonists for Diabetes

Policy # 00324

Original Effective Date: 11/16/2011

Current Effective Date: 10/12/2020

7. DeFronzo RA, Ratner RE, Han J, Kim DD, Fineman MS, Baron AD. Effects of exenatide (Exendin-4) on glycemic control and weight over 30 weeks in metformin-treated patients with type 2 diabetes. *Diabetes Care*. 2005;28(5):1092-1100.
8. Buse JB, Henry RR, Han J, et al. Effects of exenatide (Exendin-4) on glycemic control over 30 weeks in sulfonylurea-treated patients with type 2 diabetes. *Diabetes Care*. 2004;27(11):2628-2635.
9. Kendall DM, Riddle MC, Rosenstock J, et al. Effects of exenatide (Exendin-4) on glycemic control over 30 weeks in patients with type 2 diabetes treated with metformin and a sulfonylurea. *Diabetes Care*. 2005;28(5):1083-1091.
10. Blonde L, Klein EJ, Han J, et al. Interim analysis of the effects of exenatide treatment on A1C, weight and cardiovascular risk factors over 82 weeks in 314 overweight patients with type 2 diabetes. *Diabetes Obes Metab*. 2006;8:436-447.
11. Ratner RE, Maggs D, Nielsen LL, et al. Long-term effects of exenatide therapy over 82 weeks on glycaemic control and weight in over-weight metformin-treated patients with type 2 diabetes mellitus. *Diabetes Obes Metab*. 2006;8:419-428.
12. Heine RJ, Van Gaal LF, Johns D, et al. Exenatide versus insulin glargine in patients with suboptimally controlled type 2 diabetes. A randomized trial. *Ann Intern Med*. 2005;143:559-569.
13. Cretkovic RS, Plosker GL. Exenatide: a review of its use in patients with type 2 diabetes mellitus (as an adjunct to metformin and/or a sulfonylurea). *Drugs*. 2007;67(6):935-954.
14. Garber A, Henry R, Ratner R, et al; for the LEAD-3 (Mono) study group. Liraglutide versus glimeperide monotherapy for type 2 diabetes (LEAD-3 mono): a randomized, 52-week, phase III, double-blind, parallel-treatment trial. *Lancet*. 2009;373:473-481.
15. Nauck M, Frid A, Hermansen K, et al; for the LEAD-2 study group. Efficacy and safety comparison of liraglutide, glimeperide, and placebo, all in combination with metformin, in type 2 diabetes. *Diabetes Care*. 2009;32:84-90.
16. Marre M, Shaw J, Brandle M, et al; on behalf of the LEAD-1 SU study group. Liraglutide, a once-daily human GLP-1 analogue, added to a sulphonylurea over 26 weeks produces greater improvements in glycaemic and weight control compared with adding rosiglitazone or placebo in subjects with type 2 diabetes (LEAD-1 SU). *Diabet Med*. 2009;26:268-278.
17. Russel-Jones D, Vaag A, Schmitz O, et al; on behalf of the Liraglutide Effect and Action in Diabetes 5 (LEAD-5) met + SU study group. *Diabetologia*. 2009;52:2046-2055.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

GLP-1 Agonists for Diabetes

Policy # 00324

Original Effective Date: 11/16/2011

Current Effective Date: 10/12/2020

18. Zinman B, Gerich J, Buse JB, et al. Efficacy and safety of the human glucagon-like peptide-1 analog liraglutide in combination with metformin and thiazolidinediones in patients with type 2 diabetes (LEAD-4 Met + TZD). *Diabetes Care*. 2009;32(7):1224-1230.
19. Buse JB, Rosenstock J, Sesti G, et al; for the LEAD-6 study group. Liraglutide once a day versus exenatide twice a day for type 2 diabetes: a 26-week randomized, parallel-group, multinational, open-label trial (LEAD-6). *Lancet*. 2009;374:39-47.
20. Rodbard, HW, Jellinger PS, Davidson JA, et al. Statement by an American Association of Clinical Endocrinologists/American College of Endocrinology consensus panel on type 2 diabetes mellitus: An algorithm for glycemic control. *Endocr Pract*. 2009;15(6):541-559.
21. Astrup A, Rossner S, Van Gaal L et al. Effects of liraglutide in the treatment of obesity: a randomized, double-blind, placebo controlled study. *Lancet*. 2009;1606-1616.
22. Rosenstock J, Kladd LJ, Northrup J, et al. Effects of exenatide and lifestyle modification on body weight and glucose tolerance in obese subjects with and without pre-diabetes. *Diabetes Care*. 2010;33(6):1173-1175.
23. Raman VS, Mason KJ, Rodriguez LM, et al. The role of adjunctive exenatide in pediatric type 1 diabetes. *Diabetes Care*. 2010;22(6):1294-1296.
24. Kielgast U, Krarup T, Holst JJ and Madsbad S. Four weeks of treatment with liraglutide reduces insulin dose without loss of glycemic control in type 1 diabetic patients with and without residual β -cell function. *Diabetes Care*. 2011;34:1463-1468.
25. Varanasi A, Bellini N, Rawal D, et al. Liraglutide as additional treatment for type 1 diabetes. *Eur J Endocrinol*. 2011;165:77-84.
26. Trulicity [package insert]. Eli Lilly and Company. Indianapolis, IN. March 2015.
27. Adylin [package insert]. Sanofi-Aventis. Bridgewater, New Jersey. July 2016.
28. Ozempic [package insert]. Novo Nordisk. Plainsboro, New Jersey. December 2017.
29. Rybelsus [package insert]. Novo Nordisk. Plainsboro, New Jersey. September 2019.

Policy History

Original Effective Date: 11/16/2011

Current Effective Date: 10/12/2020

11/03/2011 Medical Policy Committee review

11/16/2011 Medical Policy Implementation Committee approval. New policy.

11/01/2012 Medical Policy Committee review

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

GLP-1 Agonists for Diabetes

Policy # 00324

Original Effective Date: 11/16/2011

Current Effective Date: 10/12/2020

| | |
|------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 11/28/2012 | Medical Policy Implementation Committee approval. Added Bydureon (exenatide ER) to the title and coverage statement. |
| 11/07/2013 | Medical Policy Committee review |
| 11/20/2013 | Medical Policy Implementation Committee approval. Revision to coverage language without changing the intent of the policy. Coverage eligibility unchanged. |
| 11/06/2014 | Medical Policy Committee review |
| 11/21/2014 | Medical Policy Implementation Committee approval. Changed title. Added Tanzeum to the policy. Updated background information and rationale to reflect new product and title change. |
| 04/02/2015 | Medical Policy Committee review |
| 04/20/2015 | Medical Policy Implementation Committee approval. Changed title. Added Trulicity to policy. Updated background and rationale. |
| 04/07/2016 | Medical Policy Committee review |
| 04/20/2016 | Medical Policy Implementation Committee approval. No change to coverage. |
| 10/06/2016 | Medical Policy Committee review |
| 10/19/2016 | Medical Policy Implementation Committee approval. Chose preferred products in this class (Byetta, Bydureon, Victoza, and Trulicity). |
| 03/02/2017 | Medical Policy Committee review |
| 03/15/2017 | Medical Policy Implementation Committee approval. Clarified to use two preferred products. |
| 03/01/2018 | Medical Policy Committee review |
| 03/21/2018 | Medical Policy Implementation Committee approval. No change to coverage. |
| 07/05/2018 | Medical Policy Committee review |
| 07/11/2018 | Medical Policy Implementation Committee approval. Added Ozempic and Bydureon BCise to the policy. |
| 09/06/2018 | Medical Policy Committee review |
| 09/19/2018 | Medical Policy Implementation Committee approval. Moved Byetta and Bydureon/Bydureon Bcise to non-preferred |
| 09/05/2019 | Medical Policy Committee review |
| 09/11/2019 | Medical Policy Implementation Committee approval. No change to coverage. |
| 12/05/2019 | Medical Policy Committee review |
| 12/11/2019 | Medical Policy Implementation Committee approval. Added a new drug, Rybelsus, as a preferred option. Removed Tanzeum from the policy as it has been discontinued. |

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

GLP-1 Agonists for Diabetes

Policy # 00324

Original Effective Date: 11/16/2011

Current Effective Date: 10/12/2020

09/03/2020 Medical Policy Committee review

09/09/2020 Medical Policy Implementation Committee approval. No change to coverage.

Next Scheduled Review Date: 09/2021

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
 1. Consultation with the Blue Cross and Blue Shield Association technology assessment program (TEC) or other nonaffiliated technology evaluation center(s);
 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
 3. Reference to federal regulations.

**Medically Necessary (or “Medical Necessity”) - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. In accordance with nationally accepted standards of medical practice;
- B. Clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and
- C. Not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

GLP-1 Agonists for Diabetes

Policy # 00324

Original Effective Date: 11/16/2011

Current Effective Date: 10/12/2020

For these purposes, “nationally accepted standards of medical practice” means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

‡ Indicated trademarks are the registered trademarks of their respective owners.

NOTICE: If the Patient’s health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.