



Louisiana

Genetic Testing for the Diagnosis of inherited Peripheral Neuropathies

Policy # 00378

Original Effective Date: 08/21/2013

Current Effective Date: 05/11/2020

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

When Services May Be Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Based on review of available data, the Company may consider genetic testing when the diagnosis of an inherited peripheral motor or sensory neuropathy is suspected due to signs and/or symptoms but a definitive diagnosis cannot be made to be **eligible for coverage.****

Patient Selection Criteria:

Coverage eligibility will be considered in individuals when the following criteria are met:

- Patients with cryptogenic polyneuropathy who exhibit a hereditary neuropathy phenotype may be considered for initial genetic testing of the most common genetic abnormalities, e.g. PMP22 (CMT1A duplication/ deletion), GJB1, or MFN2. Initial genetic testing should be guided by the clinical phenotype, inheritance pattern, and electrodiagnostic features. If these initial tests are negative, then second bullet will be considered.
- Evaluation of rarer genetic causes would be appropriate only if initial testing is negative.

When Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

The use of genetic testing when the diagnosis of an inherited peripheral motor or sensory neuropathy is suspected due to signs and/or symptoms but a definitive diagnosis cannot be made when patient selection criteria are not met is considered to be **investigational.***

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic Testing for the Diagnosis of Inherited Peripheral Neuropathies

Policy # 00378

Original Effective Date: 08/21/2013

Current Effective Date: 05/11/2020

Based on review of available data, the Company considers genetic testing for an inherited peripheral motor or sensory neuropathy for all other indications to be **investigational**.*

Based on review of available data, the Company considers initial testing with comprehensive multigene panels that test most known genes related to hereditary neuropathies to be **investigational**.*

Policy Guidelines

This policy addresses the hereditary motor and sensory peripheral neuropathies, of which peripheral neuropathy is the primary clinical manifestation. A number of other hereditary disorders may have neuropathy as an associated finding but typically have other central nervous system and occasional other systemic findings. Examples include Refsum disease, various lysosomal storage diseases, and mitochondrial disorders.

Genetic Counseling

Genetic counseling is primarily aimed at patients who are at risk for inherited disorders, and experts recommend formal genetic counseling in most cases when genetic testing for an inherited condition is considered. The interpretation of the results of genetic tests and the understanding of risk factors can be very difficult and complex. Therefore, genetic counseling will assist individuals in understanding the possible benefits and harms of genetic testing, including the possible impact of the information on the individual's family. Genetic counseling may alter the utilization of genetic testing substantially and may reduce inappropriate testing. Genetic counseling should be performed by an individual with experience and expertise in genetic medicine and genetic testing methods.

Background/Overview

Inherited Peripheral Neuropathies

Inherited peripheral neuropathies are a clinically and genetically heterogeneous group of disorders. The estimated prevalence in aggregate is 1 in 2500 persons, making inherited peripheral neuropathies the most common inherited neuromuscular disease.

Peripheral neuropathies can be subdivided into two major categories: primary axonopathies and primary myelinopathies, depending on which portion of the nerve fiber is affected. The further anatomic classification includes fiber type (eg, motor vs sensory, large vs small) and gross distribution of the nerves affected (eg, symmetry, length-dependency).

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic Testing for the Diagnosis of Inherited Peripheral Neuropathies

Policy # 00378

Original Effective Date: 08/21/2013

Current Effective Date: 05/11/2020

Inherited peripheral neuropathies are divided into the hereditary motor and sensory neuropathies, hereditary neuropathy with liability to pressure palsies (HNPP), and other miscellaneous, rare types (eg, hereditary brachial plexopathy, hereditary sensory, autonomic neuropathies). Other hereditary metabolic disorders, such as Friedreich ataxia, Refsum disease, and Krabbe disease, may be associated with motor and/or sensory neuropathies but typically have other predominating symptoms. This evidence review focuses on the hereditary motor and sensory neuropathies and HNPP.

A genetic etiology of peripheral neuropathy is typically suggested by generalized polyneuropathy, family history, lack of positive sensory symptoms, early age of onset, symmetry, associated skeletal abnormalities, and very slowly progressive clinical course. A family history of at least three generations with details on health issues, the cause of death, and age at death should be collected.

Charcot-Marie-Tooth Disease

Hereditary Motor and Sensory Neuropathies

Most inherited polyneuropathies were originally described clinically as variants of CMT disease. The clinical phenotype of CMT is highly variable, ranging from minimal neurologic findings to the classic picture with pes cavus and “stork legs” to a severe polyneuropathy with respiratory failure. CMT disease is genetically and clinically heterogeneous. Variants in more than 30 genes and more than 44 different genetic loci have been associated with the inherited neuropathies. Also, different pathogenic variants in a single gene can lead to different inherited neuropathy phenotypes and inheritance patterns. A 2016 cross-sectional study of 520 children and adolescents with CMT found variability in CMT-related symptoms across the 5 most commonly represented subtypes.

CMT subtypes are characterized by variants in one of several myelin genes, which lead to abnormalities in myelin structure, function, or upkeep. There are seven subtypes of CMT, with type 1 and 2 representing the most common hereditary peripheral neuropathies.

Most cases of CMT are autosomal dominant, although autosomal recessive and X-linked dominant forms exist. Most cases are CMT type 1 (approximately 40%-50% of all CMT cases, with 78%-80% of those due to *PMP22* variants). CMT type 2 is associated with 10% to 15% of CMT cases, with 20% of those due to *MFN2* variants.

A summary of the molecular genetics of CMT is outlined in Table 1.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic Testing for the Diagnosis of Inherited Peripheral Neuropathies

Policy # 00378

Original Effective Date: 08/21/2013

Current Effective Date: 05/11/2020

Table 1. Molecular Genetics of CMT Variants

Locus	Gene	Protein Product	Prevalence (if known)
CMT type 1			
CMT1A	<i>PMP22</i>	Peripheral myelin protein 22	70%-80% of CMT1
CMT1B	<i>MPZ</i>	Myelin P0 protein	10%-12% of CMT1
CMT1C	<i>LITAF</i>	Lipopolysaccharide-induced tumor necrosis factor- α factor	\approx 1% of CMT1
CMT1D	<i>EGR2</i>	Early growth response protein 2	
CMT1E	<i>PMP22</i>	Peripheral myelin protein 22 (sequence changes)	\approx 1% of CMT1
CMT1F/2E	<i>NEFL</i>	Neurofilament light polypeptide	
CMT type 2			
CMT2A1	<i>KIF1B</i>	Kinesin-like protein KIF1B	
CMT2A2	<i>MFN2</i>	Mitofusin-2	20% of CMT2
CMT2B	<i>RAB7A</i>	Ras-related protein Rab-7	
CMT2B1	<i>LMNA</i>	Lamin A/C	
CMT2B2	<i>MED25</i>	Mediator of RNA polymerase II transcription subunit 25	
CMT2C	<i>TRPV4</i>	Transient receptor potential cation channel subfamily V member 4	
CMT2D	<i>GARS</i>	Glycyl-tRNA synthetase	
CMT2E/1F	<i>NEFL</i>	Neurofilament light polypeptide	
CMT2F	<i>HSPB1</i>	Heat-shock protein beta-1	
CMT2G	12q12-q13	Unknown	
CMT2H/2K	<i>GDAP1</i>	Ganglioside-induced differentiation-associated protein 1	
CMT2I/2J	<i>MPZ</i>	Myelin P0 protein	
CMT2L	<i>HSPB8</i>	Heat-shock protein beta-8	
CMT2N	<i>AARS</i>	Alanyl-tRNA synthetase, cytoplasmic	
CMT2O	<i>DYNC1H1</i>	Cytoplasmic dynein 1 heavy chain 1	
CMT2P	<i>LRSAM1</i>	E3 ubiquitin-protein ligase LRSAM1	

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic Testing for the Diagnosis of Inherited Peripheral Neuropathies

Policy # 00378

Original Effective Date: 08/21/2013

Current Effective Date: 05/11/2020

Locus	Gene	Protein Product	Prevalence (if known)
CMT2S	<i>IGHMBP2</i>	DNA-binding protein SMUBP-2	
CMT2T	<i>DNAJB2</i>	DnaJ homolog subfamily B member 2	
CMT2U	<i>MARS</i>	Methionine-tRNA ligase, cytoplasmic	
CMT type 4			
CMT4A	<i>GDAP1</i>	Ganglioside-induced differentiation-associated protein 1	
CMT4B1	<i>MTMR2</i>	Myotubularin-related protein 2	
CMT4B2	<i>SBF2</i>	Myotubularin-related protein 13	
CMT4C	<i>SH3TC2</i>	SH3 domain and tetratricopeptide repeats-containing protein 2	
CMT4D	<i>NDRG1</i>	Protein NDRG1	
CMT4E	<i>EGR2</i>	Early growth response protein 2	
CMT4F	<i>PRX</i>	Periaxin	
CMT4H	<i>FGD4</i>	FYVE, RhoGEF and PH domain-containing protein 4	
CMT4J	<i>FIG4</i>	Phosphatidylinositol 3, 5-biphosphate	
X-linked CMT			
CMTX1	<i>GJB1</i>	Gap junction beta-1 protein (connexin 32)	90% of X-linked CMT
CMTX2	<i>Xp22.2</i>	Unknown	
CMTX3	<i>Xq26</i>	Unknown	
CMTX4	<i>AIFM1</i>	Apoptosis-inducing factor 1	
CMTX5	<i>PRPS1</i>	Ribose-phosphate pyrophosphokinase 1	
CMTX6	<i>PKD3</i>	Pyruvate dehydrogenase kinase isoform 3	

Adapted from Bird (2016).

CMT: Charcot-Marie-Tooth.

The clinical features of CMT are briefly summarized.

CMT Type 1

CMT1 is an autosomal dominant, demyelinating peripheral neuropathy characterized by distal muscle weakness and atrophy, sensory loss, and slow nerve conduction velocity. It is usually slowly progressive

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic Testing for the Diagnosis of Inherited Peripheral Neuropathies

Policy # 00378

Original Effective Date: 08/21/2013

Current Effective Date: 05/11/2020

and often associated with pes cavus foot deformity, bilateral foot drop, and palpably enlarged nerves, especially the ulnar nerve at the olecranon groove and the greater auricular nerve. Affected people usually become symptomatic between ages 5 and 25 years, and lifespan is not shortened. Less than 5% of people become wheelchair-dependent. CMT1 is inherited in an autosomal dominant manner. The CMT1 subtypes (CMT 1A-E) are separated by molecular findings and are often clinically indistinguishable. CMT1A accounts for 70% to 80% of all CMT1, and about two-thirds of probands with CMT1A have inherited the disease-causing variant, and about one-third have CMT1A as the result of a de novo variant.

CMT1A involves duplication of the *PMP22* gene. *PMP22* encodes an integral membrane protein, peripheral membrane protein 22, which is a major component of myelin in the peripheral nervous system. The phenotypes associated with this disease arise because of abnormal *PMP22* gene dosage effects. Two normal alleles represent the normal wild-type condition. Four normal alleles (as in the homozygous CMT1A duplication) result in the most severe phenotype, whereas three normal alleles (as in the heterozygous CMT1A duplication) cause a less severe phenotype.

CMT Type 2

CMT2 is a non-demyelinating (axonal) peripheral neuropathy characterized by distal muscle weakness and atrophy, mild sensory loss, and normal or near-normal nerve conduction velocities. Clinically, CMT2 is similar to CMT1, although typically less severe. The subtypes of CMT2 are similar clinically and distinguished only by molecular genetic findings. CMT2B1, CMT2B2, and CMT2H/K are inherited in an autosomal recessive manner; all other subtypes of CMT2 are inherited in an autosomal dominant manner. The most common subtype of CMT2 is CMT2A, which accounts for approximately 20% of CMT2 cases and is associated with variants in the *MFN2* gene.

X-Linked CMT

CMT X type 1 is characterized by a moderate-to-severe motor and sensory neuropathy in affected males and mild to no symptoms in carrier females. Sensorineural deafness and central nervous system symptoms also occur in some families. CMT X type 1 is inherited in an X-linked dominant manner. Molecular genetic testing of *GJB1* (Cx32), which is available on a clinical basis, detects about 90% of cases of CMT X type 1.

CMT Type 4

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic Testing for the Diagnosis of Inherited Peripheral Neuropathies

Policy # 00378

Original Effective Date: 08/21/2013

Current Effective Date: 05/11/2020

CMT type 4 is a form of hereditary motor and sensory neuropathy that is inherited in an autosomal recessive fashion and occurs secondary to myelinopathy or axonopathy. It occurs more rarely than the other forms of CMT neuropathy, but some forms may be rapidly progressive and/or associated with severe weakness.

Hereditary Neuropathy with Liability to Pressure Palsies

The largest proportion of CMT1 cases are due to variants in *PMP22*. In HNPP (also called tomaculous neuropathy), inadequate production of *PMP22* causes nerves to be more susceptible to trauma or minor compression or entrapment. HNPP patients rarely present symptoms before the second or third decade of life. However, some have reported presentation as early as birth or as late as the seventh decade of life. The prevalence is estimated at 16 persons per 100000, although some authors have indicated a potential for under diagnosis of the disease. An estimated 50% of carriers are asymptomatic and do not display abnormal neurologic findings on clinical examination. HNPP is characterized by repeated focal pressure neuropathies such as carpal tunnel syndrome and peroneal palsy with foot drop and episodes of numbness, muscular weakness, atrophy, and palsies due to minor compression or trauma to the peripheral nerves. The disease is benign with complete recovery occurring within a period of days to months in most cases, although an estimated 15% of patients have residual weakness following an episode. Poor recovery usually involves a history of prolonged pressure on a nerve, but, in these cases, the remaining symptoms are typically mild.

PMP22 is the only gene for which a variant is known to cause HNPP. A large deletion occurs in approximately 80% of patients, and the remaining 20% of patients have single nucleotide variants (SNVs) and small deletions in the *PMP22* gene. One normal allele (due to a 17p11.2 deletion) results in HNPP and a mild phenotype. SNVs in *PMP22* have been associated with a variable spectrum of HNPP phenotypes ranging from mild symptoms to representing a more severe, CMT1-like syndrome. Studies have also reported that the SNV frequency may vary considerably by ethnicity. About 10% to 15% of variant carriers remain clinically asymptomatic, suggesting incomplete penetrance.

Treatment

Currently, there is no therapy to slow the progression of neuropathy for the inherited peripheral neuropathies. A 2015 systematic review of exercise therapies for CMT including 9 studies described in 11 articles reported significant improvements with functional activities and physiological adaptations with exercise. Supportive treatment, if necessary, is generally provided by a multidisciplinary team including neurologists, physiatrists, orthopedic surgeons, and physical and occupational therapists.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic Testing for the Diagnosis of Inherited Peripheral Neuropathies

Policy # 00378

Original Effective Date: 08/21/2013

Current Effective Date: 05/11/2020

Treatment choices are limited to physical therapy, use of orthotics, surgical treatment for skeletal or soft tissue abnormalities, and drug treatment for pain. Avoidance of obesity and drugs associated with nerve damage (eg, vincristine, paclitaxel, cisplatin, isoniazid, nitrofurantoin) is recommended in CMT patients.

Supportive treatment for HNPP can include transient bracing (eg, wrist splint or ankle-foot orthosis), which may become permanent in some cases of foot drop. Prevention of HNPP manifestations can be accomplished by wearing protective padding (eg, elbow or knee pads) to prevent trauma to nerves during activity. Some have reported that vincristine should also be avoided in HNPP patients. Ascorbic acid has been investigated as a treatment for CMT1A based on animal models, but a 2013 trial in humans did not demonstrate significant clinical benefit. Attarian et al (2014) reported results of an exploratory phase 2 randomized, double-blind, placebo-controlled trial of PXT3003, a low-dose combination of 3 approved compounds (baclofen, naltrexone, sorbitol) in 80 adults with CMT1A. The trial demonstrated the safety and tolerability of the drug. Mandel et al (2015) included this randomized controlled trial and 3 other trials (1 of ascorbic acid, 2 of PXT3003) in a meta-analysis.

Molecular Genetic Testing

Multiple laboratories offer individual variant testing for genes involved in hereditary sensory and motor neuropathies, which would typically involve sequencing analysis via Sanger sequencing or next-generation sequencing followed by deletion/duplication analysis (ie, with array comparative genomic hybridization) to detect large deletions or duplications. For the detection of variants in *MFN2*, whole gene or select exome sequence analysis is typically used to identify SNVs, in addition to or followed by deletion or duplication analysis for the detection of large deletions or duplications.

A number of genetic panel tests for the assessment of peripheral neuropathies are commercially available. For example, GeneDx (Gaithersburg, MD) offers an Axonal CMT panel, which uses next-generation sequencing and exon array comparative genomic hybridization. The genes tested include: *AARS*, *BSCL2*, *DNM2*, *DYNC1H1*, *GARS*, *GDAP1*, *GJB1*, *HSPB1*, *HSPB8*, *LMNA*, *LRSAM1*, *MED25*, *MFN2*, *MPZ*, *NEFL*, *PRPS1*, *RAB7A*, and *TRPV4*. InterGenetics (Athens, Greece) offers a next-generation sequencing panel for neuropathy that includes 42 genes involved in CMT, along with other hereditary neuropathies. Fulgent Clinical Diagnostics Lab offers a broader NGS panel for CMT that includes 48 genes associated with CMT and other neuropathies and myopathies.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic Testing for the Diagnosis of Inherited Peripheral Neuropathies

Policy # 00378

Original Effective Date: 08/21/2013

Current Effective Date: 05/11/2020

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratory-developed tests must meet the general regulatory standards of the Clinical Laboratory Improvement Amendments. Genetic testing for the diagnosis of inherited peripheral neuropathies is available under the auspices of the Clinical Laboratory Improvement Amendments. Laboratories that offer laboratory-developed tests must be licensed by the Clinical Laboratory Improvement Amendments for high-complexity testing. To date, the U.S. Food and Drug Administration has chosen not to require any regulatory review of this test.

Rationale/Source

The inherited peripheral neuropathies are a heterogeneous group of diseases that may be inherited in an autosomal dominant, autosomal recessive, or X-linked dominant manner. These diseases can generally be diagnosed based on clinical presentation, nerve conduction studies, and family history. Genetic testing has been used to diagnose specific inherited peripheral neuropathies.

For individuals with suspected inherited motor and sensory peripheral neuropathy who receive testing for genes associated with inherited peripheral neuropathies, the evidence includes case-control and genome-wide association studies. The relevant outcomes are test validity, symptoms, and change in disease status. For the evaluation of hereditary motor and sensory peripheral neuropathies and hereditary neuropathy with liability to pressure palsies, the diagnostic testing yield is likely to be high, particularly when sequential testing is used based on patient phenotype. However, the clinical utility of genetic testing to confirm a diagnosis in a patient with a clinical diagnosis of an inherited peripheral neuropathy is unknown. No direct evidence for improved outcomes with the use of genetic testing for hereditary motor and sensory peripheral neuropathies and hereditary neuropathy with liability to pressure palsies was identified. However, a chain of evidence supports the use of genetic testing to establish a diagnosis in cases of suspected inherited motor or sensory neuropathy, when a diagnosis cannot be made by other methods, to initiate supportive therapies. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

Supplemental Information

Practice Guidelines and Position Statements

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic Testing for the Diagnosis of Inherited Peripheral Neuropathies

Policy # 00378

Original Effective Date: 08/21/2013

Current Effective Date: 05/11/2020

American Academy of Neurology et al

The American Academy of Neurology and 2 other specialty societies (2009) published an evidence-based, tiered approach for the evaluation of distal symmetric polyneuropathy and suspected hereditary neuropathies, which concluded the following (see Table 2).

Table 2. Recommendations on Distal Symmetric Polyneuropathy and Suspected Hereditary Neuropathies

Recommendation	

LOE: level of evidence.

^a Grade A: established as effective, ineffective, or harmful for the given condition in the specified population; grade C: possibly effective, ineffective, or harmful for the given condition in the specified population; grade U: data inadequate or conflicting; given current knowledge.

The American Academy of Neurology website indicates the recommendations were reaffirmed in 2013 and in November 2017 indicated an update is in progress.

American Academy of Family Physicians

The American Academy of Family Physicians (2010) recommended genetic testing for a patient with suspected peripheral neuropathy, if basic blood tests are negative, electrodiagnostic studies suggest an axonal etiology, and diseases such as diabetes, toxic medications, thyroid disease, and vasculitides can be ruled out.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic Testing for the Diagnosis of Inherited Peripheral Neuropathies

Policy # 00378

Original Effective Date: 08/21/2013

Current Effective Date: 05/11/2020

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 3.

Table 3. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
Ongoing			
NCT01193075	Natural History Evaluation of Charcot Marie Tooth Disease (CMT) Type (CMT1B), 2A (CMT2A), 4A (CMT4A), 4C (CMT4C), and Others	5000	Dec 2019
NCT01193088	Genetics of Charcot Marie Tooth Disease (CMT) - Modifiers of CMT1A, New Causes of CMT	1050	Dec 2019

NCT: national clinical trial.

References

1. Blue Cross and Blue Shield Association, Medical Policy Reference Manual, “Genetic Testing for the Diagnosis of Inherited Peripheral Neuropathies”, 2.04.89, February, 2020.
2. Burgunder JM, Schols L, Baets J, et al. EFNS guidelines for the molecular diagnosis of neurogenetic disorders: motoneuron, peripheral nerve and muscle disorders. *Eur J Neurol.* Feb 2011;18(2):207-217. PMID 20500522
3. Alport AR, Sander HW. Clinical approach to peripheral neuropathy: anatomic localization and diagnostic testing. *Continuum (Minneap Minn).* Feb 2012;18(1):13-38. PMID 22810068
4. England JD, Gronseth GS, Franklin G, et al. Practice Parameter: evaluation of distal symmetric polyneuropathy: role of laboratory and genetic testing (an evidence-based review). Report of the American Academy of Neurology, American Association of Neuromuscular and Electrodiagnostic

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic Testing for the Diagnosis of Inherited Peripheral Neuropathies

Policy # 00378

Original Effective Date: 08/21/2013

Current Effective Date: 05/11/2020

- Medicine, and American Academy of Physical Medicine and Rehabilitation. *Neurology*. Jan 13 2009;72(2):185-192. PMID 19056666
5. Saporta AS, Sottile SL, Miller LJ, et al. Charcot-Marie-Tooth disease subtypes and genetic testing strategies. *Ann Neurol*. Jan 2011;69(1):22-33. PMID 21280073
 6. Cornett KM, Menezes MP, Bray P, et al. Phenotypic variability of childhood Charcot-Marie-Tooth disease. *JAMA Neurol*. Jun 01 2016;73(6):645-651. PMID 27043305
 7. Bird TD. Charcot-Marie-Tooth Hereditary Neuropathy Overview. In: Pagon RA, Bird TD, Dolan CR, et al., eds. *GeneReviews*. Seattle, WA: University of Washington; 2016.
 8. Stankiewicz P, Lupski JR. The genomic basis of disease, mechanisms and assays for genomic disorders. *Genome Dyn*. Aug 2006;1:1-16. PMID 18724050
 9. Bird TD. Charcot-Marie-Tooth Neuropathy Type 1. In: Adams M, Ardinger HH, Pagon RA, et al., eds. *GeneReviews*. Seattle, WA: University of Washington; 2015.
 10. Bird TD. Charcot-Marie-Tooth Neuropathy Type 2. In: Adam MP, Ardinger HH, Pagon RA, et al., eds. *GeneReviews*. Seattle, WA: University of Washington; 2016.
 11. Bird TD. Charcot-Marie-Tooth Neuropathy X Type 1. In: Adam MP, Ardinger HH, Pagon RA, et al., eds. *GeneReviews*. Seattle, WA: University of Washington; 2016.
 12. Meretoja P, Silander K, Kalimo H, et al. Epidemiology of hereditary neuropathy with liability to pressure palsies (HNPP) in south western Finland. *Neuromuscul Disord*. Dec 1997;7(8):529-532. PMID 9447611
 13. Celik Y, Kilincer C, Hamamcioglu MK, et al. Hereditary neuropathy with liability to pressure palsies in a Turkish patient (HNPP): a rare cause of entrapment neuropathies in young adults. *Turk Neurosurg*. Jan 2008;18(1):82-84. PMID 18382985
 14. Taioli F, Cabrini I, Cavallaro T, et al. Inherited demyelinating neuropathies with micromutations of peripheral myelin protein 22 gene. *Brain*. Feb 2011;134(Pt 2):608-617. PMID 21252112
 15. Bissar-Tadmouri N, Parman Y, Boutrand L, et al. Mutational analysis and genotype/phenotype correlation in Turkish Charcot-Marie-Tooth Type 1 and HNPP patients. *Clin Genet*. Nov 2000;58(5):396-402. PMID 11140841
 16. Dubourg O, Mouton P, Brice A, et al. Guidelines for diagnosis of hereditary neuropathy with liability to pressure palsies. *Neuromuscul Disord*. Mar 2000;10(3):206-208. PMID 10734269
 17. Sman AD, Hackett D, Fiatarone Singh M, et al. Systematic review of exercise for Charcot-Marie-Tooth disease. *J Peripher Nerv Syst*. Dec 2015;20(4):347-362. PMID 26010435
 18. Pareyson D, Marchesi C. Natural history and treatment of peripheral inherited neuropathies. *Adv Exp Med Biol*. Jan 2009;652:207-224. PMID 20225028

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic Testing for the Diagnosis of Inherited Peripheral Neuropathies

Policy # 00378

Original Effective Date: 08/21/2013

Current Effective Date: 05/11/2020

19. Bird TD. Hereditary neuropathy with liability to pressure palsies. In: Adam MP, Ardinger HH, Pagon RA, et al., eds. Gene Reviews. Seattle, WA: University of Washington; 2014.
20. Lewis RA, McDermott MP, Herrmann DN, et al. High-dosage ascorbic acid treatment in Charcot-Marie-Tooth disease type 1A: results of a randomized, double-masked, controlled trial. *JAMA Neurol.* Aug 2013;70(8):981-987. PMID 23797954
21. Attarian S, Vallat JM, Magy L, et al. An exploratory randomised double-blind and placebo-controlled phase 2 study of a combination of baclofen, naltrexone and sorbitol (PXT3003) in patients with Charcot-Marie-Tooth disease type 1A. *Orphanet J Rare Dis.* Dec 18 2014;9:199. PMID 25519680
22. Mandel J, Bertrand V, Lehert P, et al. A meta-analysis of randomized double-blind clinical trials in CMT1A to assess the change from baseline in CMTNS and ONLS scales after one year of treatment. *Orphanet J Rare Dis.* Jun 13 2015;10:74. PMID 26070802
23. GeneDx. Axonal CMT Panel. <https://www.genedx.com/test-catalog/available-tests/axonal-cmt-panel-1/>.
24. Aretz S, Rautenstrauss B, Timmerman V. Clinical utility gene card for: HMSN/HNPP HMSN types 1, 2, 3, 6 (CMT1,2,4, DSN, CHN, GAN, CCFDN, HNA); HNPP. *Eur J Hum Genet.* Sep 2010;18(9). PMID 20512157
25. Rudnik-Schoneborn S, Tolle D, Senderek J, et al. Diagnostic algorithms in Charcot-Marie-Tooth neuropathies: experiences from a German genetic laboratory on the basis of 1206 index patients. *Clin Genet.* Jan 2016;89(1):34-43. PMID 25850958
26. Gess B, Schirmacher A, Boentert M, et al. Charcot-Marie-Tooth disease: frequency of genetic subtypes in a German neuromuscular center population. *Neuromuscul Disord.* Aug 2013;23(8):647-651. PMID 23743332
27. Ostern R, Fagerheim T, Hjellnes H, et al. Diagnostic laboratory testing for Charcot Marie Tooth disease (CMT): the spectrum of gene defects in Norwegian patients with CMT and its implications for future genetic test strategies. *BMC Med Genet.* 2013;14:94. PMID 24053775
28. Murphy SM, Laura M, Fawcett K, et al. Charcot-Marie-Tooth disease: frequency of genetic subtypes and guidelines for genetic testing. *J Neurol Neurosurg Psychiatry.* Jul 2012;83(7):706-710. PMID 22577229
29. Antoniadi T, Buxton C, Dennis G, et al. Application of targeted multi-gene panel testing for the diagnosis of inherited peripheral neuropathy provides a high diagnostic yield with unexpected phenotype-genotype variability. *BMC Med Genet.* Sep 21 2015;16:84. PMID 26392352

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic Testing for the Diagnosis of Inherited Peripheral Neuropathies

Policy # 00378

Original Effective Date: 08/21/2013

Current Effective Date: 05/11/2020

30. DiVincenzo C, Elzinga CD, Medeiros AC, et al. The allelic spectrum of Charcot-Marie-Tooth disease in over 17,000 individuals with neuropathy. *Mol Genet Genomic Med.* Nov 2014;2(6):522-529. PMID 25614874
31. Sanmaneechai O, Feely S, Scherer SS, et al. Genotype-phenotype characteristics and baseline natural history of heritable neuropathies caused by mutations in the MPZ gene. *Brain.* Nov 2015;138(Pt 11):3180-3192. PMID 26310628
32. Karadima G, Koutsis G, Raftopoulou M, et al. Mutational analysis of Greek patients with suspected hereditary neuropathy with liability to pressure palsies (HNPP): a 15-year experience. *J Peripher Nerv Syst.* Jun 2015;20(2):79-85. PMID 26110377
33. Azhary H, Farooq MU, Bhanushali M, et al. Peripheral neuropathy: differential diagnosis and management. *Am Fam Physician.* Apr 1 2010;81(7):887-892. PMID 20353146

Policy History

Original Effective Date: 08/21/2013

Current Effective Date: 05/11/2020

08/01/2013	Medical Policy Committee review
08/21/2013	Medical Policy Implementation Committee approval. New policy.
08/07/2014	Medical Policy Committee review
08/20/2014	Medical Policy Implementation Committee approval. No change to coverage.
08/06/2015	Medical Policy Committee review
08/19/2015	Medical Policy Implementation Committee approval. No change to coverage.
08/04/2016	Medical Policy Committee review
08/17/2016	Medical Policy Implementation Committee approval. No change to coverage.
01/01/2017	Coding update: Removing ICD-9 Diagnosis Codes
04/06/2017	Medical Policy Committee review
04/19/2017	Medical Policy Implementation Committee approval. Coverage eligibility statements rewritten.
01/01/2018	Coding update
04/05/2018	Medical Policy Committee review
04/18/2018	Medical Policy Implementation Committee approval. No change to coverage.
04/04/2019	Medical Policy Committee review
04/24/2019	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
04/02/2020	Medical Policy Committee review

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic Testing for the Diagnosis of Inherited Peripheral Neuropathies

Policy # 00378

Original Effective Date: 08/21/2013

Current Effective Date: 05/11/2020

04/08/2020 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
Next Scheduled Review Date: 04/2021

Coding

The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)‡, copyright 2019 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	81324, 81325, 81326, 81403, 81404, 81405, 81406, 81448, 81479
HCPCS	No codes
ICD-10 Diagnosis	Z31.430, Z31.440

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic Testing for the Diagnosis of Inherited Peripheral Neuropathies

Policy # 00378

Original Effective Date: 08/21/2013

Current Effective Date: 05/11/2020

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
 1. Consultation with the Blue Cross and Blue Shield Association technology assessment program (TEC) or other nonaffiliated technology evaluation center(s);
 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
 3. Reference to federal regulations.

**Medically Necessary (or “Medical Necessity”) - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. In accordance with nationally accepted standards of medical practice;
- B. Clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and
- C. Not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.

For these purposes, “nationally accepted standards of medical practice” means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Genetic Testing for the Diagnosis of Inherited Peripheral Neuropathies

Policy # 00378

Original Effective Date: 08/21/2013

Current Effective Date: 05/11/2020

‡ Indicated trademarks are the registered trademarks of their respective owners.

NOTICE: If the Patient's health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.