



Louisiana

Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia

Policy # 00048

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Note: BCR-ABL1 Testing in Chronic Myelogenous Leukemia and Acute Lymphoblastic Leukemia is addressed separately in medical policy 00428.

When Services Are Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Children

Based on review of available data, the Company may consider allogeneic or autologous hematopoietic cell transplantation (HCT) to treat childhood acute lymphoblastic leukemia (ALL) in first complete remission but at high risk of relapse to be **eligible for coverage.** **

Based on review of available data, the Company may consider autologous or allogeneic hematopoietic cell transplantation (HCT) to treat childhood acute lymphoblastic leukemia (ALL) in second or greater remission or refractory acute lymphoblastic leukemia (ALL) to be **eligible for coverage.****

Based on review of available data, the Company considers allogeneic hematopoietic cell transplantation (HCT) to treat relapsing acute lymphoblastic leukemia (ALL) after a prior autologous hematopoietic cell transplantation (HCT) to be **eligible for coverage.****

Adults

Based on review of available data, the Company may consider autologous hematopoietic cell transplantation (HCT) to treat adult acute lymphoblastic leukemia (ALL) in first complete remission but at high risk of relapse to be **eligible for coverage.****

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia

Policy # 00048

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

Based on review of available data, the Company may consider allogeneic hematopoietic cell transplantation (HCT) to treat adult acute lymphoblastic leukemia (ALL) in first complete remission for any risk level to be **eligible for coverage.****

Based on review of available data, the Company may consider allogeneic hematopoietic cell transplantation (HCT) to treat adult acute lymphoblastic leukemia (ALL) in second or greater remissions, or in adults with relapsed or refractory acute lymphoblastic leukemia (ALL) to be **eligible for coverage.****

Based on review of available data, the Company may consider reduced-intensity conditioning (RIC) allogeneic hematopoietic cell transplantation (HCT) as a treatment of acute lymphoblastic leukemia (ALL) in patients who are in complete marrow and extramedullary first or second remission, and who, for medical reasons, would be unable to tolerate a standard myeloablative conditioning regimen to be **eligible for coverage.****

Based on review of available data, the Company considers allogeneic hematopoietic cell transplantation (HCT) to treat relapsing acute lymphoblastic leukemia (ALL) after a prior autologous hematopoietic cell transplantation (HCT) to be **eligible for coverage.****

When Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Based on review of available data, the Company considers autologous hematopoietic cell transplantation (HCT) to treat adult acute lymphoblastic leukemia (ALL) in second or greater remission or those with refractory disease to be **investigational.***

Policy Guidelines

Relapse Risk Prognostic Factors

Childhood Acute Lymphoblastic Leukemia

Adverse prognostic factors in children include the following: age younger than 1 year or more than 9 years, male sex, white blood cell count at presentation above 50,000/ μ L, hypodiploidy (<45

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia

Policy # 00048

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

chromosomes), translocation involving chromosomes 9 and 22 (t[9;22]) or *BCR-ABL* fusion, translocation involving chromosomes 4 and 11 (t[4;11]) or *MLL-AF4* fusion, and ProB or T-lineage immunophenotype. Several risk-stratification schema exist, but, in general, the following findings help define children at high-risk of relapse: (1) poor response to initial therapy including poor response to prednisone prophase defined as an absolute blast count of 1000/ μ L or greater, or poor treatment response to induction therapy at 6 weeks with high-risk having 1% or higher minimal residual disease measured by flow cytometry; (2) all children with T-cell phenotype; and (3) patients with either the t(9;22) or t(4;11) regardless of early response measures.

Adult Acute Lymphoblastic Leukemia

Risk factors for relapse are less well-defined in adults, but a patient with any of the following may be considered at high-risk for relapse: age older than 35 years, leukocytosis at presentation of greater than 30000/ μ L (B-cell lineage) or greater than 100000/ μ L (T-cell lineage), “poor prognosis” genetic abnormalities like the Philadelphia chromosome (t[9;22]), extramedullary disease, and time to attain complete remission longer than 4 weeks.

Reduced-Intensity Conditioning

Some patients for whom a conventional myeloablative allogeneic hematopoietic cell transplantation (HCT) could be curative may be considered candidates for reduced-intensity conditioning allogeneic HCT (see Background section). Such patients include those whose age (typically >60 years) or comorbidities (eg, liver or kidney dysfunction, generalized debilitation, prior intensive chemotherapy including autologous or allogeneic HCT, low Karnofsky Performance Status score) preclude the use of a standard myeloablative conditioning regimen.

The ideal allogeneic donors are human leukocyte antigen (HLA)â€identical siblings, matched at the HLA-A, -B, and DR (antigen-D related) loci on each arm of chromosome 6. Related donors mismatched at one locus are also considered suitable donors. A matched, unrelated donor identified through the National Marrow Donor Registry is typically the next option considered. Recently, there has been interest in haploidentical donors, typically a parent or a child of the patient, where usually there is sharing of only three of the six major histocompatibility antigens. Most patients will have such a donor. The risk of morbidity (eg, graft-versus-host disease) may be higher than with HLA-matched donors; however, as medical treatments improve, the risks of graft-versus-host disease with haploidentical donors are approaching those similar to HLA-matched donors.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia

Policy # 00048

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

Background/Overview

Acute Lymphoblastic Leukemia

Childhood Acute Lymphoblastic Leukemia

ALL is the most common cancer diagnosed in children; it represents nearly 25% of cancers in children younger than 15 years. Remission of disease is now typically achieved with pediatric chemotherapy regimens in 98% of children with ALL, with up to 85% long-term survival rates. Survival rates have improved with the identification of effective drugs and combination chemotherapy through large randomized trials, integration of presymptomatic central nervous system prophylaxis, and intensification and risk-based stratification of treatment. The prognosis after the first relapse is related to the length of the original remission. For example, leukemia-free survival is 40% to 50% for children whose first remission was longer than 3 years compared with 10% to 15% for those who relapse less than 3 years after treatment. Thus, hematopoietic cell transplantation (HCT) may be a strong consideration in those with short remissions. At present, the comparative outcomes with autologous or allogeneic HCT (allo-HCT) are unknown.

ALL is a heterogeneous disease with different genetic variations resulting in distinct biologic subtypes. Patients are stratified by certain clinical and genetic risk factors that predict an outcome, with risk-adapted therapy tailoring treatment based on the predicted risk of relapse. Two of the most important factors predictive of risk are patient age and white blood cell count at diagnosis. Certain genetic characteristics of leukemic cells strongly influence prognosis. Clinical and biologic factors predicting clinical outcomes and relapse risk are summarized in the Policy Guidelines section.

Adult ALL

ALL accounts for 20% of acute leukemias in adults. Between 60% and 80% of adults with ALL can be expected to achieve a complete response after induction chemotherapy; however, only 35% to 40% can be expected to survive 2 years. Differences in the frequency of genetic abnormalities that characterize adult ALL versus childhood ALL help, in part, explain differences in outcomes between the 2 groups. For example, the “good prognosis” genetic abnormalities, such as hyperdiploidy and translocation of chromosomes 12 and 21, are seen much less commonly in adult ALL, whereas they are some of the most common in childhood ALL. Conversely, “poor prognosis” genetic abnormalities such as the Philadelphia chromosome (translocation of chromosomes 9 and 22) are seen in 25% to 30% of adult ALL but infrequently in childhood ALL. Other adverse prognostic

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia

Policy # 00048

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

factors in adult ALL include age greater than 35 years, poor performance status, male sex, and leukocytosis at presentation of greater than 30000/ μ L (B-cell lineage) or greater than 100000/ μ L (T-cell lineage).

Hematopoietic Cell Transplantation

HCT) is a procedure in which hematopoietic stem cells are intravenously infused to restore bone marrow and immune function in cancer patients who receive bone marrow-toxic doses of cytotoxic drugs with or without whole-body radiotherapy. Hematopoietic stem cells may be obtained from the transplant recipient (autologous HCT) or a donor (allo-HCT). They can be harvested from bone marrow, peripheral blood, or umbilical cord blood shortly after delivery of neonates.

Immunologic compatibility between infused hematopoietic stem cells and the recipient is not an issue in autologous HCT. In allogeneic stem cell transplantation, immunologic compatibility between donor and patient is a critical factor for achieving a successful outcome. Compatibility is established by typing of human leukocyte antigens (HLA) using cellular, serologic, or molecular techniques. HLA refers to the gene complex expressed at the HLA-A, -B, and -DR (antigen-D related) loci on each arm of chromosome 6. An acceptable donor will match the patient at all or most of the HLA loci.

Conditioning for Hematopoietic Cell Transplantation

Conventional Conditioning

The conventional (“classical”) practice of allo-HCT involves administration of cytotoxic agents (e.g., cyclophosphamide, busulfan) with or without total body irradiation at doses sufficient to cause bone marrow ablation in the recipient. The beneficial treatment effect of this procedure is due to a combination of the initial eradication of malignant cells and subsequent graft-versus-malignancy effect mediated by non-self-immunologic effector cells. While the slower graft-versus-malignancy effect is considered the potentially curative component, it may be overwhelmed by existing disease in the absence of pre transplant conditioning. Intense conditioning regimens are limited to patients who are sufficiently medically fit to tolerate substantial adverse effects. These include opportunistic infections secondary to loss of endogenous bone marrow function and organ damage or failure caused by cytotoxic drugs. Subsequent to graft infusion in allo-HCT, immunosuppressant drugs are required to minimize graft rejection and graft-versus-host disease, which increases susceptibility to opportunistic infections.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia

Policy # 00048

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

The success of autologous HCT is predicated on the potential of cytotoxic chemotherapy, with or without radiotherapy, to eradicate cancerous cells from the blood and bone marrow. This permits subsequent engraftment and repopulation of the bone marrow with presumably normal hematopoietic stem cells obtained from the patient before undergoing bone marrow ablation. Therefore, autologous HCT is typically performed as consolidation therapy when the patient's disease is in complete remission. Patients who undergo autologous HCT are also susceptible to chemotherapy-related toxicities and opportunistic infections before engraftment, but not graft-versus-host disease.

Reduced-Intensity Conditioning Allogeneic Hematopoietic Cell Transplantation

RIC refers to the pre transplant use of lower doses of cytotoxic drugs or less intense regimens of radiotherapy than are used in traditional full-dose myeloablative conditioning treatments. Although the definition of RIC is variable, with numerous versions employed, all regimens seek to balance the competing effects of relapse due to residual disease and non-relapse mortality. The goal of RIC is to reduce disease burden and to minimize associated treatment-related morbidity and non-relapse mortality in the period during which the beneficial graft-versus-malignancy effect of allogeneic transplantation develops. RIC regimens range from nearly total myeloablative to minimally myeloablative with lymphoablation, with intensity tailored to specific diseases and patient condition. Patients who undergo RIC with allo-HCT initially demonstrate donor cell engraftment and bone marrow mixed chimerism. Most will subsequently convert to full-donor chimerism. In this review, the term reduced-intensity conditioning will refer to all conditioning regimens intended to be nonmyeloablative.

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

The FDA regulates human cells and tissues intended for implantation, transplantation, or infusion through the Center for Biologics Evaluation and Research, under Code of Federal Regulation, Title 21, parts 1270 and 1271. Hematopoietic stem cells are included in these regulations.

Rationale/Source

Acute lymphoblastic leukemia (ALL) is a heterogeneous disease with different genetic variations resulting in distinct biologic subtypes. Patients are stratified to risk-adapted therapy according to

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia

Policy # 00048

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

certain clinical and genetic risk factors that predict an outcome. Therapy may include hematopoietic cell transplantation (HCT).

For individuals who have childhood ALL in first complete remission (CR1) at high-risk of relapse, remission, or refractory ALL who receive autologous HCT, the evidence includes RCTs and systematic reviews. The relevant outcomes are overall survival (OS), disease-specific survival (DSS), and treatment-related mortality (TRM) and morbidity. For children with high-risk ALL in CR1 or with relapsed ALL, studies have suggested that HCT is associated with fewer relapses but higher death rates due to treatment-related toxicity. However, for a subset of high-risk patients in second complete remission or beyond or with relapsed disease, autologous HCT is a treatment option. This conclusion is further supported by an evidence-based systematic review and position statement from the American Society for Blood and Marrow Transplantation. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have childhood ALL in CR1 at high-risk of relapse, remission, or refractory ALL who receive allogeneic HCT (allo-HCT), the evidence includes RCTs and systematic reviews. The relevant outcomes are OS, DSS, and TRM and morbidity. For children with high-risk ALL in CR1 or with relapsed ALL, studies have suggested that allo-HCT is associated with fewer relapses but higher death rates due to treatment-related toxicity. However, for a subset of high-risk patients in second complete remission or beyond or with relapsed disease, allo-HCT is a treatment option. This conclusion is further supported by an evidence-based systematic review and position statement from the American Society for Blood and Marrow Transplantation. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have adult ALL in CR1, subsequent remission, or refractory ALL who receive autologous HCT, the evidence includes RCTs and systematic reviews. The relevant outcomes are OS, DSS, and TRM and morbidity. Current evidence supports the use of autologous HCT for adults with high-risk ALL in CR1, whose health status is sufficient to tolerate the procedure. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have adult ALL in CR1 or subsequent remission or refractory ALL who receive allo-HCT, the evidence includes RCTs and systematic reviews. The relevant outcomes are OS,

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia

Policy # 00048

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

DSS, and TRM and morbidity. Current evidence supports the use of myeloablative allo-HCT for adults with any risk level ALL, whose health status is sufficient to tolerate the procedure. Reduced-intensity conditioning allo-HCT may be considered for patients who demonstrate complete marrow and extramedullary first or second remission and who could be expected to benefit from a myeloablative allo-HCT, but for medical reasons would not tolerate a myeloablative conditioning regimen. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have relapsed after a prior autologous HCT for ALL who receive allo-HCT, the evidence includes case series and systematic reviews. The relevant outcomes are OS, DSS, and TRM and morbidity. Evidence reviews have identified only small case series with short-term follow-up, which was considered inadequate evidence of benefit. The evidence is insufficient to determine the effects of the technology on health outcomes.

Allo-HCT after failed autologous HCT has been shown to be of clinical benefit in other hematologic malignancies and is potentially curative. In addition, clinical input has supported the use of allo-HCT to treat relapsing ALL after a failed, prior autologous HCT, particularly with reduced-intensity conditioning regimens, in adults or children. Thus, these indications may be considered medically necessary.

Supplemental Information

Clinical Input From Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

In response to requests, input was received from 1 medical society, 2 academic medical centers, and 3 physicians from Blue Distinction Centers while this policy was under review in 2013. In general, input supported most existing policy statements. However, most reviewers disagreed that allogeneic hematopoietic cell transplantation (allo-HCT) is considered investigational to treat relapsing acute lymphoblastic leukemia (ALL) after a prior autologous HCT in either children or adults. Given a scarcity of evidence on this topic, with no substantial trials likely to be forthcoming,

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia

Policy # 00048

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

and that reduced-intensity conditioning allo- HCT is considered medically necessary to treat ALL in second or greater remission or relapsed or refractory ALL, the policy statements were revised to medical necessity for this indication in children and adults.

Practice Guidelines and Position Statements

National Comprehensive Cancer Network

Current National Comprehensive Cancer Network guidelines (v.2.2019) for ALL indicate allo-HCT is appropriate for consolidation treatment of most poor risk (eg, the Philadelphia chromosome-positive, relapsed, or refractory) patients with ALL. The guidelines state that for appropriately fit older adults with ALL who are achieving remission, “consideration of autologous or reduced-intensity allogeneic stem cell transplantation may be appropriate.” In addition, the guidelines note that chronologic age is not a good surrogate for fitness for therapy and that patient should be evaluated on an individual basis.

Current National Comprehensive Cancer Network guidelines (v.2.2020) for pediatric ALL say that "Allogeneic HSCT has demonstrated improved clinical outcomes in pediatric ALL patients with evidence of certain high-risk features and/or persistent disease. In addition, survival rates appear to be comparable regardless of the stem cell source (matched related, matched unrelated, cord blood, or haploidentical donor)."

American Society for Blood and Marrow Transplantation

The guidelines from the American Society for Blood and Marrow Transplantation (2015) were published on indications for autologous and allo- HCT. Recommendations were intended to describe the current consensus on the use of HCT in and out of the clinical trial setting. Recommendations on ALL are listed in Table 1.

Table 1. Guidelines for Autologous and Allogeneic HCT in ALL

Indication	Children (Age <18 Years)		Adults (Age ≥18 Years)	
	Allogeneic HCT	Autologous HCT	Allogeneic HCT	Autologous HCT

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia

Policy # 00048

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

First complete response, standard-risk	N	N	S	C
First complete response, high-risk	S	N	S	N
Second complete response	S	N	S	C
At least third complete response	C	N	C	N
Not in remission	C	N	C	N

ALL: acute lymphoblastic leukemia; C: clinical evidence available; HCT: hematopoietic cell transplantation; N: not generally recommended; S: standard of care.

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is a national coverage determination for stem cell transplantation (110.23; formerly 110.81), portions of which are highlighted below:

Nationally Covered Indications

“I. Allogeneic Hematopoietic Stem Cell Transplantation (HSCT)

- a) Effective ... 1978, for the treatment of leukemia, leukemia in remission, or aplastic anemia when it is reasonable and necessary,
- b) Effective ... 1985, for the treatment of severe combined immunodeficiency disease (SCID) and for the treatment of Wiskott-Aldrich syndrome.
- c) Effective ... 2010, for the treatment of Myelodysplastic Syndromes (MDS) pursuant to Coverage with Evidence Development (CED) in the context of a Medicare-approved, prospective clinical study...
- d) Effective for claims with dates of service on or after January 27, 2016, allogeneic HSCT for multiple myeloma is covered by Medicare only for beneficiaries with Durie-Salmon Stage II or III

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia

Policy # 00048

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

multiple myeloma, or International Staging System (ISS) Stage II or Stage III multiple myeloma, and participating in an approved prospective clinical study that meets the criteria below. There must be appropriate statistical techniques to control for selection bias and confounding by age, duration of diagnosis, disease classification, International Myeloma Working Group (IMWG) classification, ISS stage, comorbid conditions, type of preparative/conditioning regimen, graft vs. host disease (GVHD) prophylaxis, donor type and cell source...

e) Effective for claims with dates of service on or after January 27, 2016, allogeneic HSCT for myelofibrosis (MF) is covered by Medicare only for beneficiaries with Dynamic International Prognostic Scoring System (DIPSSplus) intermediate-2 or High primary or secondary MF and participating in an approved prospective clinical study. All Medicare-approved studies must use appropriate statistical techniques in the analysis to control for selection bias and potential confounding by age, duration of diagnosis, disease classification, DIPSSplus score, comorbid conditions, type of preparative/conditioning regimen, graft vs. host disease (GVHD) prophylaxis, donor type and cell source...

f) Effective for claims with dates of service on or after January 27, 2016, allogeneic HSCT for sickle cell disease (SCD) is covered by Medicare only for beneficiaries with severe, symptomatic SCD who participate in an approved prospective clinical study....

II. Autologous Stem Cell Transplantation (AuSCT)

a) Effective ... 1989, AuSCT is considered reasonable and necessary ... for the following conditions and is covered under Medicare for patients with:

1. Acute leukemia in remission who have a high probability of relapse and who have no human leukocyte antigens (HLA)-matched;
2. Resistant non-Hodgkin's lymphomas or those presenting with poor prognostic features following an initial response;
3. Recurrent or refractory neuroblastoma; or,
4. Advanced Hodgkin's disease who have failed conventional therapy and have no HLA-matched donor.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia

Policy # 00048

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

b) Effective ... 2000, single AuSCT is only covered for Durie-Salmon Stage II or III patients that fit the following requirements:

- Newly diagnosed or responsive multiple myeloma. This includes those patients with previously untreated disease, those with at least a partial response to prior chemotherapy (defined as a 50% decrease either in measurable paraprotein [serum and/or urine] or in bone marrow infiltration, sustained for at least 1 month), and those in responsive relapse; and
- Adequate cardiac, renal, pulmonary, and hepatic function.

c) Effective ... 2005, when recognized clinical risk factors are employed to select patients for transplantation, high dose melphalan (HDM) together with AuSCT is reasonable and necessary for Medicare beneficiaries of any age group with primary amyloid light chain (AL) amyloidosis who meet the following criteria:

- Amyloid deposition in 2 or fewer organs; and,
- Cardiac left ventricular ejection fraction (EF) greater than 45%.”

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 2.

Table 2. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT02042690	Phase III Study to Compare Haplo-identical HSCT Versus Chemotherapy in First Remission for Standard-risk Adult Acute Lymphoblastic Leukemia	300	Apr 2019 (completed)
NCT01597778	A Multi-Center, Phase III, Randomized Trial of Reduced Intensity Conditioning (RIC) and Transplantation of Double Unrelated Umbilical Cord Blood (dUCB) Versus HLA-Haploidentical	410	Jun 2020

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia

Policy # 00048

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

	Related Bone Marrow (Haplo) for Patients With Hematologic Malignancies		
NCT01700946	A Phase II Study of Therapy for Pediatric Relapsed or Refractory Precursor B-Cell Acute Lymphoblastic Leukemia and Lymphoma	40	Apr 2021
NCT03314974	Myeloablative Allogeneic Hematopoietic Cell Transplantation Using a Related or Unrelated Donor for the Treatment of Hematological Diseases	40	Nov 2024
NCT01949129	Allogeneic Stem Cell Transplantation for Children and Adolescents With Acute Lymphoblastic Leukaemia	1000	Apr 2026

NCT: national clinical trial.

References

1. Blue Cross and Blue Shield Association, Medical Policy Reference Manual, “Hematopoietic Stem-Cell Transplantation for Acute Lymphoblastic Leukemia”, 8.01.32, February 2020.
2. National Cancer Institute. Childhood Acute Lymphoblastic Leukemia Treatment (PDQ)Health Professional Version. 2017; https://www.cancer.gov/types/leukemia/hp/child-all-treatment-pdq#section/_1.
3. Pieters R, Carroll WL. Biology and treatment of acute lymphoblastic leukemia. *Pediatr Clin North Am.* Feb 2008;55(1):1-20, ix. PMID 18242313
4. Carroll WL, Bhojwani D, Min DJ, et al. Pediatric acute lymphoblastic leukemia. *Hematology Am Soc Hematol Educ Program.* 2003:102-131. PMID 14633779
5. National Cancer Institute. Adult Acute Lymphoblastic Leukemia Treatment (PDQ)Health Professional Version. 2017; <https://www.cancer.gov/types/leukemia/hp/adult-all-treatment-pdq>.
6. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). 1987 TEC Evaluations. Page 243.
7. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). 1990 TEC Evaluations. Page 254.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia

Policy # 00048

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

8. Harrison G, Richards S, Lawson S, et al. Comparison of allogeneic transplant versus chemotherapy for relapsed childhood acute lymphoblastic leukaemia in the MRC UKALL R1 trial. MRC Childhood Leukaemia Working Party. *Ann Oncol.* Aug 2000;11(8):999-1006. PMID 11038037
9. Lawson SE, Harrison G, Richards S, et al. The UK experience in treating relapsed childhood acute lymphoblastic leukaemia: a report on the Medical Research Council UKALLR1 study. *Br J Haematol.* Mar 2000;108(3):531-543. PMID 10759711
10. Wheeler KA, Richards SM, Bailey CC, et al. Bone marrow transplantation versus chemotherapy in the treatment of very high-risk childhood acute lymphoblastic leukemia in first remission: results from Medical Research Council UKALL X and XI. *Blood.* Oct 1 2000;96(7):2412-2418. PMID 11001892
11. Ribera JM, Ortega JJ, Oriol A, et al. Comparison of intensive chemotherapy, allogeneic, or autologous stem-cell transplantation as postremission treatment for children with very high risk acute lymphoblastic leukemia: PETHEMA ALL-93 Trial. *J Clin Oncol.* Jan 1 2007;25(1):16-24. PMID 17194902
12. Oliansky DM, Camitta B, Gaynon P, et al. Role of cytotoxic therapy with hematopoietic stem cell transplantation in the treatment of pediatric acute lymphoblastic leukemia: update of the 2005 evidence-based review. *Biol Blood Marrow Transplant.* Apr 2012;18(4):505-522. PMID 22209888
13. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). High-dose chemotherapy with autologous stem-cell support in the treatment of adult acute lymphoblastic leukemia. *TEC Assessments.* 1997;Volume 12:Tab 25.
14. Yanada M, Matsuo K, Suzuki T, et al. Allogeneic hematopoietic stem cell transplantation as part of postremission therapy improves survival for adult patients with high-risk acute lymphoblastic leukemia: a meta-analysis. *Cancer.* Jun 15 2006;106(12):2657-2663. PMID 16703597
15. Hahn T, Wall D, Camitta B, et al. The role of cytotoxic therapy with hematopoietic stem cell transplantation in the therapy of acute lymphoblastic leukemia in adults: an evidence-based review. *Biol Blood Marrow Transplant.* Jan 2006;12(1):1-30. PMID 16399566
16. Attal M, Blaise D, Marit G, et al. Consolidation treatment of adult acute lymphoblastic leukemia: a prospective, randomized trial comparing allogeneic versus autologous bone marrow transplantation and testing the impact of recombinant interleukin-2 after autologous bone marrow transplantation. *BGMT Group. Blood.* Aug 15 1995;86(4):1619-1628. PMID 7632972

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia

Policy # 00048

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

17. Dombret H, Gabert J, Boiron JM, et al. Outcome of treatment in adults with Philadelphia chromosome-positive acute lymphoblastic leukemia--results of the prospective multicenter LALA-94 trial. *Blood*. Oct 1 2002;100(7):2357-2366. PMID 12239143
18. Hunault M, Harousseau JL, Delain M, et al. Better outcome of adult acute lymphoblastic leukemia after early genotypical allogeneic bone marrow transplantation (BMT) than after late high-dose therapy and autologous BMT: a GOELAMS trial. *Blood*. Nov 15 2004;104(10):3028-3037. PMID 15256423
19. Gupta V, Richards S, Rowe J, et al. Allogeneic, but not autologous, hematopoietic cell transplantation improves survival only among younger adults with acute lymphoblastic leukemia in first remission: an individual patient data meta-analysis. *Blood*. Jan 10 2013;121(2):339-350. PMID 23165481
20. Ribera JM, Oriol A, Bethencourt C, et al. Comparison of intensive chemotherapy, allogeneic or autologous stem cell transplantation as postremission treatment for adult patients with high-risk acute lymphoblastic leukemia. Results of the PETHEMA ALL-93 trial. *Haematologica*. Oct 2005;90(10):1346-1356. PMID 16219571
21. Goldstone AH, Richards SM, Lazarus HM, et al. In adults with standard-risk acute lymphoblastic leukemia, the greatest benefit is achieved from a matched sibling allogeneic transplantation in first complete remission, and an autologous transplantation is less effective than conventional consolidation/maintenance chemotherapy in all patients: final results of the International ALL Trial (MRC UKALL XII/ECOG E2993). *Blood*. Feb 15 2008;111(4):1827-1833. PMID 18048644
22. Fielding AK, Rowe JM, Richards SM, et al. Prospective outcome data on 267 unselected adult patients with Philadelphia chromosome-positive acute lymphoblastic leukemia confirms superiority of allogeneic transplantation over chemotherapy in the pre-imatinib era: results from the International ALL Trial MRC UKALLXII/ECOG2993. *Blood*. May 7 2009;113(19):4489-4496. PMID 19244158
23. Cornelissen JJ, van der Holt B, Verhoef GE, et al. Myeloablative allogeneic versus autologous stem cell transplantation in adult patients with acute lymphoblastic leukemia in first remission: a prospective sibling donor versus no-donor comparison. *Blood*. Feb 5 2009;113(6):1375-1382. PMID 18988865
24. Giebel S, Labopin M, Socie G, et al. Improving results of allogeneic hematopoietic cell transplantation for adults with acute lymphoblastic leukemia in first complete remission: an analysis from the Acute Leukemia Working Party of the European Society for Blood and Marrow Transplantation. *Haematologica*. Jan 2017;102(1):139-149. PMID 27686376

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia

Policy # 00048

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

25. Dinmohamed AG, Szabo A, van der Mark M, et al. Improved survival in adult patients with acute lymphoblastic leukemia in the Netherlands: a population-based study on treatment, trial participation and survival. *Leukemia*. Feb 2016;30(2):310-317. PMID 26286115
26. Pavlu J, Labopin M, Zoellner AK, et al. Allogeneic hematopoietic cell transplantation for primary refractory acute lymphoblastic leukemia: A report from the Acute Leukemia Working Party of the EBMT. *Cancer*. Jun 01 2017;123(11):1965-1970. PMID 28211939
27. Pidala J, Djulbegovic B, Anasetti C, et al. Allogeneic hematopoietic cell transplantation for adult acute lymphoblastic leukemia (ALL) in first complete remission. *Cochrane Database Syst Rev*. Oct 5 2011(10):CD008818. PMID 21975786
28. Abdul Wahid SF, Ismail NA, Mohd-Idris MR, et al. Comparison of reduced-intensity and myeloablative conditioning regimens for allogeneic hematopoietic stem cell transplantation in patients with acute myeloid leukemia and acute lymphoblastic leukemia: a meta-analysis. *Stem Cells Dev*. Nov 1 2014;23(21):2535-2552. PMID 25072307
29. Gutierrez-Aguirre CH, Gomez-Almaguer D, Cantu-Rodriguez OG, et al. Non-myeloablative stem cell transplantation in patients with relapsed acute lymphoblastic leukemia: results of a multicenter study. *Bone Marrow Transplant*. Sep 2007;40(6):535-539. PMID 17618317
30. Mohty M, Labopin M, Tabrizzi R, et al. Reduced-intensity conditioning allogeneic stem cell transplantation for adult patients with acute lymphoblastic leukemia: a retrospective study from the European Group for Blood and Marrow Transplantation. *Haematologica*. Feb 2008;93(2):303-306. PMID 18245655
31. Cho BS, Lee S, Kim YJ, et al. Reduced-intensity conditioning allogeneic stem cell transplantation is a potential therapeutic approach for adults with high-risk acute lymphoblastic leukemia in remission: results of a prospective phase 2 study. *Leukemia*. Oct 2009;23(10):1763-1770. PMID 19440217
32. Pulsipher MA, Boucher KM, Wall D, et al. Reduced-intensity allogeneic transplantation in pediatric patients ineligible for myeloablative therapy: results of the Pediatric Blood and Marrow Transplant Consortium Study ONC0313. *Blood*. Aug 13 2009;114(7):1429-1436. PMID 19528536
33. Rosko A, Wang HL, de Lima M, et al. Reduced-intensity conditioned allograft yields favorable survival for older adults with B-cell acute lymphoblastic leukemia. *Am J Hematol*. Jan 2017;92(1):42-49. PMID 27712033
34. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Salvage high-dose chemotherapy with allogeneic stem-cell support for relapse or incomplete remission

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia

Policy # 00048

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

following high-dose chemotherapy with autologous stem-cell transplantation for hematologic malignancies. TEC Assessments. 2000;Volume 15:Tab 9.

- 35. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Acute lymphoblastic leukemia. Version 2.2019. https://www.nccn.org/professionals/physician_gls/pdf/all.pdf.
- 36. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Pediatric Acute Lymphoblastic Leukemia. Version 2.2020. https://www.nccn.org/professionals/physician_gls/pdf/ped_all.pdf.
- 37. Majhail NS, Farnia SH, Carpenter PA, et al. Indications for autologous and allogeneic hematopoietic cell transplantation: guidelines from the American Society for Blood and Marrow Transplantation. Biol Blood Marrow Transplant. Nov 2015;21(11):1863-1869. PMID 26256941
- 38. Centers for Medicare and Medicaid Services. National Coverage Determination (NCD) for Stem Cell Transplantation Formerly 110.8.1 (110.23). 2016; [https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=366&ncdver=1&MCDIndexType=2&mcctypename=Potential+National+Coverage+Determination+\(NCD\)+Topics&bc=AAAAQAAAAAA&](https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=366&ncdver=1&MCDIndexType=2&mcctypename=Potential+National+Coverage+Determination+(NCD)+Topics&bc=AAAAQAAAAAA&).

Policy History

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

- 12/06/2001 Medical Policy Committee review
- 03/25/2002 Managed Care Advisory Council approval
- 06/24/2002 Format revision. Policy addresses only Acute Lymphocytic Leukemia. Replaces High Dose Chemotherapy with Hematopoietic Stem Cell Support.
- 03/31/2004 Medical Director review
- 04/20/2004 Medical Policy Committee review. Format revision. No changes to coverage eligibility.
- 04/26/2004 Managed Care Advisory Council approval
- 04/05/2005 Medical Director review
- 04/19/2005 Medical Policy Committee review. Format revisions only. Coverage eligibility unchanged. 05/23/2005 Managed Care Advisory Council approval
- 09/06/2006 Medical Director review

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia

Policy # 00048

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

09/20/2006	Medical Policy Committee approval. Format changes only. Coverage eligibility unchanged.
07/11/2007	Medical Director review
07/18/2007	Medical Policy Committee approval. Format revision only. Coverage eligibility unchanged.
07/02/2008	Medical Director review
07/16/2008	Medical Policy Committee approval. No change to coverage eligibility.
07/02/2009	Medical Director review
07/22/2009	Medical Policy Committee approval. Title changed to Hematopoietic Stem-Cell Transplantation for Acute Lymphoblastic Leukemia. Revised policy based on updated research.
07/01/2010	Medical Policy Committee approval.
07/21/2010	Medical Policy Implementation Committee approval. Policy statement regarding treatment of adult ALL in first complete remission but at high risk of relapse split to address allogeneic and autologous transplant separately.
07/07/2011	Medical Policy Committee approval.
07/20/2011	Medical Policy Implementation Committee approval. No change to coverage eligibility.
06/28/2012	Medical Policy Committee review.
07/27/2012	Medical Policy Implementation Committee approval. No change to coverage eligibility.
03/04/2013	Coding updated
08/01/2013	Medical Policy Committee review
08/21/2013	Medical Policy Implementation Committee approval. Revised coverage statements so that allogeneic HSCT may be considered eligible for coverage following a failed autologous HSCT in children or adult patients.
09/04/2014	Medical Policy Committee review
09/17/2014	Medical Policy Implementation Committee approval. No change to coverage.
09/03/2015	Medical Policy Committee review
09/23/2015	Medical Policy Implementation Committee approval. No change to coverage.
09/08/2016	Medical Policy Committee review
09/21/2016	Medical Policy Implementation Committee approval. No change to coverage.
01/01/2017	Coding update: Removing ICD-9 Diagnosis Codes
09/07/2017	Medical Policy Committee review

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia

Policy # 00048

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

- 09/20/2017 Medical Policy Implementation Committee approval. The word stem was removed from title and body of policy.
- 09/06/2018 Medical Policy Committee review
- 09/19/2018 Medical Policy Implementation Committee approval. No change to coverage.
- 09/05/2019 Medical Policy Committee review
- 09/11/2019 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
- 09/03/2020 Medical Policy Committee review
- 09/09/2020 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

Next Scheduled Review Date: 09/2021

Coding

The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)‡, copyright 2019 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia

Policy # 00048

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	38204, 38205, 38206, 38207, 38208, 38209, 38210, 38211, 38212, 38213, 38214, 38215, 38230, 38240, 38241, 38242, 38243
HCPCS	S2140, S2142, S2150
ICD-10 Diagnosis	C91.00-C91.02

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
 - 1. Consultation with the Blue Cross and Blue Shield Association technology assessment program (TEC) or other nonaffiliated technology evaluation center(s);
 - 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
 - 3. Reference to federal regulations.

**Medically Necessary (or “Medical Necessity”) - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. In accordance with nationally accepted standards of medical practice;

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia

Policy # 00048

Original Effective Date: 01/28/2002

Current Effective Date: 10/12/2020

- B. Clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and
- C. Not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.

For these purposes, "nationally accepted standards of medical practice" means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

‡ Indicated trademarks are the registered trademarks of their respective owners.

NOTICE: If the Patient's health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.