



Louisiana

Image-Guided Minimally Invasive Decompression for Spinal Stenosis

Policy # 00278

Original Effective Date: 11/16/2010

Current Effective Date: 03/08/2020

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the “Company”), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Based on review of available data, the Company considers image-guided minimally invasive spinal decompression to be **investigational**.*

Background/Overview

Spinal Stenosis

In spinal stenosis, the space around the spinal cord narrows, compressing the spinal cord and its nerve roots. The goal of surgical treatment is to “decompress” the spinal cord and/or nerve roots.

The most common symptoms of lumbar spinal stenosis (LSS) are back pain with neurogenic claudication (ie, pain, numbness, weakness) in the legs that worsens with standing or walking and is alleviated by sitting or leaning forward. Compression of neural elements generally occurs from a combination of degenerative changes, including ligamentum flavum hypertrophy, bulging of the intervertebral disc, and facet thickening with arthropathy. Spinal stenosis is often linked to age-related changes in disc height and arthritis of the facet joints. LSS is among the most common reasons for back surgery and the most common reason for lumbar spine surgery in adults over the age of 65.

The most common symptoms of cervical/thoracic spinal stenosis are neck pain and radiculopathy of the shoulder and arm. The most common cause of cervical radiculopathy is degenerative changes, including disc herniation.

©2019 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Image-Guided Minimally Invasive Decompression for Spinal Stenosis

Policy # 00278

Original Effective Date: 11/16/2010

Current Effective Date: 03/08/2020

Treatment

Conventional Posterior Decompression Surgery

For patients with LSS, surgical laminectomy has established benefits in reducing pain and improving quality of life.

For patients with cervical or thoracic stenosis, surgical treatment includes discectomy or foraminal decompression.

A systematic review by Chou et al (2009) assessed surgery for back pain; it was commissioned by the American Pain Society and conducted by an evidence-based center. Four higher quality randomized trials were reviewed; they compared surgery with nonsurgical therapy for spinal stenosis, including two studies from the multicenter Spine Patient Outcomes Research Trial that evaluated laminectomy for spinal stenosis (specifically with or without degenerative spondylolisthesis). All 4 studies found that initial decompressive surgery (laminectomy) was slightly to moderately superior to initial nonsurgical therapy (eg, average 8- to 18-point differences on the 36-Item Short-Form Health Survey and Oswestry Disability Index). However, there was insufficient evidence to determine the optimal adjunctive surgical methods for laminectomy (ie, with or without fusion, instrumented vs non instrumented fusion) in patients with or without degenerative spondylolisthesis. Spine Patient Outcomes Research Trial continues to be referenced as the highest quality evidence published on decompressive surgery.

Less invasive surgical procedures include open laminotomy and microendoscopic laminotomy. In general, the literature comparing surgical procedures is limited. The literature has suggested that less invasive surgical decompression may reduce perioperative morbidity without impairing long-term outcomes when performed in appropriately selected patients. Posterior decompressive surgical procedures include: decompressive laminectomy, hemilaminotomy and laminotomy, and microendoscopic decompressive laminotomy.

Decompressive laminectomy, the classic treatment for LSS, unroofs the spinal canal by extensive resection of posterior spinal elements, including the lamina, spinous processes, portions of the facet joints, ligamentum flavum, and the interspinous ligaments. Wide muscular dissection and retraction is needed to achieve adequate surgical visualization. The extensive resection and injury to the posterior spine and supporting musculature can lead to instability with significant morbidity, both

©2019 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Image-Guided Minimally Invasive Decompression for Spinal Stenosis

Policy # 00278

Original Effective Date: 11/16/2010

Current Effective Date: 03/08/2020

postoperatively and longer term. Spinal fusion, performed at the same time as laminectomy or after symptoms have developed, may be required to reduce resultant instability. Laminectomy may also be used for extensive multilevel decompression.

Hemilaminotomy and laminotomy, sometimes termed *laminoforaminotomy*, are less invasive than laminectomy. These procedures focus on the interlaminar space, where most of the pathologic changes are concentrated, minimizing resection of the stabilizing posterior spine. A laminotomy typically removes the inferior aspect of the cranial lamina, superior aspect of the subjacent lamina, ligamentum flavum, and the medial aspect of the facet joint. Unlike laminectomy, laminotomy does not disrupt the facet joints, supra- and interspinous ligaments, a major portion of the lamina, or the muscular attachments. Muscular dissection and retraction are required to achieve adequate surgical visualization.

Microendoscopic decompressive laminotomy, similar to laminotomy, uses endoscopic visualization. The position of the tubular working channel is confirmed by fluoroscopic guidance, and serial dilators are used to dilate the musculature and expand the fascia. For microendoscopic decompressive laminotomy, an endoscopic curette, rongeur, and drill are used for the laminotomy, facetectomy, and foraminotomy. The working channel may be repositioned from a single incision for multilevel and bilateral dissections.

Image-Guided Minimally Invasive Lumbar Decompression

Posterior decompression for LSS has been evolving toward increasingly minimally invasive procedures in an attempt to reduce postoperative morbidity and spinal instability. Unlike conventional surgical decompression, the percutaneous mild^{®‡} decompressive procedure is performed solely under fluoroscopic guidance (eg, without endoscopic or microscopic visualization of the work area). This procedure is indicated for central stenosis only, without the capability of addressing nerve root compression or disc herniation, should either be required.

Percutaneous image-guided minimally invasive lumbar decompression using a specially designed tool kit (mild) has been proposed as an ultra-minimally invasive treatment of central LSS. In this procedure, the epidural space is filled with contrast medium under fluoroscopic guidance. Using a 6-gauge cannula clamped in place with a back plate, single-use tools (portal cannula, surgical guide, bone rongeur, tissue sculpter, trocar) are used to resect thickened ligamentum flavum and small pieces of lamina. The tissue and bone sculpting is conducted entirely under fluoroscopic guidance,

©2019 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Image-Guided Minimally Invasive Decompression for Spinal Stenosis

Policy # 00278

Original Effective Date: 11/16/2010

Current Effective Date: 03/08/2020

with contrast media added throughout the procedure to aid visualization of the decompression. The process is repeated on the opposite side for bilateral decompression of the central canal. The devices are not intended for use near the lateral neural elements and are contraindicated for disc procedures.

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

In 2006, the X-Sten MILD Tool Kit now the mild device kit (X-Sten Corp. renamed Vertos Medical) was cleared for marketing by the U.S. Food and Drug Administration through the 510(k) process for treatment of various spinal conditions. This set of specialized surgical instruments is used to perform percutaneous lumbar decompressive procedures.

Vertos's mild instructions state that the device is not intended for disc procedures but rather for tissue resection at the perilaminar space, within the interlaminar space, and at the ventral aspect of the lamina. The device is not intended for use near the lateral neural elements and remains dorsal to the dura using image guidance and anatomic landmarks.

Food and Drug Administration product code: HRX.

Rationale/Source

Image-guided minimally invasive lumbar decompression (IG-MLD) describes a percutaneous procedure for decompression of the central spinal canal in patients with spinal stenosis and hypertrophy of the ligamentum flavum. In this procedure, a specialized cannula and surgical tools (mild) are used under fluoroscopic guidance for bone and tissue sculpting near the spinal canal. IG-MLD is proposed as an alternative to existing posterior decompression procedures.

For individuals who have lumbar spinal stenosis, or cervical or thoracic spinal stenosis who receive IG-MLD, the evidence includes a large, ongoing randomized controlled trial (n=302), a systematic review of a small randomized controlled trial (n=38), and a number of prospective and retrospective cohort studies and case series. The relevant outcomes are symptoms, functional outcomes, health status measures, and treatment-related morbidity. The largest randomized controlled trial compared IG-MLD with epidural steroid injections (control) in patients who had ligamentum flavum hypertrophy and who failed conservative therapy. Early results have suggested reductions in pain and improvements in function scores in the IG-MLD group vs the control group. The trial was

©2019 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Image-Guided Minimally Invasive Decompression for Spinal Stenosis

Policy # 00278

Original Effective Date: 11/16/2010

Current Effective Date: 03/08/2020

unblinded and there is evidence of differing expectations and follow-up in the two groups, suggesting a high-risk of bias. The available evidence is insufficient to determine the efficacy of mild compared with placebo or to determine the efficacy of IG-MLD compared with open decompression. Trials with relevant control groups could provide greater certainty on the risks and benefits of this procedure. The evidence is insufficient to determine the effects of the technology on health outcomes.

Supplemental Information

Practice Guidelines and Position Statements

No guidelines or statements were identified.

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

Effective for services performed on or after January 9, 2014, the Centers for Medicare & Medicaid Services has determined that percutaneous image-guided lumbar decompression for lumbar spinal stenosis is not reasonable and necessary.

The Centers for Medicare & Medicaid Services determined that percutaneous image-guided lumbar decompression would be covered by Medicare when provided in a clinical study through coverage with evidence development for beneficiaries with lumbar spinal stenosis enrolled in an approved clinical study meeting criteria in the decision memo.

According to the national coverage decision, percutaneous image-guided lumbar decompression is a posterior decompression of the lumbar spine performed under indirect image guidance without any direct visualization of the surgical area. This procedure is proposed as a treatment for symptomatic lumbar spinal stenosis unresponsive to conservative therapy. This procedure is generally described as a noninvasive procedure using specially designed instruments to percutaneously remove a portion of the lamina and debulk the ligamentum flavum. The procedure is performed under x-ray guidance (eg, fluoroscopic, computed tomography) with contrast media to identify and monitor the compressed area via epidurogram.

©2019 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Image-Guided Minimally Invasive Decompression for Spinal Stenosis

Policy # 00278

Original Effective Date: 11/16/2010

Current Effective Date: 03/08/2020

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 2.

Table 2. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
Ongoing			
NCT03072927 ^a	MILD Percutaneous Image-Guided Lumbar Decompression: A Medicare Claims Study	4000	Dec 2020

NCT: national clinical trial.

^a Denotes industry-sponsored or cosponsored trial.

References

1. Blue Cross and Blue Shield Association, Medical Policy Reference Manual, “Image-Guided Minimally Invasive Lumbar Decompression (IG-MLD) for Spinal Stenosis”, 7.01.126, 5:2019
2. Chou R, Baisden J, Carragee EJ, et al. Surgery for low back pain: a review of the evidence for an American Pain Society Clinical Practice Guideline. Spine. May 1 2009;34(10):1094-1109. PMID 19363455
3. Chou R, Loeser JD, Owens DK, et al. Interventional therapies, surgery, and interdisciplinary rehabilitation for low back pain: an evidence-based clinical practice guideline from the American Pain Society. Spine. May 1 2009;34(10):1066-1077. PMID 19363457
4. Weinstein JN, Lurie JD, Tosteson TD, et al. Surgical versus nonsurgical treatment for lumbar degenerative spondylolisthesis. N Engl J Med. May 31 2007;356(22):2257-2270. PMID 17538085
5. Weinstein JN, Tosteson TD, Lurie JD, et al. Surgical versus nonsurgical therapy for lumbar spinal stenosis. N Engl J Med. Feb 21 2008;358(8):794-810. PMID 18287602

©2019 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Image-Guided Minimally Invasive Decompression for Spinal Stenosis

Policy # 00278

Original Effective Date: 11/16/2010

Current Effective Date: 03/08/2020

6. Staats PS, Benyamin RM. MiDAS ENCORE: randomized controlled clinical trial report of 6-month results. *Pain Physician*. Feb 2016;19(2):25-38. PMID 26815247
7. Benyamin RM, Staats PS, Mi DASEI. MILD(R) is an effective treatment for lumbar spinal stenosis with neurogenic claudication: MiDAS ENCORE randomized controlled trial. *Pain Physician*. May 2016;19(4):229-242. PMID 27228511
8. Kreiner DS, Macvicar J, Duszynski B, et al. The mild(R) Procedure: a systematic review of the current literature. *Pain Med*. Feb 2014;15(2):196-205. PMID 24308292
9. Brown LL. A double-blind, randomized, prospective study of epidural steroid injection vs. the mild(R) procedure in patients with symptomatic lumbar spinal stenosis. *Pain Pract*. Jun 2012;12(5):333-341. PMID 22272730
10. Chopko BW. Long-term results of percutaneous lumbar decompression for LSS: two-year outcomes. *Clin J Pain*. Nov 2013;29(11):939-943. PMID 23446067
11. Chopko BW. A novel method for treatment of lumbar spinal stenosis in high-risk surgical candidates: pilot study experience with percutaneous remodeling of ligamentum flavum and lamina. *J Neurosurg Spine*. Jan 2011;14(1):46-50. PMID 21142460
12. Lingreen R, Grider JS. Retrospective review of patient self-reported improvement and post-procedure findings for mild (minimally invasive lumbar decompression). *Pain Physician*. Nov-Dec 2010;13(6):555-560. PMID 21102968
13. Centers for Medicare & Medicaid Services. National Coverage Determination (NCD) for percutaneous image-guided lumbar decompression for lumbar spinal stenosis (150.13). 2014; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=358&ncdver=1&CoverageSelection=Both&ArticleType=All&PolicyType=Final&s=All&Keyword=decompression&KeywordLookUp=Title&KeywordSearchType=And&bc=gAAAABAAAAAAAAA>.

Policy History

Original Effective Date: 11/16/2010

Current Effective Date: 03/08/2020

- 11/04/2010 Medical Policy Committee review
- 11/16/2010 Medical Policy Implementation Committee approval. New policy.
- 11/03/2011 Medical Policy Committee review.
- 11/16/2011 Medical Policy Implementation Committee.
- 12/06/2011 Posted as Archived policy.

©2019 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Image-Guided Minimally Invasive Decompression for Spinal Stenosis

Policy # 00278

Original Effective Date: 11/16/2010

Current Effective Date: 03/08/2020

03/05/2012 Re-posted as active policy.
11/01/2012 Medical Policy Committee review.
11/28/2012 Medical Policy Implementation Committee. No change to coverage.
11/07/2013 Medical Policy Committee review.
11/20/2013 Medical Policy Implementation Committee. No change to coverage.
11/06/2014 Medical Policy Committee review.
11/21/2014 Medical Policy Implementation Committee. No change to coverage.
08/03/2015 Coding update: ICD10 Diagnosis code section added; ICD9 Procedure code section removed.
10/29/2015 Medical Policy Committee review.
11/16/2015 Medical Policy Implementation Committee. No change to coverage.
11/03/2016 Medical Policy Committee review.
11/16/2016 Medical Policy Implementation Committee. No change to coverage.
11/02/2017 Medical Policy Committee review.
11/15/2017 Medical Policy Implementation Committee. Policy statement changed from “lumbar” to “spinal” to include cervical/thoracic decompression. “Lumbar” removed from title.
11/08/2018 Medical Policy Committee review.
11/21/2018 Medical Policy Implementation Committee. No change to coverage.
11/07/2019 Medical Policy Committee review.
11/13/2019 Medical Policy Implementation Committee. No change to coverage.
Next Scheduled Review Date: 11/2020

Coding

The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)‡, copyright 2018 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability

©2019 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Image-Guided Minimally Invasive Decompression for Spinal Stenosis

Policy # 00278

Original Effective Date: 11/16/2010

Current Effective Date: 03/08/2020

attributable or related to any use, nonuse or interpretation of information contained in Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	0274T, 0275T
HCPCS	G0276
ICD-10 Diagnosis	All related diagnoses

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:

©2019 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Image-Guided Minimally Invasive Decompression for Spinal Stenosis

Policy # 00278

Original Effective Date: 11/16/2010

Current Effective Date: 03/08/2020

1. Consultation with the Blue Cross and Blue Shield Association technology assessment program (TEC) or other nonaffiliated technology evaluation center(s);
2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
3. Reference to federal regulations.

‡ Indicated trademarks are the registered trademarks of their respective owners.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

©2019 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.