



Louisiana

Laboratory and Genetic Testing for Use of 5-Fluorouracil in Patients With Cancer

Policy # 00291

Original Effective Date: 03/16/2011

Current Effective Date: 09/14/2020

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Based on review of available data, the Company considers My5-FU^{TM†} testing or other types of assays for determining 5-fluorouracil (5-FU) area under the curve (AUC) in order to adjust 5-FU dose for colorectal cancer (CRC) patients or other cancer patients to be **investigational**.*

Based on review of available data, the Company considers testing for genetic variants in dihydropyrimidine dehydrogenase (*DPYD*) or thymidylate synthase (*TYMS*) to guide 5-FU dosing and/or treatment choice in patients with cancer to be **investigational**.*

Policy Guidelines

Genetics Nomenclature Update

The Human Genome Variation Society nomenclature is used to report information on variants found in DNA and serves as an international standard in DNA diagnostics. It is being implemented for genetic testing medical evidence review updates starting in 2017 (see Table PG1). The Society's nomenclature is recommended by the Human Variome Project, the Human Genome Organization, and by the Human Genome Variation Society itself.

The American College of Medical Genetics and Genomics and the Association for Molecular Pathology standards and guidelines for interpretation of sequence variants represent expert opinion from both organizations, in addition to the College of American Pathologists. These recommendations primarily apply to genetic tests used in clinical laboratories, including genotyping, single genes, panels, exomes, and genomes. Table PG2 shows the recommended standard

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Laboratory and Genetic Testing for Use of 5-Fluorouracil in Patients With Cancer

Policy # 00291

Original Effective Date: 03/16/2011

Current Effective Date: 09/14/2020

terminology-“pathogenic,” “likely pathogenic,” “uncertain significance,” “likely benign,” and “benign”-to describe variants identified that cause Mendelian disorders.

Table PG1. Nomenclature to Report on Variants Found in DNA

Previous	Updated	Definition
Mutation	Disease-associated variant	Disease-associated change in the DNA sequence
	Variant	Change in the DNA sequence
	Familial variant	Disease-associated variant identified in a proband for use in subsequent targeted genetic testing in first-degree relatives

Table PG2. ACMG - AMP Standards and Guidelines for Variant Classification

Variant Classification	Definition
Pathogenic	Disease-causing change in the DNA sequence
Likely pathogenic	Likely disease-causing change in the DNA sequence
Variant of uncertain significance	Change in DNA sequence with uncertain effects on disease
Likely benign	Likely benign change in the DNA sequence
Benign	Benign change in the DNA sequence

ACMG: American College of Medical Genetics and Genomics; AMP: Association for Molecular Pathology.

Genetic Counseling

Experts recommend formal genetic counseling for patients who are at risk for inherited disorders and who wish to undergo genetic testing. Interpreting the results of genetic tests and understanding risk factors can be difficult for some patients; genetic counseling helps individuals understand the impact of genetic testing, including the possible effects the test results could have on the individual or their family members. It should be noted that genetic counseling may alter the utilization of genetic testing substantially and may reduce inappropriate testing; further, genetic counseling should

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Laboratory and Genetic Testing for Use of 5-Fluorouracil in Patients With Cancer

Policy # 00291

Original Effective Date: 03/16/2011

Current Effective Date: 09/14/2020

be performed by an individual with experience and expertise in genetic medicine and genetic testing methods.

Background/Overview

5-fluorouracil

The agent 5-fluorouracil is a widely used antineoplastic chemotherapy drug that targets thymidylate synthase (*TYMS*) enzyme, which is involved in DNA production. 5-fluorouracil has been used for many years to treat solid tumors (eg, colon and rectal cancer, head and neck cancer). In general, the incidence of grade 3 or 4 toxicity (mainly neutropenia, diarrhea, mucositis, and hand-foot syndrome) increases with higher systemic exposure to 5-fluorouracil. Several studies also have reported statistically significant positive associations between 5-fluorouracil exposure and tumor response. In current practice, however, 5-fluorouracil dose is reduced when symptoms of severe toxicity appear but is seldom increased to promote efficacy.

Based on known 5-fluorouracil pharmacology, it is possible to determine a sampling scheme for the area under the curve determination and to optimize an area under the curve target and dose-adjustment algorithm for a particular 5-fluorouracil chemotherapy regimen and patient population. For each area under the curve value or range, the algorithm defines the dose adjustment during the next chemotherapy cycle most likely to achieve the target area under the curve without overshooting and causing severe toxicity.

In clinical research studies, 5-fluorouracil blood plasma levels most recently have been determined by high-performance liquid chromatography or liquid chromatography coupled with tandem mass spectrometry. Both methods require expertise to develop an in-house assay and may be less amenable to routine clinical laboratory settings.

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratory-developed tests must meet the general regulatory standards of the Clinical Laboratory Improvement Amendments (CLIA). My5-fluorouracil^{TM†} (Saladax Biomedical) and genetic testing for variants in *DPYD* and *TYMS* for predicting the risk of 5-fluorouracil toxicity and chemotherapeutic response (ARUP Laboratories) are available under the auspices of the CLIA.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Laboratory and Genetic Testing for Use of 5-Fluorouracil in Patients With Cancer

Policy # 00291

Original Effective Date: 03/16/2011

Current Effective Date: 09/14/2020

Laboratories that offer laboratory-developed tests must be licensed by the CLIA for high-complexity testing. To date, the U.S. Food and Drug Administration has chosen not to require any regulatory review of this test.

Rationale/Source

Variability in systemic exposure to 5-fluorouracil chemotherapy is thought to directly impact 5-fluorouracil tolerability and efficacy. The standard approach is dosing according to body surface area. Two alternative approaches have been proposed for modifying use of 5-fluorouracil: (1) dosing based on the determined area under the curve serum concentration target and (2) genetic testing for variants affecting 5-fluorouracil metabolism. For genetic testing, currently available polymerase chain reaction tests assess specific variants in genes encoding dihydropyrimidine reductase (*DPYD*) and thymidylate synthase (*TYMS*) in the catabolic and anabolic pathways of 5-fluorouracil metabolism, respectively.

For individuals who have cancer for whom treatment with 5-fluorouracil is indicated who receive laboratory assays to determine 5-fluorouracil area under the curve, the evidence includes randomized controlled trials (RCTs), observational studies, and systematic reviews. Relevant outcomes are overall survival, disease-specific survival, test accuracy and validity, and treatment-related morbidity. Several analyses of patients with colorectal cancer have evaluated clinical validity. One study, for example, found that the rate of severe toxicity was significantly lower in patients with stage II and III cancer who chose pharmacokinetic monitoring versus body surface area monitoring but progression-free survival did not differ between groups in patients with stage IV or recurrent cancer. No RCTs or nonrandomized comparative studies were identified comparing health outcomes in cancer patients who did and did not have 5-fluorouracil dose adjustment using the My 5-fluorouracil assay and who were treated with chemotherapy regimens used in current clinical practice. A systematic review of the available literature found a significantly higher response rate with body surface area based monitoring and no significant difference in toxicity. Most data derived from observational studies and the RCTs were conducted in the 1980s when different chemotherapy protocols were used. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have cancer for whom treatment with 5-fluorouracil is indicated who receive genetic testing for variants (eg, in *DPYD* and *TYMS*) affecting 5-fluorouracil metabolism, the

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Laboratory and Genetic Testing for Use of 5-Fluorouracil in Patients With Cancer

Policy # 00291

Original Effective Date: 03/16/2011

Current Effective Date: 09/14/2020

evidence includes observational studies and systematic reviews. Relevant outcomes are overall survival, disease-specific survival, test accuracy and validity, and treatment-related morbidity. A TEC Assessment (2010) concluded that *DPYD* and *TYMS* variant testing had poor prognostic capacity to identify patients likely to experience severe 5-fluorouracil toxicity. Since the publication of that Assessment, no prospective trials comparing the efficacy and toxicity outcomes in patients who did and did not undergo pretreatment *DPYD* and/or *TYMS* testing have been published. Three prospective observational studies used a historical control group and 1 also used a matched-pairs analysis to compare outcomes in patients who received genotype-based dosing to those who received standard dosing. No differences in overall survival, progression-free survival, or tumor progression were observed. Risk of serious toxicity was higher in *DPYD* allele carriers who received genotype-based dosing compared to wild-type patients but lower when compared to historical controls who were carriers but received standard dosing. The evidence is limited by retrospective data collection, use of historical control groups, small sample sizes, and missing data. The evidence is insufficient to determine the effects of the technology on health outcomes.

Supplemental Information

Practice Guidelines and Position Statements

National Comprehensive Cancer Network Guidelines

National Comprehensive Cancer Network guidelines do not recommend use of area under the curve guidance for 5-fluorouracil dosing or genetic testing for *DPYD* and/or *TYMS* variants in patients with colon, rectal, breast, gastric, pancreatic cancer, or head and neck cancers.

International Association of Therapeutic Drug Monitoring and Clinical Toxicology

In 2019, the International Association of Therapeutic Drug Monitoring and Clinical Toxicology published recommendations for therapeutic drug monitoring of 5-Fluorouracil therapy. The work was supported in part by grants from the National Cancer Institute National Institutes of Health. Several authors reported relationships with Saladax, the manufacturer of the My5-fluorouracil test. The committee concluded that there was sufficient evidence to strongly recommend therapeutic drug monitoring for the management of 5-fluorouracil therapy in patients with early or advanced colorectal cancer and patients with squamous cell carcinoma of head-and-neck cancer receiving common 5-fluorouracil dosing regimens.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Laboratory and Genetic Testing for Use of 5-Fluorouracil in Patients With Cancer

Policy # 00291

Original Effective Date: 03/16/2011

Current Effective Date: 09/14/2020

Clinical Pharmacogenetics Implementation Consortium

In 2009, the Clinical Pharmacogenetics Implementation Consortium was formed as a shared project between PharmGKB, an internet research tool developed by Stanford University, and the Pharmacogenomics Research Network of the National Institutes of Health. In 2013, the Clinical Pharmacogenetics Implementation Consortium published evidence-based guidelines for *DPYD* genotype and fluoropyrimidine dosing. The guidelines did not address testing.

An update to the Clinical Pharmacogenetics Implementation Consortium (2017) guidelines was published by Amstutz et al (2018). As in 2013, the primary focus of the guidelines was on the *DPYD* genotype and implications for dosing of fluoropyrimidine. In the 2017 update, the Clinical Pharmacogenetics Implementation Consortium noted that genetic testing for *DPYD* may include “resequencing of the complete coding regions” or may be confined to analysis of particular risk variants, among which Clinical Pharmacogenetics Implementation Consortium listed the c.190511G>A, c.1679T>G, c.2846A>T, and c.1129-5923C>G variants, as affecting 5-fluorouracil toxicity. The guideline further noted that, while other genes (*TYMS*, *MTHFR*) may be tested for variants, the clinical utility of such tests is yet unproven. In patients who have undergone genetic testing and who are known carriers of a *DPYD* risk variant, the guidelines recommended that caregivers strongly reduce the dosage of 5-fluorouracil-based treatments, or exclude them, depending on the patient’s level of *DPYD* activity. CPIC advised follow-up therapeutic drug monitoring to guard against underdosing and cautioned that genetic tests could be limited to known risk variants and, therefore, not identify other *DPYD* variants.

National Institute for Health and Care Excellence

In 2014, the National Institute of Health and Care Excellence published evidence-based diagnostics guidance on the 5-fluorouracil assay for 5-fluorouracil chemotherapy dose adjustment. The guidance stated: “The My5-fluorouracil assay is only recommended for use in research for guiding dose adjustment in people having fluorouracil chemotherapy by continuous infusion. The My5-fluorouracil assay shows promise and the development of robust evidence is recommended to demonstrate its utility in clinical practice.”

U.S. Preventive Services Task Force Recommendations

Not applicable.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Laboratory and Genetic Testing for Use of 5-Fluorouracil in Patients With Cancer

Policy # 00291

Original Effective Date: 03/16/2011

Current Effective Date: 09/14/2020

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

Ongoing and Unpublished Clinical Trials

There are currently no relevant ongoing trials. Some unpublished trials that might influence this review are listed in Table 1.

Table 1. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Unpublished</i>			
<i>Area under the curve-guided dosing of 5-fluorouracil</i>			
NCT00943137	The Optimisation of 5-Fluorouracil Dose by Pharmacokinetic Monitoring in Asian Patients With Advanced Stage Cancer	55	June 2017
NCT02055560a	Retrospective Data Comparison of Toxicity and Efficacy in Colorectal Cancer (CRC) Patients Managed With and Without 5-fluorouracil Exposure Optimization Testing	350	Dec 2017(unknown)

NCT: national clinical trial.

^a Denotes industry-sponsored or cosponsored trial.

References

1. Blue Cross and Blue Shield Association, Medical Policy Reference Manual, “Laboratory and Genetic Testing for Use of 5-Fluorouracil in Patients with Cancer”, 2.04.68, April 2020.
2. Saladax Biomedical Inc. My5-FU: 5-Fluorouracil (My5-FUTM) Assay. 2018. <https://mycaretests.com/oncology/products>
3. Kline CL, Schiccitano A, Zhu J, et al. Personalized dosing via pharmacokinetic monitoring of 5-fluorouracil might reduce toxicity in early- or late-stage colorectal cancer patients treated with

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Laboratory and Genetic Testing for Use of 5-Fluorouracil in Patients With Cancer

Policy # 00291

Original Effective Date: 03/16/2011

Current Effective Date: 09/14/2020

- infusional 5-fluorouracil-based chemotherapy regimens. Clin Colorectal Cancer. Jun 2014;13(2):119-126. PMID 24461492
4. Saam J, Critchfield GC, Hamilton SA, et al. Body surface area-based dosing of 5-fluorouracil results in extensive interindividual variability in 5-fluorouracil exposure in colorectal cancer patients on FOLFOX regimens. Clin Colorectal Cancer. Sep 2011;10(3):203-206. PMID 21855044
 5. Gamelin E, Boisdron-Celle M, Delva R, et al. Long-term weekly treatment of colorectal metastatic cancer with fluorouracil and leucovorin: results of a multicentric prospective trial of fluorouracil dosage optimization by pharmacokinetic monitoring in 152 patients. J Clin Oncol. Apr 1998;16(4):1470-1478. PMID 9552054
 6. Boisdron-Celle M, Craipeau C, Brienza S, et al. Influence of oxaliplatin on 5-fluorouracil plasma clearance and clinical consequences. Cancer Chemother Pharmacol. Mar 2002;49(3):235-243. PMID 11935216
 7. Wilhelm M, Mueller L, Miller MC, et al. Prospective, Multicenter Study of 5-Fluorouracil Therapeutic Drug Monitoring in Metastatic Colorectal Cancer Treated in Routine Clinical Practice. Clin Colorectal Cancer. 2016 Dec;15(4). PMID 27256667
 8. Milano G, Etienne MC, Renee N, et al. Relationship between fluorouracil systemic exposure and tumor response and patient survival. J Clin Oncol. Jun 1994;12(6):1291-1295. PMID 8201391
 9. Santini J, Milano G, Thyss A, et al. 5-FU therapeutic monitoring with dose adjustment leads to an improved therapeutic index in head and neck cancer. Br J Cancer. Feb 1989;59(2):287-290. PMID 2930694
 10. Gamelin EC, Danquechin-Dorval EM, Dumesnil YF, et al. Relationship between 5-fluorouracil (5-FU) dose intensity and therapeutic response in patients with advanced colorectal cancer receiving infusional therapy containing 5-FU. Cancer. Feb 1 1996;77(3):441-451. PMID 8630950
 11. Gamelin E, Delva R, Jacob J, et al. Individual fluorouracil dose adjustment based on pharmacokinetic follow-up compared with conventional dosage: results of a multicenter randomized trial of patients with metastatic colorectal cancer. J Clin Oncol. May 1 2008;26(13):2099-2105. PMID 18445839
 12. Andre T, Boni C, Navarro M, et al. Improved overall survival with oxaliplatin, fluorouracil, and leucovorin as adjuvant treatment in stage II or III colon cancer in the MOSAIC trial. J Clin Oncol. Jul 1 2009;27(19):3109-3116. PMID 19451431

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Laboratory and Genetic Testing for Use of 5-Fluorouracil in Patients With Cancer

Policy # 00291

Original Effective Date: 03/16/2011

Current Effective Date: 09/14/2020

13. de Gramont A, Figer A, Seymour M, et al. Leucovorin and fluorouracil with or without oxaliplatin as first-line treatment in advanced colorectal cancer. *J Clin Oncol*. Aug 2000;18(16):2938-2947. PMID 10944126
14. Fety R, Rolland F, Barberi-Heyob M, et al. Clinical impact of pharmacokinetically-guided dose adaptation of 5- fluorouracil: results from a multicentric randomized trial in patients with locally advanced head and neck carcinomas. *Clin Cancer Res*. Sep 1998;4(9):2039-2045. PMID 9748117
15. Yang R, Zhang Y, Zhou H, et al. Individual 5-fluorouracil dose adjustment via pharmacokinetic monitoring versus conventional body-area-surface method: a meta-analysis. *Ther Drug Monit*. Feb 2016;38(1):79-86. PMID 26309030
16. Grem JL. 5-Fluorouracil and its biomodulation in the management of colorectal cancer. In: Saltz LB, ed. *Colorectal Cancer: Multimodality Management*. Totowa, NJ: Humana Press; 2002.
17. Caudle KE, Thorn CF, Klein TE, et al. Clinical Pharmacogenetics Implementation Consortium guidelines for dihydropyrimidine dehydrogenase genotype and fluoropyrimidine dosing. *Clin Pharmacol Ther*. Dec 2013;94(6):640-645. PMID 23988873
18. ARUP Laboratories. 5-Fluorouracil Toxicity and Chemotherapeutic Response Panel. 2016; <http://ltd.aruplab.com/Tests/Pdf/128..>
19. Li Q, Liu Y, Zhang HM, et al. Influence of DPYD genetic polymorphisms on 5-fluorouracil toxicities in patients with colorectal cancer: a meta-analysis. *Gastroenterol Res Pract*. Jan 2014;2014:827989. PMID 25614737
20. Rosmarin D, Palles C, Church D, et al. Genetic markers of toxicity from capecitabine and other fluorouracil- based regimens: investigation in the QUASAR2 study, systematic review, and meta-analysis. *J Clin Oncol*. Apr 1 2014;32(10):1031-1039. PMID 24590654
21. Schwab M, Zanger UM, Marx C, et al. Role of genetic and nongenetic factors for fluorouracil treatment-related severe toxicity: a prospective clinical trial by the German 5-FU Toxicity Study Group. *J Clin Oncol*. May 1 2008;26(13):2131-2138. PMID 18299612
22. Boige V, Vincent M, Alexandre P, et al. DPYD genotyping to predict adverse events following treatment with fluorouracil-based adjuvant chemotherapy in patients with stage III colon cancer: a secondary analysis of the PETACC-8 Randomized Clinical Trial. *JAMA Oncol*. Jan 21 2016;2(5):655-662. PMID 26794347
23. Vazquez C, Orlova M, Angriman F, et al. Prediction of severe toxicity in adult patients under treatment with 5- fluorouracil: a prospective cohort study. *Anticancer Drugs*. Oct 2017;28(9):1039-1046. PMID 28723867

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Laboratory and Genetic Testing for Use of 5-Fluorouracil in Patients With Cancer

Policy # 00291

Original Effective Date: 03/16/2011

Current Effective Date: 09/14/2020

24. Wang YC, Xue HP, Wang ZH, et al. An integrated analysis of the association between Ts gene polymorphisms and clinical outcome in gastric and colorectal cancer patients treated with 5-FU-based regimens. *Mol Biol Rep.* Jul 2013;40(7):4637-4644. PMID 23645036
25. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Pharmacogenetic Testing to Predict Serious Toxicity From 5-Fluorouracil (5-FU) for Patients Administered 5-FU-Based Chemotherapy for Cancer. *TEC Assessments.* 2010;24:Tab 13.
26. Deenen MJ, Meulendijks D, Cats A, et al. Upfront genotyping of DPYD*2A to individualize fluoropyrimidine therapy: a safety and cost analysis. *J Clin Oncol.* Jan 20 2016;34(3):227-234. PMID 26573078
27. Henricks, LL, van Merendonk, LL, Meulendijks, DD, Deenen, MM, Beijnen, JJ, de Boer, AA, Cats, AA, Schellens, JJ. Effectiveness and safety of reduced-dose fluoropyrimidine therapy in patients carrying the DPYD*2A variant: A matched pair analysis. *Int. J. Cancer,* 2018 Nov 30;144(9). PMID 30485432
28. Henricks, LL, Lunenburg, CC, de Man, FF, Meulendijks, DD, Frederix, GG, Kienhuis, EE, Creemers, GG, Baars, AA, Dezentj, VV, Imholz, AA, Jeurissen, FF, Portielje, JJ, Jansen, RR, Hamberg, PP, Ten Tije, AA, Droogendijk, HH, Koopman, MM, Nieboer, PP, van de Poel, MM, Mandigers, CC, Rosing, HH, Beijnen, JJ, Werkhoven, EE, van Kuilenburg, AA, van Schaik, RR, Mathijssen, RR, Swen, JJ, Gelderblom, HH, Cats, AA, Guchelaar, HH, Schellens, JJ. DPYD genotype-guided dose individualisation of fluoropyrimidine therapy in patients with cancer: a prospective safety analysis. *Lancet Oncol.,* 2018 Oct 24;19(11). PMID 30348537
29. Goff LW, Thakkar N, Du L, et al. Thymidylate synthase genotype-directed chemotherapy for patients with gastric and gastroesophageal junction cancers. *PLoS One.* Sep 2014;9(9):e107424. PMID 25232828
30. Magnani E, Farnetti E, Nicoli D, et al. Fluoropyrimidine toxicity in patients with dihydropyrimidine dehydrogenase splice site variant: the need for further revision of dose and schedule. *Intern Emerg Med.* Aug 2013;8(5):417-423. PMID 23585145
31. Cremolini, CC, Del Re, MM, Antoniotti, CC, Lonardi, SS, Bergamo, FF, Loupakis, FF, Borelli, BB, Marmorino, FF, Citi, VV, Cortesi, EE, Moretto, RR, Ronzoni, MM, Tomasello, GG, Zaniboni, AA, Racca, PP, Buonadonna, AA, Allegrini, GG, Ricci, VV, Di Donato, SS, Zagonel, VV, Boni, LL, Falcone, AA, Danesi, RR. *NA. Oncotarget,* 2018 Mar 1;9(8). PMID 29487697
32. National Comprehensive Cancer Network (NCCN). *NCCN Clinical Practice Guidelines in Oncology: Colon Cancer. Version 1.2020.*
http://www.nccn.org/professionals/physician_gls/pdf/colon.pdf.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Laboratory and Genetic Testing for Use of 5-Fluorouracil in Patients With Cancer

Policy # 00291

Original Effective Date: 03/16/2011

Current Effective Date: 09/14/2020

33. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Rectal Cancer. Version 1.2020.
http://www.nccn.org/professionals/physician_gls/pdf/rectal.pdf.
34. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Breast Cancer. Version 1.2020.
http://www.nccn.org/professionals/physician_gls/pdf/breast.pdf.
35. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Gastric Cancer. Version 4.2019.
http://www.nccn.org/professionals/physician_gls/pdf/gastric.pdf.
36. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Pancreatic Cancer. Version 1.2020.
http://www.nccn.org/professionals/physician_gls/pdf/pancreatic.pdf.
37. Beumer JH, Chu E, Allegra C, et al. Therapeutic Drug Monitoring in Oncology: International Association of Therapeutic Drug Monitoring and Clinical Toxicology Recommendations for 5-Fluorouracil Therapy. Clin. Pharmacol. Ther. 2019 Mar;105(3). PMID 29923599
38. Amstutz U, Henricks LM, Offer SM, et al. Clinical Pharmacogenetics Implementation Consortium (CPIC) guideline for dihydropyrimidine dehydrogenase genotype and fluoropyrimidine dosing: 2017 update. Clin Pharmacol Ther. Feb 2018;103(2):210-216. PMID 29152729
39. National Institute for Health and Care Excellence (NICE). Fluorouracil chemotherapy: The My5-FU assay for guiding dose adjustment [DG16]. 2014; <https://www.nice.org.uk/guidance/dg16>.

Policy History

Original Effective Date: 03/16/2011

Current Effective Date: 09/14/2020

- | | |
|------------|---|
| 03/03/2011 | Medical Policy Committee review |
| 03/16/2011 | Medical Policy Implementation Committee approval. New Policy. |
| 03/01/2012 | Medical Policy Committee review |
| 03/21/2012 | Medical Policy Implementation Committee approval. Coverage eligibility unchanged. |
| 05/02/2013 | Medical Policy Committee review |
| 05/22/2013 | Medical Policy Implementation Committee approval. Coverage eligibility unchanged. |

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Laboratory and Genetic Testing for Use of 5-Fluorouracil in Patients With Cancer

Policy # 00291

Original Effective Date: 03/16/2011

Current Effective Date: 09/14/2020

06/05/2014 Medical Policy Committee review

06/18/2014 Medical Policy Implementation Committee approval. Investigational OnDose statement modified to reflect new test name, My5-FU. Investigational statement for TheraGuide testing for genetic mutations in *DPYD* or *TYMS* added. Title changed from “Laboratory Testing to Allow Area Under the Curve (AUC) Targeted 5-Fluorouracil (5-FU) Dosing for Patients Administered 5-FU for Cancer” to “Laboratory and Genetic Testing for Use of 5-Fluorouracil in Patients With Cancer” to reflect incorporation of new test.

08/06/2015 Medical Policy Committee review

08/19/2015 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

08/04/2016 Medical Policy Committee review

08/17/2016 Medical Policy Implementation Committee approval. Removed TheraGuide test from policy statement as it is no longer available.

01/01/2017 Coding update: Removing ICD-9 Diagnosis Codes

08/03/2017 Medical Policy Committee review

08/23/2017 Medical Policy Implementation Committee approval. Genetic testing nomenclature updated. “Mutations” changed to “variants” throughout the policy. Coverage eligibility unchanged.

08/09/2018 Medical Policy Committee review

08/15/2018 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

03/01/2019 Coding update

08/01/2019 Medical Policy Committee review

08/14/2019 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

08/06/2020 Medical Policy Committee review

08/12/2020 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

Next Scheduled Review Date: 08/2021

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Laboratory and Genetic Testing for Use of 5-Fluorouracil in Patients With Cancer

Policy # 00291

Original Effective Date: 03/16/2011

Current Effective Date: 09/14/2020

Coding

The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)‡, copyright 2019 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	81232, 81346, 81400, 81401, 84999
HCPCS	S3722
ICD-10 Diagnosis	All related diagnoses

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Laboratory and Genetic Testing for Use of 5-Fluorouracil in Patients With Cancer

Policy # 00291

Original Effective Date: 03/16/2011

Current Effective Date: 09/14/2020

standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
 1. Consultation with the Blue Cross and Blue Shield Association technology assessment program (TEC) or other nonaffiliated technology evaluation center(s);
 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
 3. Reference to federal regulations.

‡ Indicated trademarks are the registered trademarks of their respective owners.

NOTICE: If the Patient's health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.