



# Louisiana

## **Navigated Transcranial Magnetic Stimulation**

**Policy #** 00407

**Original Effective Date:** 03/19/2014

**Current Effective Date:** 10/12/2020

*Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.*

*Note: Magnetoencephalography/Magnetic Source Imaging is addressed separately in medical policy 00082.*

### **Services Are Considered Investigational**

*Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.*

Based on review of available data, the Company considers navigated transcranial magnetic stimulation (nTMS) for all purposes, including but not limited to the preoperative evaluation of patients being considered for brain surgery, when localization of eloquent areas of the brain (e.g., controlling verbal or motor function) is an important consideration in surgical planning to be **investigational**.\*

### **Background/Overview**

#### **Management of Brain Tumors**

Surgical management of brain tumors involves resecting the brain tumor and preserving essential brain function. "Mapping" of brain functions, such as body movement and language, is most accurately achieved with direct cortical stimulation (DCS), an intraoperative procedure that lengthens operating times and requires a wide surgical opening. Even if not completely accurate compared with DCS, preoperative techniques that map brain functions may aid in planning the extent of resection and the surgical approach. Although DCS is still usually performed to confirm the brain locations associated with specific functions, preoperative mapping techniques may provide useful information that improves patient outcomes.

#### **Noninvasive Mapping Techniques**

The most commonly used tool for the noninvasive localization of brain functions is functional magnetic resonance imaging (fMRI). Functional MRI identifies regions of the brain where there are changes in localized cortical blood oxygenation, which correlate with the neuronal activity

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Navigated Transcranial Magnetic Stimulation

Policy # 00407

Original Effective Date: 03/19/2014

Current Effective Date: 10/12/2020

associated with a specific motor or speech task being performed as the image is obtained. The accuracy and precision of fMRI depend on the patient's ability to perform the isolated motor task, such as moving the single assigned muscle without moving others. This may be difficult in patients in whom brain tumors have caused partial or complete paresis. The reliability of fMRI in mapping language areas has been questioned. Guissani et al (2010) reviewed several studies comparing fMRI with DCS of language areas and found large variability in the sensitivity and specificity rates of fMRI. Reviewers also pointed out a major conceptual point in how fMRI and DCS "map" language areas: fMRI identifies regional oxygenation changes, which show that a particular region of the brain is involved in the capacity of interest, whereas DCS locates specific areas in which the activity of interest is disrupted. Regions of the brain involved in a certain activity may not necessarily be required for that activity and could theoretically be safely resected.

Magnetoencephalography (MEG) is also used to map brain activity. In this procedure, electromagnetic recorders are attached to the scalp. Unlike electroencephalography, MEG records magnetic fields generated by electric currents in the brain, rather than the electric currents themselves. Magnetic fields tend to be less distorted by the skull and scalp than electric currents, yielding an improved spatial resolution. MEG is conducted in a magnetically shielded room to screen out environmental electric or magnetic noises that could interfere with the MEG recording. (See medical policy 00082 for additional information on MEG and magnetic resonance imaging.)

Navigated transcranial magnetic stimulation (nTMS) is a noninvasive imaging method for evaluating eloquent brain areas. Transcranial magnetic pulses are delivered to the patient as a navigation system calculates the strength, location, and direction of the stimulating magnetic field. The locations of these pulses are registered to a magnetic resonance image of the patient's brain. Surface electromyography electrodes are attached to various limb muscles of the patient. Moving the magnetic stimulation source to various parts of the brain causes electromyography electrodes to respond, indicating the part of the cortex involved in particular muscle movements. For evaluation of language areas, magnetic stimulation areas that disrupt specific speech tasks are thought to identify parts of the brain involved in speech function. Navigated TMS can be considered a noninvasive alternative to DCS, in which electrodes are directly applied to the surface of the cortex during craniotomy. Navigated TMS is being evaluated as an alternative to other noninvasive cortical mapping techniques (eg, fMRI, MEG) for presurgical identification of cortical areas involved in motor and language functions. Navigated TMS, used for cortical language area mapping, is also

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

Navigated Transcranial Magnetic Stimulation

Policy # 00407

Original Effective Date: 03/19/2014

Current Effective Date: 10/12/2020

being investigated in combination with diffusion tensor imaging tractography for subcortical white matter tract mapping.

## **FDA or Other Governmental Regulatory Approval**

### **U.S. Food and Drug Administration (FDA)**

In 2009, the eXimia Navigated Brain Stimulation System (Nexstim) was cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process for noninvasive mapping of the primary motor cortex of the brain to its cortical gyrus for preprocedural planning.

Similarly, in May 2012, the Nexstim Navigated Brain Stimulation System 4 and Navigated Brain Stimulation System 4 with NexSpeech<sup>®†</sup> were cleared for marketing by the FDA through the 510(k) process for noninvasive mapping of the primary motor cortex and for localization of cortical areas that do not contain speech function for preprocedural planning.

## **Rationale/Source**

For individuals who have brain lesion(s) undergoing preoperative evaluation for localization of eloquent areas of the brain who receive nTMS, the evidence includes controlled observational studies and case series. Relevant outcomes are overall survival, test accuracy, morbid events, and functional outcomes. Several small studies have evaluated the distance between nTMS hotspots and direct cortical stimulation hotspots for the same muscle. Although the average distance in most studies is 10 mm or less, this does not take into account the error margin in this average distance or whether hotspots are missed. It is difficult to verify nTMS hotspots fully because only exposed cortical areas can be verified with direct cortical stimulation. Limited studies of nTMS evaluating language areas have shown high false-positive rates (low specificity) and sensitivity that may be insufficient for clinical use. Several controlled observational studies have compared outcomes in patients undergoing nTMS with those (generally pre-TMS historical controls) who did not undergo nTMS. Findings of the studies were mixed; outcomes were not consistently better in patients who underwent presurgical nTMS. For example, overall survival did not differ significantly between groups in 2 studies and 1 reporting postoperative language deficits found significantly fewer deficits in the group that had presurgical nTMS. The controlled observational studies had various methodologic limitations and, being nonrandomized, might not have adequately controlled for differences in patient groups, which could have biased outcomes. The evidence is insufficient to determine the effects of the technology on health outcomes.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

Navigated Transcranial Magnetic Stimulation

Policy # 00407

Original Effective Date: 03/19/2014

Current Effective Date: 10/12/2020

## **Supplemental Information**

### **Clinical Input From Physician Specialty Societies and Academic Medical Centers**

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

In response to requests, input was received from 1 physician specialty society (2 reviewers) and 2 academic medical centers while this policy was under review in 2013. Most reviewers considered navigated transcranial magnetic stimulation to be investigational.

### **Practice Guidelines and Position Statements**

No guidelines or statements were identified.

### **U.S. Preventive Services Task Force Recommendations**

Not applicable.

### **Medicare National Coverage**

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

### **Ongoing and Unpublished Clinical Trials**

Some currently unpublished trials that might influence this review are listed in Table 1.

**Table 1. Summary of Key Trials**

| <b>NCT No.</b>           | <b>Trial Name</b>  | <b>Planned Enrollment</b> | <b>Completion Date</b> |
|--------------------------|--|---------------------------|------------------------|
| <i>Ongoing</i>           |  |                           |                        |
| NCT02741193 <sup>a</sup> | Validation of Presurgical Motor Mapping With Transcranial Magnetic Stimulation (TMS) in Patients With Epilepsy | 14                        | Jun 2021               |

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Navigated Transcranial Magnetic Stimulation

Policy # 00407

Original Effective Date: 03/19/2014

Current Effective Date: 10/12/2020

NCT: national clinical trial.

<sup>a</sup> Denotes industry-sponsored or cosponsored trial.

## **References**

1. Blue Cross and Blue Shield Association, Medical Policy Reference Manual, “Navigated Transcranial Magnetic Stimulation”, Policy 2.01.90, August 2020.
2. Mangraviti A, Casali C, Cordella R, et al. Practical assessment of preoperative functional mapping techniques: navigated transcranial magnetic stimulation and functional magnetic resonance imaging. *Neurol Sci*. Sep 2013; 34(9): 1551-7. PMID 23266868
3. Opitz A, Zafar N, Bockermann V, et al. Validating computationally predicted TMS stimulation areas using direct electrical stimulation in patients with brain tumors near precentral regions. *Neuroimage Clin*. 2014; 4: 500-7. PMID 24818076
4. Rizzo V, Terranova C, Conti A, et al. Preoperative functional mapping for rolandic brain tumor surgery. *Neurosci Lett*. Nov 07 2014; 583: 136-41. PMID 25224631
5. Forster MT, Limbart M, Seifert V, et al. Test-retest reliability of navigated transcranial magnetic stimulation of the motor cortex. *Neurosurgery*. Mar 2014; 10 Suppl 1: 51-5; discussion 55-6. PMID 23842557
6. Kato N, Schilt S, Schneider H, et al. Functional brain mapping of patients with arteriovenous malformations using navigated transcranial magnetic stimulation: first experience in ten patients. *Acta Neurochir (Wien)*. May 2014; 156(5): 885-95. PMID 24639144
7. Weiss C, Nettekoven C, Rehme AK, et al. Mapping the hand, foot and face representations in the primary motor cortex - retest reliability of neuronavigated TMS versus functional MRI. *Neuroimage*. Feb 01 2013; 66: 531-42. PMID 23116812
8. Schmidt S, Bathe-Peters R, Fleischmann R, et al. Nonphysiological factors in navigated TMS studies; confounding covariates and valid intracortical estimates. *Hum Brain Mapp*. Jan 2015; 36(1): 40-9. PMID 25168635
9. Sollmann N, Ille S, Boeckh-Behrens T, et al. Mapping of cortical language function by functional magnetic resonance imaging and repetitive navigated transcranial magnetic stimulation in 40 healthy subjects. *Acta Neurochir (Wien)*. Jul 2016; 158(7): 1303-16. PMID 27138329
10. Sollmann N, Tanigawa N, Tussis L, et al. Cortical regions involved in semantic processing investigated by repetitive navigated transcranial magnetic stimulation and object naming. *Neuropsychologia*. Apr 2015; 70: 185-95. PMID 25731903

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Navigated Transcranial Magnetic Stimulation

Policy # 00407

Original Effective Date: 03/19/2014

Current Effective Date: 10/12/2020

11. Picht T, Schmidt S, Brandt S, et al. Preoperative functional mapping for rolandic brain tumor surgery: comparison of navigated transcranial magnetic stimulation to direct cortical stimulation. *Neurosurgery*. Sep 2011; 69(3): 581-8; discussion 588. PMID 21430587
12. Forster MT, Hattingen E, Senft C, et al. Navigated transcranial magnetic stimulation and functional magnetic resonance imaging: advanced adjuncts in preoperative planning for central region tumors. *Neurosurgery*. May 2011; 68(5): 1317-24; discussion 1324-5. PMID 21273929
13. Tarapore PE, Tate MC, Findlay AM, et al. Preoperative multimodal motor mapping: a comparison of magnetoencephalography imaging, navigated transcranial magnetic stimulation, and direct cortical stimulation. *J Neurosurg*. Aug 2012; 117(2): 354-62. PMID 22702484
14. Krieg SM, Shibani E, Buchmann N, et al. Utility of presurgical navigated transcranial magnetic brain stimulation for the resection of tumors in eloquent motor areas. *J Neurosurg*. May 2012; 116(5): 994-1001. PMID 22304452
15. Picht T, Krieg SM, Sollmann N, et al. A comparison of language mapping by preoperative navigated transcranial magnetic stimulation and direct cortical stimulation during awake surgery. *Neurosurgery*. May 2013; 72(5): 808-19. PMID 23385773
16. Tarapore PE, Findlay AM, Honma SM, et al. Language mapping with navigated repetitive TMS: proof of technique and validation. *Neuroimage*. Nov 15 2013; 82: 260-72. PMID 23702420
17. Hendrix P, Senger S, Simgen A, et al. Preoperative rTMS Language Mapping in Speech-Eloquent Brain Lesions Resected Under General Anesthesia: A Pair-Matched Cohort Study. *World Neurosurg*. Apr 2017; 100: 425-433. PMID 28109861
18. Krieg SM, Sabih J, Bulbasova L, et al. Preoperative motor mapping by navigated transcranial magnetic brain stimulation improves outcome for motor eloquent lesions. *Neuro-oncology*. Sep 2014; 16(9): 1274-82. PMID 24516237
19. Krieg SM, Sollmann N, Obermueller T, et al. Changing the clinical course of glioma patients by preoperative motor mapping with navigated transcranial magnetic brain stimulation. *BMC Cancer*. Apr 08 2015; 15: 231. PMID 25884404
20. Frey D, Schilt S, Strack V, et al. Navigated transcranial magnetic stimulation improves the treatment outcome in patients with brain tumors in motor eloquent locations. *Neuro-oncology*. Oct 2014; 16(10): 1365-72. PMID 24923875
21. Picht T, Schulz J, Hanna M, et al. Assessment of the influence of navigated transcranial magnetic stimulation on surgical planning for tumors in or near the motor cortex. *Neurosurgery*. May 2012; 70(5): 1248-56; discussion 1256-7. PMID 22127045

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Navigated Transcranial Magnetic Stimulation

Policy # 00407

Original Effective Date: 03/19/2014

Current Effective Date: 10/12/2020

### **Policy History**

Original Effective Date: 03/19/2014

Current Effective Date: 10/12/2020

- 03/06/2014 Medical Policy Committee review
- 03/19/2014 Medical Policy Implementation Committee approval. New policy.
- 08/03/2015 Coding update: ICD10 Diagnosis code section added; ICD9 Procedure code section removed.
- 08/06/2015 Medical Policy Committee review
- 08/19/2015 Medical Policy Implementation Committee approval. Updated rationale and references. Coverage eligibility unchanged.
- 08/04/2016 Medical Policy Committee review
- 08/17/2016 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
- 01/01/2017 Coding update: Removing ICD-9 Diagnosis Codes
- 08/03/2017 Medical Policy Committee review
- 08/23/2017 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
- 01/01/2018 Coding update
- 08/09/2018 Medical Policy Committee review
- 08/15/2018 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
- 08/01/2019 Medical Policy Committee review
- 08/14/2019 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
- 09/03/2020 Medical Policy Committee review
- 09/09/2020 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

Next Scheduled Review Date: 09/2021

### **Coding**

*The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)‡, copyright 2019 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of*

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Navigated Transcranial Magnetic Stimulation

Policy # 00407

Original Effective Date: 03/19/2014

Current Effective Date: 10/12/2020

*descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.*

*The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.*

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

| Code Type        | Code                  |
|------------------|-----------------------|
| CPT              | 64999                 |
| HCPCS            | No codes              |
| ICD-10 Diagnosis | All related diagnoses |

\*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.





# Louisiana

## Navigated Transcranial Magnetic Stimulation

Policy # 00407

Original Effective Date: 03/19/2014

Current Effective Date: 10/12/2020

- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
1. Consultation with the Blue Cross and Blue Shield Association technology assessment program (TEC) or other nonaffiliated technology evaluation center(s);
  2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
  3. Reference to federal regulations.

‡ Indicated trademarks are the registered trademarks of their respective owners.

**NOTICE:** If the Patient's health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

**NOTICE:** Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.