



# Louisiana

## Serum Biomarker Tests for Multiple Sclerosis

**Policy #** 00433

Original Effective Date: 10/15/2014

Current Effective Date: 03/09/2020

Archived Date: 10/19/2016

Returned to Active Status: 03/09/2020

*Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.*

### Services Are Considered Investigational

*Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.*

Based on review of available data, the Company considers serum biomarker tests for multiple sclerosis (MS) are considered **investigational\*** in all situations.

### Background/Overview

Serum antibodies to polysaccharide-containing molecules, called glycans, and other serum biomarkers are potential biomarker testes for the diagnosis and prognosis of MS. Multiple sclerosis is diagnosed according to criteria that incorporate clinical symptoms and magnetic resonance imaging (MRI) and cerebrospinal fluid findings. Currently, there is no biomarker available to inform diagnosis or prognosis. A serum biomarker is particularly desirable because of ease of repeat measurements.

### Disease Description

Estimated prevalence of MS in North America varies regionally and ranges from 240 of 100,000 in Canada to 191 of 100,000 in Minnesota and 40 of 100,000 in Texas. Women are affected twice as often as men, and median age of onset is 24 years. Most patients (85%) have the relapsing remitting form of MS (RRMS), and of these, 60% to 70% will progress to secondary progressive MS, usually 10 to 30 years after disease onset. Rarer forms are primary progressive MS and progressive relapsing MS.

Multiple sclerosis is characterized by destruction of myelin in the central nervous system.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Serum Biomarker Tests for Multiple Sclerosis

Policy # 00433

Original Effective Date: 10/15/2014

Current Effective Date: 03/09/2020

Archived Date: 10/19/2016

Returned to Active Status: 03/09/2020

symptoms are diverse and may include cognitive, speech, or vision deficits; numbness; pain; weakness or dyscoordination; and bowel or bladder dysfunction. Diagnosis is made by clinical symptoms, typical MRI findings, and oligoclonal antibodies in the cerebrospinal fluid according to current McDonald criteria. Diagnosis requires 2 clinical episodes occurring at 2 discreet points in time, or 1 clinical episode (CIS, defined next) with MRI lesions indicating development at 2 discreet points in time (ie, simultaneous appearance of old and new lesions). Disability progression is quantified in practice and in clinical trials by the Kurtzke Expanded Disability Status Scale (EDSS). Patients with scores less than 5 are fully ambulatory; scores of 5 to 10 are defined by incrementally decreasing ability to walk.

The term clinically isolated syndrome (CIS) describes patients who have suffered a first episode suggestive of MS but do not meet diagnostic criteria for definite MS. Studies indicated that early treatment with interferon beta-1b (IFN $\beta$ -1b) may delay relapse (ie, a second episode), although long-term disability outcomes were unaffected.

In addition to IFN $\beta$ -1b, 8 other disease-modifying drugs are currently U.S. Food and Drug Administration (FDA)-approved for first- or second-line treatment of MS with varying degrees of efficacy for reducing relapses and preventing neurologic deterioration. First-line treatments include self-injectable drugs (interferon and glatiramer acetate) and newer oral agents, such as fingolimod, teriflunomide, and dimethyl fumarate. Choice of first-line agent depends on severity of initial presentation, patient preference, and adverse effect profile. Patients with more active or refractory disease are more likely to tolerate greater risk for greater efficacy, for example with second- or third-line agents, natalizumab and alemtuzumab.

### **Biomarkers**

Glycominds Ltd., based in Israel, markets the diagnostic test, gMS<sup>®†</sup> Dx, for patients with a first episode or CIS, and the multi-marker prognostic test, gMS Pro EDSS, for predicting deterioration in patients diagnosed with MS. Both tests are based on detection of serum antibodies to glycans, which are polysaccharide- or carbohydrate-containing molecules on the surface of immune and other cells. gMS Dx detects immunoglobulin M (IgM) antibodies to the disaccharide glycan, glucose ( $\alpha$ 1,4)glucose( $\alpha$ ) (GAGA4), and gMS Pro EDSS detects IgM antibodies to GAGA2, -3, -4, and -6.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Serum Biomarker Tests for Multiple Sclerosis

Policy # 00433

Original Effective Date: 10/15/2014

Current Effective Date: 03/09/2020

Archived Date: 10/19/2016

Returned to Active Status: 03/09/2020

These anti-glycan antibodies are thought to interfere with normal function of the immune system. Temperature controls are implemented during assay runs to prevent IgM precipitation.

Several other serum biomarkers for MS have been investigated, but no other commercially-available tests were identified.

## **FDA or Other Governmental Regulatory Approval**

### **U.S. Food and Drug Administration (FDA)**

The FDA-approved tests for serum biomarkers in MS are currently unavailable. Glycominds Ltd offered gMS Dx and gMS Pro EDSS as laboratory-developed (in-house) tests at its Clinical Laboratory Improvement Act (CLIA)-certified laboratory in Simi Valley, California. However, current status of the tests is unknown because links to the company website are inactive, and ordering information is not readily available through the parent company, Coronis Partners. Although commercial versions of other biomarker assays were not identified, clinical laboratories may offer in-house assays to measure serum biomarkers in MS.

Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratories offering such tests as a clinical service must meet general regulatory standards of CLIA and must be licensed by CLIA for high-complexity testing.

## **Rationale/Source**

Serum antibodies to polysaccharide-containing molecules, called glycans, and other serum biomarkers are potential biomarker tests for the diagnosis and prognosis of multiple sclerosis (MS). MS is diagnosed according to criteria that incorporate clinical symptoms and magnetic resonance imaging and cerebrospinal fluid findings. Currently, there is no biomarker available to inform diagnosis or prognosis. A serum biomarker is particularly desirable because of ease of repeat measurements.

For individuals who have signs and/or symptoms of MS who receive serum biomarker tests for MS, the evidence includes cross-sectional studies of diagnostic accuracy and cohort studies. Relevant outcomes are test accuracy and validity, other test performance measures, symptoms, functional

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Serum Biomarker Tests for Multiple Sclerosis

Policy # 00433

Original Effective Date: 10/15/2014

Current Effective Date: 03/09/2020

Archived Date: 10/19/2016

Returned to Active Status: 03/09/2020

outcomes, health status measures, and quality of life. Antibodies to glycan molecules are thought to impair immune function. They include antibodies to 1 (glucose[ $\alpha$ 1,4]glucose[ $\alpha$ ] [GAGA4]) or several (GAGA2, -3, -4, and - 6) glycan molecules. The gMS Dx and gMS Pro EDSS tests may aid in the diagnosis and prognosis in MS, respectively. Tests for serum levels of other potential MS biomarkers, including but not limited to apoptosis-related molecules, intercellular adhesion molecules, and myelin peptides, have also been described. Current evidence for these other biomarkers makes it difficult to assess their utility in diagnosis and prognosis. The evidence is insufficient to determine the effects of the technology on health outcomes.

## **Supplemental Information**

### **Practice Guidelines and Position Statements**

#### **Multiple Sclerosis Think Tank**

In 2013, the Multiple Sclerosis Think Tank, a group of approximately 40 hospital neurologists in France, published consensus recommendations for serum tests useful to diagnose MS. Recommendations were developed by systematic review of the literature and a Delphi consensus process. Panelists concurred that “there is currently no useful biological blood test for the positive diagnosis of MS.”

#### **International Advisory Committee on Clinical Trials in Multiple Sclerosis**

In 2014, the International Advisory Committee on Clinical Trials in Multiple Sclerosis, jointly sponsored by the U.S. National Multiple Sclerosis Society, the European Committee for Treatment and Research in Multiple Sclerosis, and the MS Phenotype Group, published results of its deliberations on the MS clinical course descriptions in light of current evidence for improved descriptive terminology (eg, incorporating evidence for serum and other biomarkers). The Committee concluded: “To date, there are no clear clinical, imaging, immunologic, or pathologic criteria to determine the transition point when RRMS [relapse-remitting MS] converts to SPMS [secondary progressive MS]; the transition is usually gradual. This has limited our ability to study the imaging and biomarker characteristics that may distinguish this course.”

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Serum Biomarker Tests for Multiple Sclerosis

Policy # 00433

Original Effective Date: 10/15/2014

Current Effective Date: 03/09/2020

Archived Date: 10/19/2016

Returned to Active Status: 03/09/2020

### U.S. Preventive Services Task Force Recommendations

Not applicable.

### Medicare National Coverage

There is no national coverage determination (NCD). In the absence of an NCD, coverage decisions are left to the discretion of local Medicare carriers.

## References

1. Blue Cross and Blue Shield Association, Medical Policy Reference Manual, “Serum Biomarker Tests for Multiple Sclerosis”, 2.04.118, Archived July 2016.
2. Evans C, Beland SG, Kulaga S et al. Incidence and prevalence of multiple sclerosis in the Americas: a systematic review. *Neuroepidemiology* 2013; 40(3):195-210.
3. Wingerchuk DM, Carter JL. Multiple sclerosis: current and emerging disease-modifying therapies and treatment strategies. *Mayo Clin Proc* 2014; 89(2):225-40.
4. Polman CH, Reingold SC, Banwell B et al. Diagnostic criteria for multiple sclerosis: 2010 revisions to the McDonald criteria. *Ann Neurol* 2011; 69(2):292-302.
5. Kurtzke JF. Rating neurologic impairment in multiple sclerosis: an expanded disability status scale (EDSS). *Neurology* 1983; 33(11):1444-52.
6. Kappos L, Freedman MS, Polman CH et al. Effect of early versus delayed interferon beta-1b treatment on disability after a first clinical event suggestive of multiple sclerosis: a 3-year follow-up analysis of the BENEFIT study. *Lancet* 2007; 370(9585):389-97.
7. Kappos L, Freedman MS, Polman CH et al. Long-term effect of early treatment with interferon beta-1b after a first clinical event suggestive of multiple sclerosis: 5-year active treatment extension of the phase 3 BENEFIT trial. *Lancet Neurol* 2009; 8(11):987-97.
8. Keegan BM. Therapeutic decision making in a new drug era in multiple sclerosis. *Semin Neurol* 2013; 33(1):5-12.
9. Hadjigeorgiou GM, Doxani C, Miligkos M et al. A network meta-analysis of randomized controlled trials for comparing the effectiveness and safety profile of treatments with marketing authorization for relapsing multiple sclerosis. *J Clin Pharm Ther* 2013; 38(6):433-9.
10. Schwarz M, Spector L, Gortler M et al. Serum anti-Glc(alpha1,4)Glc(alpha) antibodies as a biomarker for relapsing-remitting multiple sclerosis. *J Neurol Sci* 2006; 244(1-2):59-68.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Serum Biomarker Tests for Multiple Sclerosis

Policy # 00433

Original Effective Date: 10/15/2014

Current Effective Date: 03/09/2020

Archived Date: 10/19/2016

Returned to Active Status: 03/09/2020

11. Brettschneider J, Jaskowski TD, Tumani H et al. Serum anti-GAGA4 IgM antibodies differentiate relapsing remitting and secondary progressive multiple sclerosis from primary progressive multiple sclerosis and other neurological diseases. *J Neuroimmunol* 2009; 217(1-2):95-101.
12. Freedman MS, Laks J, Dotan N et al. Anti-alpha-glucose-based glycan IgM antibodies predict relapse activity in multiple sclerosis after the first neurological event. *Mult Scler* 2009; 15(4):422-30.
13. Freedman MS, Metzigg C, Kappos L et al. Predictive nature of IgM anti-alpha-glucose serum biomarker for relapse activity and EDSS progression in CIS patients: a BENEFIT study analysis. *Mult Scler* 2012; 18(7):966-73.
14. Polman CH, Reingold SC, Edan G et al. Diagnostic criteria for multiple sclerosis: 2005 revisions to the "McDonald Criteria". *Ann Neurol* 2005; 58(6):840-6.
15. Comabella M, Montalban X. Body fluid biomarkers in multiple sclerosis. *Lancet Neurol* 2014; 13(1):113-26.
16. Ouallet JC, Bodiguel E, Bensa C et al. Recommendations for useful serum testing with suspected multiple sclerosis. *Rev Neurol (Paris)* 2013; 169(1):37-46.
17. Brill L, Goldberg L, Karni A, et al. Increased anti-KIR4.1 antibodies in multiple sclerosis: Could it be a marker of disease relapse? *Mult Scler*. Apr 2015;21(5):572-579. PMID 25392324
18. Findling O, Durot I, Weck A, et al. Antimyelin antibodies as predictors of disability after clinically isolated syndrome. *Int J Neurosci*. Aug 2014;124(8):567-572. PMID 24274327
19. Koudriavtseva T, D'Agosto G, Mandoj C, et al. High frequency of antiphospholipid antibodies in relapse of multiple sclerosis: a possible indicator of inflammatory-thrombotic processes. *Neurol Sci*. Nov 2014;35(11):1737-1741. PMID 24847961
20. Colomba P, Fontana S, Salemi G, et al. Identification of biomarkers in cerebrospinal fluid and serum of multiple sclerosis patients by immunoproteomics approach. *Int J Mol Sci*. 2014;15(12):23269-23282. PMID 25517032
21. Moreno C, Prieto P, Macias A et al. Modulation of voltage-dependent and inward rectifier potassium channels by 15-epi-lipoxin-A4 in activated murine macrophages: implications in innate immunity. *J Immunol* 2013; 191(12):6136-46.
22. Holmoy T, Loken-Amsrud KI, Bakke SJ et al. Inflammation markers in multiple sclerosis: CXCL16 reflects and may also predict disease activity. *PLoS One* 2013; 8(9):e75021.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.





# Louisiana

## Serum Biomarker Tests for Multiple Sclerosis

Policy # 00433

Original Effective Date: 10/15/2014

Current Effective Date: 03/09/2020

Archived Date: 10/19/2016

Returned to Active Status: 03/09/2020

23. Ingram G, Hakobyan S, Hirst CL et al. Complement regulator factor H as a serum biomarker of multiple sclerosis disease state. *Brain* 2010; 133(Pt 6):1602-11.
24. Kvistad S, Myhr KM, Holmoy T, et al. Antibodies to Epstein-Barr virus and MRI disease activity in multiple sclerosis. *Mult Scler*. Dec 2014;20(14):1833-1840. PMID 24842958
25. Sternberg Z, Sternberg D, Drake A, et al. Disease modifying drugs modulate endogenous secretory receptor for advanced glycation end-products, a new biomarker of clinical relapse in multiple sclerosis. *J Neuroimmunol*. Sep 15 2014;274(1-2):197-201. PMID 25064498
26. Gironi M, Solaro C, Meazza C et al. Growth hormone and disease severity in early stage of multiple sclerosis. *Mult Scler Int* 2013; 2013:836486.
27. Ortega-Madueno I, Garcia-Montojo M, Dominguez-Mozo MI, et al. Anti-human herpesvirus 6A/B IgG correlates with relapses and progression in multiple sclerosis. *PLoS One*. 2014;9(8):e104836. PMID 25110949
28. Dimisianos N, Rodi M, Kalavrizioti D, et al. Cytokines as Biomarkers of Treatment Response to IFN beta in Relapsing-Remitting Multiple Sclerosis. *Mult Scler Int*. 2014;2014:436764. PMID 25152817
29. Fissolo N, Canto E, Vidal-Jordana A, et al. Levels of soluble TNF-RII are increased in serum of patients with primary progressive multiple sclerosis. *J Neuroimmunol*. Jun 15 2014;271(1-2):56-59. PMID 24794503
30. Polachini CR, Spanevello RM, Casali EA, et al. Alterations in the cholinesterase and adenosine deaminase activities and inflammation biomarker levels in patients with multiple sclerosis. *Neuroscience*. Apr 25 2014;266:266-274. PMID 24508813
31. Uysal S, Meric Yilmaz F, Bogdaycioglu N, et al. Increased serum levels of some inflammatory markers in patients with multiple sclerosis. *Minerva Med*. Jun 2014;105(3):229-235. PMID 24988088
32. Trenova AG, Slavov GS, Manova MG, et al. Cytokines and disability in interferon-beta-1b treated and untreated women with multiple sclerosis. *Arch Med Res*. Aug 2014;45(6):495-500. PMID 25130430
33. Hartung HP, Reiners K, Archelos JJ et al. Circulating adhesion molecules and tumor necrosis factor receptor in multiple sclerosis: correlation with magnetic resonance imaging. *Ann Neurol* 1995; 38(2):186-93.
34. Trojano M, Avolio C, Simone IL et al. Soluble intercellular adhesion molecule-1 in serum and cerebrospinal fluid of clinically active relapsing-remitting multiple sclerosis: correlation with

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Serum Biomarker Tests for Multiple Sclerosis

Policy # 00433

Original Effective Date: 10/15/2014

Current Effective Date: 03/09/2020

Archived Date: 10/19/2016

Returned to Active Status: 03/09/2020

- Gd-DTPA magnetic resonance imaging-enhancement and cerebrospinal fluid findings. *Neurology* 1996; 47(6):1535-41.
35. Aydin O, Ellidag HY, Eren E, et al. Ischemia modified albumin is an indicator of oxidative stress in multiple sclerosis. *Biochem Med (Zagreb)*. 2014;24(3):383-389. PMID 25351357
  36. Amorini AM, Nociti V, Petzold A, et al. Serum lactate as a novel potential biomarker in multiple sclerosis. *Biochim Biophys Acta*. Jul 2014;1842(7):1137-1143. PMID 24726946
  37. Jafarzadeh A, Ebrahimi HA, Bagherzadeh S, et al. Lower serum levels of Th2-related chemokine CCL22 in women patients with multiple sclerosis: a comparison between patients and healthy women. *Inflammation*. Apr 2014;37(2):604-610. PMID 24254331
  38. Stilund M, Reuschlein AK, Christensen T, et al. Soluble CD163 as a marker of macrophage activity in newly diagnosed patients with multiple sclerosis. *PLoS One*. 2014;9(6):e98588. PMID 24886843
  39. Dickens AM, Larkin JR, Griffin JL, et al. A type 2 biomarker separates relapsing-remitting from secondary progressive multiple sclerosis. *Neurology*. Oct 21 2014;83(17):1492-1499. PMID 25253
  40. Waubant E, Goodkin DE, Gee L et al. Serum MMP-9 and TIMP-1 levels are related to MRI activity in relapsing multiple sclerosis. *Neurology* 1999; 53(7):1397-401.
  41. Trentini A, Manfrinato MC, Castellazzi M, et al. TIMP-1 resistant matrix metalloproteinase-9 is the predominant serum active isoform associated with MRI activity in patients with multiple sclerosis. *Mult Scler*. Feb 6 2015. PMID 25662349
  42. Kacperska MJ, Jastrzebski K, Tomasik B, et al. Selected Extracellular microRNA as Potential Biomarkers of Multiple Sclerosis Activity-Preliminary Study. *J Mol Neurosci*. Dec 10 2014. PMID 25487315
  43. Berger T, Rubner P, Schautzer F et al. Antimyelin antibodies as a predictor of clinically definite multiple sclerosis after a first demyelinating event. *N Engl J Med* 2003; 349(2):139-45.
  44. Kuhle J, Pohl C, Mehling M et al. Lack of association between antimyelin antibodies and progression to multiple sclerosis. *N Engl J Med* 2007; 356(4):371-8.
  45. Koch MW, George S, Wall W, et al. Serum NSE level and disability progression in multiple sclerosis. *J Neurol Sci*. Mar 15 2015;350(1-2):46-50. PMID 25686504
  46. Kivisakk P, Healy BC, Francois K et al. Evaluation of circulating osteopontin levels in an unselected cohort of patients with multiple sclerosis: relevance for biomarker development. *Mult Scler* 2013.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.





# Louisiana

## Serum Biomarker Tests for Multiple Sclerosis

Policy # 00433

Original Effective Date: 10/15/2014

Current Effective Date: 03/09/2020

Archived Date: 10/19/2016

Returned to Active Status: 03/09/2020

47. Shimizu Y, Ota K, Ikeguchi R et al. Plasma osteopontin levels are associated with disease activity in the patients with multiple sclerosis and neuromyelitis optica. J Neuroimmunol 2013; 263(1-2):148-51.
48. Siroos B, Balood M, Zahednasab H et al. Secretory phospholipase A2 activity in serum and cerebrospinal fluid of patients with relapsing-remitting multiple sclerosis. J Neuroimmunol 2013; 262(1-2):125-7.
49. Moccia M, Lanzillo R, Palladino R, et al. Uric acid: a potential biomarker of multiple sclerosis and of its disability. Clin Chem Lab Med. Sep 20 2014. PMID 25241733
50. Lublin FD, Reingold SC, Cohen JA, et al. Defining the clinical course of multiple sclerosis: the 2013 revisions. Neurology. Jul 15 2014;83(3):278-286.

### **Policy History**

Original Effective Date: 10/15/2014

Current Effective Date: 03/09/2020

- |            |  |
|------------|--|
| 10/02/2014 | Medical Policy Committee review  |
| 10/15/2014 | Medical Policy Implementation Committee approval. New policy.  |
| 08/03/2015 | Coding update: ICD10 Diagnosis code section added; ICD9 Procedure code section removed.  |
| 10/08/2015 | Medical Policy Committee review  |
| 10/21/2015 | Medical Policy Implementation Committee approval. Coverage eligibility unchanged.  |
| 10/06/2016 | Medical Policy Committee review. Recommend archiving policy.   |
| 10/19/2016 | Medical Policy Implementation Committee approval. Archived.  |
| 02/06/2020 | Medical Policy Committee review  |
| 02/12/2020 | Medical Policy Implementation Committee approval. Policy reactivated from archive status as investigational to track BCBSA's policy archived in July 2016. |

Next Scheduled Review Date: 02/2021

### **Coding**

*The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)‡, copyright 2019*

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Serum Biomarker Tests for Multiple Sclerosis

Policy # 00433

Original Effective Date: 10/15/2014

Current Effective Date: 03/09/2020

Archived Date: 10/19/2016

Returned to Active Status: 03/09/2020

*by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.*

*The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.*

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	84999
HCPCS	No codes
ICD-10 Diagnosis	G35

\*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



# Louisiana

## Serum Biomarker Tests for Multiple Sclerosis

Policy # 00433

Original Effective Date: 10/15/2014

Current Effective Date: 03/09/2020

Archived Date: 10/19/2016

Returned to Active Status: 03/09/2020

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
  - 1. Consultation with the Blue Cross and Blue Shield Association technology assessment program (TEC) or other nonaffiliated technology evaluation center(s);
  - 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
  - 3. Reference to federal regulations.

‡ Indicated trademarks are the registered trademarks of their respective owners.

**NOTICE:** Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.