



Louisiana

Sphenopalatine Ganglion Block for Headache

Policy # 00563

Original Effective Date: 08/23/2017

Current Effective Date: 09/14/2020

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Note: Botulinum Toxin is addressed separately in medical policy 00012.

Note: Transcutaneous Electrical Nerve Stimulation is addressed separately in medical policy 00142 Electrical Nerve Stimulation Devices.

Note: Occipital Nerve Stimulation is addressed separately in medical policy 00253.

Note: Surgical Deactivation of Headache Trigger Sites is addressed separately in medical policy 00683.

Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Based on review of available data, the Company considers sphenopalatine ganglion (SPG) blocks for all indications, including but not limited to the treatment of migraines and non-migraine headaches to be **investigational**.*

Background/Overview

Headaches AND Headache Treatments

Headaches are common neurologic disorders and are among the top reasons why patients seek medical care. Headaches affect approximately 50% of the general population in a given year and over 90% of people have a lifetime history of headache. The 2 most common types of headache are migraines and tension-type headaches.

Migraines are the second-most common headache disorder, with a 1-year migraine prevalence of approximately 12% in the United States. They are characterized by severe pain on one or both sides of the head, nausea, and, at times, disturbed vision. Migraines can be categorized by headache

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Sphenopalatine Ganglion Block for Headache

Policy # 00563

Original Effective Date: 08/23/2017

Current Effective Date: 09/14/2020

frequency, and by the presence or absence of aura. Chronic migraine is defined as attacks on at least 15 days per month for more than 3 months, with features of migraine on at least 8 days per month.

Tension headaches have a prevalence of approximately 40%. Diagnostic criteria include the presence of at least two of the following characteristics: bilateral headache location, nonpulsating pain, mild-to-moderate intensity, and headache not aggravated by physical activity.

Cluster headaches are less common than tension or migraine headaches, with an estimated prevalence of 0.1% of the population. They are characterized by severe unilateral orbital, supraorbital, and/or temporal pain that also includes other symptoms in the eye and/or nose on the same side (eg, rhinorrhea, eyelid edema or drooping).

Treatment

A variety of medications are used to treat acute migraine episodes. They include medications taken at the onset of an attack to abort the attack (triptans, ergotamines) and medications to treat the pain and other symptoms of migraines once they are established (nonsteroidal anti-inflammatory drugs, antiemetics). Prophylactic medication therapy may be appropriate for people with migraines that occur more than 2 days per week. In addition to medication, behavioral treatments (eg, relaxation, cognitive therapy) are used to manage migraine headache. Botulinum toxin type A injections are a U.S. Food and Drug Administration-approved treatment for chronic migraine.

Severe acute cluster headaches may be treated with abortive therapy, including breathing 100% oxygen, and triptan medications. Other medications used to treat cluster headaches include steroids, calcium channel blockers, and nerve pain medications. Due to the severity of pain associated with cluster headaches, patients may seek emergency treatment. Tension-type headaches are generally treated with over the counter pain medication.

Sphenopalatine Ganglion Block

Sphenopalatine ganglion (SPG) blocks are a proposed treatment option for chronic migraines and some severe non-migraine headaches. The SPG is a group of nerve cells located behind the bony structures of the nose. The nerve bundle is linked to the trigeminal nerve, the primary nerve involved in headache disorders. The SPG has both autonomic nerves, which in this case are associated with functions such as tearing and nasal congestion, and sensory nerves, associated with pain perception. SPG blocks involve topical application of local anesthetic to mucosa overlying the SPG. The

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Sphenopalatine Ganglion Block for Headache

Policy # 00563

Original Effective Date: 08/23/2017

Current Effective Date: 09/14/2020

rationale for using SPG blocks to treat headaches is that local anesthetics in low concentrations could block the sensory fibers and thereby reduce pain while maintaining autonomic function.

The proposed procedure for SPG blockade is to insert intranasally a catheter that is attached to a syringe carrying local anesthetic (eg, lidocaine, bupivacaine). Once the catheter is in place, the local anesthetic is applied to the posterior wall of the nasal cavity and reaches the SPG. Originally, SPG blocks were done by inserting a cotton-tipped applicator dabbed with local anesthetic into the nose; this technique may be less accurate and effective than the currently proposed procedure. Neurostimulation of the SPG and SPG blockade with radiofrequency lesioning have been used outside of the United States, but these treatments are not cleared or approved by the U.S. Food and Drug Administration.

Three catheter devices are commercially available in the United States for performing SPG blocks. The catheters have somewhat different designs but all are attached to syringes to deliver local anesthetic. The catheters are inserted intranasally and, once in place, the local anesthetic is applied through the catheter. With 2 of the 3 commercially available catheters (the SpenoCath[®], Allevio[™])[‡], patients are positioned on their back with their nose pointed vertically and their head turned to the side. With the Tx360[®][‡] device, patients remain seated.

The company marketing the Tx360 device proposes its use in the context of the MiRx[™][‡] protocol. This 2-part protocol includes a medical component for immediate pain relief and a physical component to reduce headache recurrences. The medical component involves clinical evaluation and, if the patient is considered eligible, an SPG block procedure. The physical component can include any of a number of approaches such as physical therapy, ergonomic modifications, massage, and dietary recommendations.

The optimal number and frequency of SPG treatments is unclear. Information from the American Migraine Foundation suggests that the procedure can be repeated as often as needed to control pain. A randomized controlled trial has described a course of treatment for migraines consisting of SPG blocks twice a week for 6 weeks (total, 12 treatments).

SGB blocks are proposed for both short- and long-term treatment of headaches and migraines. When used in the emergency setting in patients with severe acute headaches, the goal of treatment is to abort the current headache while the patient is in the emergency department. In the randomized

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Sphenopalatine Ganglion Block for Headache

Policy # 00563

Original Effective Date: 08/23/2017

Current Effective Date: 09/14/2020

controlled trial that provided a 6-week course of treatment with SPG blocks for chronic migraine (mentioned above), short-term outcomes were assessed up to 24 hours after each treatment, and the duration and frequency of chronic migraines were assessed at 1 and 6 months after the course of treatment.

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

The Tx360 Nasal Applicator (Tian Medical), the Allevio SPG Nerve Block Catheter (JET Medical), and the SpenoCath (Dolor Technologies) are considered class I devices by the Food and Drug Administration and are exempt from 510(k) requirements. This classification does not require submission of clinical data on efficacy but only notification of Food and Drug Administration prior to marketing. All 3 devices are used to apply numbing medication intranasally.

Rationale/Source

Chronic migraine and severe headaches are common conditions and the available treatments are not universally effective. A proposed treatment option is blocking the sphenopalatine ganglion (SPG) nerve by applying topical anesthetic intranasally. Several catheters approved by the U.S. Food and Drug Administration are available for the SPG blocking procedure.

For individuals who have chronic migraine who receive SPG block(s), the evidence includes a randomized controlled trial (RCT) and a case report. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. The randomized trial evaluated a regimen of 12 SPG blocks over 6 weeks and was double-blind and placebo-controlled. The trial found significantly greater short-term (up to 24 hours) benefits from active treatment than from placebo. There were no significant long-term effects (ie, 1 and 6 months after 12 treatments), although the trial was underpowered to detect longer term efficacy. Given that SPG blocks are being proposed as a preventive therapy for chronic migraines, evidence demonstrating reduced migraine frequency, severity, or other objective outcomes from robust trials is still needed. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have severe acute headache treated in the emergency setting who receive SPG block(s), the evidence includes 1 RCT. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. The randomized, double-blind, placebo-controlled

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Sphenopalatine Ganglion Block for Headache

Policy # 00563

Original Effective Date: 08/23/2017

Current Effective Date: 09/14/2020

trial evaluated a single SPG block for severe acute headache of mixed etiologies. There was no statistically significant difference between active treatment and placebo for the primary outcome (pain reduction 15 minutes postintervention). The trialists did not collect pain data again until 24 hours posttreatment, at which time significantly more patients were headache-free in the active treatment arm than in the placebo arm. Additional studies, preferably RCTs, are needed to determine whether SPG blocks are an effective treatment in the emergency setting. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have cluster headache who receive SPG block(s), the evidence includes case series. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Two small case series, both of which evaluated an approach for intranasal SPG blocks that differs from the intervention currently available in the United States, were identified. In these series, 40% to 50% of patients experienced complete symptom relief for a variable length of time and about 20% had treatment-related complications. However, it is not clear from these series the degree to which the procedures evaluated differ in safety and efficacy from an intranasal SPG block using a device cleared by the U.S. Food and Drug Administration. Additional studies, preferably RCTs, are needed to evaluate SPG blocks for treating cluster headaches. The evidence is insufficient to determine the effects of the technology on health outcomes.

Supplemental Information

Practice Guidelines and Position Statements

No guidelines or statements were identified.

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 1.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Sphenopalatine Ganglion Block for Headache

Policy # 00563

Original Effective Date: 08/23/2017

Current Effective Date: 09/14/2020

Table 1. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT02090998	Sphenopalatine Ganglion Nerve Block vs. Elavil for Treatment of Transformed Migraines	200	Jul 2021
<i>Unpublished</i>			
NCT02365909	Study Evaluating Sphenopalatine Ganglion Block (SPGB) for Treatment of Postdural Puncture Headache (PDPH)	6	Aug 2016 (terminated)

NCT: national clinical trial.

References

1. Blue Cross and Blue Shield Association, Medical Policy Reference Manual, “Sphenopalatine Ganglion Block for Headache”, Policy 7.01.159, December 2019.
2. International Association for the Study of Pain (IASP). Epidemiology of Headache. 2011; <https://s3.amazonaws.com/rdcms-iasp/files/production/public/Content/ContentFolders/GlobalYearAgainstPain2/HeadacheFactSheets/1-Epidemiology.pdf>.
3. Singh A, Soares WE. Management strategies for acute headache in the emergency department. *Emerg Med Pract.* Jun 2012;14(6):1-23; quiz 23-24. PMID 22830180
4. Headache Classification Committee of the International Headache S. The International Classification of Headache Disorders, 3rd edition (beta version). *Cephalalgia.* Jul 2013;33(9):629-808. PMID 23771276
5. Sanders M, Zuurmond WW. Efficacy of sphenopalatine ganglion blockade in 66 patients suffering from cluster headache: a 12- to 70-month follow-up evaluation. *J Neurosurg.* Dec 1997;87(6):876-880. PMID 9384398
6. American Migraine Foundation. Sphenopalatine Ganglion Blocks in Headache Disorders. 2016; <https://americanmigrainefoundation.org/understanding-migraine/sphenopalatine-ganglion-blocks-in-headache-disorders/>.
7. The MiRx™ Protocol. 2013; <http://www.mirxprotocol.com/>.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Sphenopalatine Ganglion Block for Headache

Policy # 00563

Original Effective Date: 08/23/2017

Current Effective Date: 09/14/2020

8. Cady R, Saper J, Dexter K, et al. A double-blind, placebo-controlled study of repetitive transnasal sphenopalatine ganglion blockade with tx360((R)) as acute treatment for chronic migraine. *Headache*. Jan 2015;55(1):101-116. PMID 25338927
9. Cady RK, Saper J, Dexter K, et al. Long-term efficacy of a double-blind, placebo-controlled, randomized study for repetitive sphenopalatine blockade with bupivacaine vs. saline with the Tx360 device for treatment of chronic migraine. *Headache*. Apr 2015;55(4):529-542. PMID 25828648
10. Candido KD, Massey ST, Sauer R, et al. A novel revision to the classical transnasal topical sphenopalatine ganglion block for the treatment of headache and facial pain. *Pain Physician*. Nov-Dec 2013;16(6):E769-778. PMID 24284858
11. Headache Classification Subcommittee of the International Headache S. The International Classification of Headache Disorders: 2nd edition. *Cephalalgia*. 2004;24(Suppl 1):9-160. PMID 14979299
12. Schaffer JT, Hunter BR, Ball KM, et al. Noninvasive sphenopalatine ganglion block for acute headache in the emergency department: a randomized placebo-controlled trial. *Ann Emerg Med*. May 2015;65(5):503-510. PMID 25577713
13. Pipolo C, Bussone G, Leone M, et al. Sphenopalatine endoscopic ganglion block in cluster headache: a reevaluation of the procedure after 5 years. *Neurol Sci*. Jun 2010;31(Suppl 1):S197-199. PMID 20464621
14. Felisati G, Arnone F, Lozza P, et al. Sphenopalatine endoscopic ganglion block: a revision of a traditional technique for cluster headache. *Laryngoscope*. Aug 2006;116(8):1447-1450. PMID 16885751

Policy History

Original Effective Date: 08/23/2017

Current Effective Date: 09/14/2020

08/03/2017 Medical Policy Committee review

08/23/2017 Medical Policy Implementation Committee approval. New policy.

08/09/2018 Medical Policy Committee review

08/15/2018 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

08/01/2019 Medical Policy Committee review

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Sphenopalatine Ganglion Block for Headache

Policy # 00563

Original Effective Date: 08/23/2017

Current Effective Date: 09/14/2020

08/14/2019 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

01/07/2020 Coding update

08/06/2020 Medical Policy Committee review

08/12/2020 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

09/16/2020 Coding update

Next Scheduled Review Date: 08/2021

Coding

The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)‡, copyright 2019 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Sphenopalatine Ganglion Block for Headache

Policy # 00563

Original Effective Date: 08/23/2017

Current Effective Date: 09/14/2020

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	64505, 64999
HCPCS	No codes
ICD-10 Diagnosis	G43.001-G43.919, G44.001-G44.029, G44.031-G44.89 Deleted the following code eff 10/1/2020: R51 Added the following codes eff 10/1/2020: R51.0, R51.9

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
 - 1. Consultation with the Blue Cross and Blue Shield Association technology assessment program (TEC) or other nonaffiliated technology evaluation center(s);
 - 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
 - 3. Reference to federal regulations.

‡ Indicated trademarks are the registered trademarks of their respective owners.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Sphenopalatine Ganglion Block for Headache

Policy # 00563

Original Effective Date: 08/23/2017

Current Effective Date: 09/14/2020

NOTICE: If the Patient's health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

©2020 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.